

Isolation, Identification, and Antibiotic Susceptibility Testing of *Salmonella* Isolated from Foodborne Outbreaks



Mohammad Mehdi Soltan Dallal^{1,2*}, Milad Abdi^{3,4}, Mahya Khalilian¹, Zahra Rajabi², Ronak Bakhtiari¹, Mohammad Kazem Sharifi Yazdi⁵, Somayeh Yaslianifard⁶, Seyedeh Masoumeh Abrishamchian Langroudi⁷

¹Division of Microbiology, Department of Pathobiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

²Food Microbiology Research Center, Tehran University of Medical Sciences, Tehran, Iran

³Student Research Committee, Faculty of Medicine, Iran University of Medical Sciences, Tehran, Iran

⁴Department of Microbiology, Faculty of Medicine, Iran University of Medical Sciences, Tehran, Iran

⁵Zoonosis Research Center, Tehran University of Medical Sciences, Tehran, Iran

⁶Department of Microbiology, Medical School, Alborz University of Medical Sciences, Karaj, Iran

⁷Pathobiology Laboratory Center, No17, Gharani Alley, Valiasr Ave, Tehran, Iran

*Corresponding Author:

Mohammad Mehdi Soltan Dallal,
Division of Food Microbiology,
Department of Pathobiology,
School of Public Health, Tehran
University of Medical Sciences,
Tehran, Iran.
Tel: +989121452646;
Email: msoltandallal@gmail.com

Published Online August 30, 2020

Keywords: *Salmonella*,
Foodborne diseases,
Outbreak, Diarrhea, Antibiotic
susceptibility testing



Abstract

Background: Foodborne diseases are a major problem worldwide. The epidemiological investigations in many parts of the world have shown an increase in infections caused by *Salmonella* serovars. Furthermore, the emergence of drug resistance among them has become a major global concern-and awareness of the resistance patterns of *Salmonella* could be very useful in treatment of diseases.

Objective: This study aimed to investigate *Salmonella* serotypes in foodborne outbreaks by sequencing of ITS region of 16S-23SrRNA gene and to determine their antimicrobial susceptibility pattern.

Materials and Methods: A total of 614 diarrheal stool samples were collected from 173 foodborne outbreaks in different provinces of Iran during one year. Identification of *Salmonella* was carried out by phenotypic and molecular (16s-23srRNA gene detection) methods and antibiotic susceptibility was performed using disc diffusion method.

Results: Out of 614 samples, 18 isolates were identified as *Salmonella* of which 16 (88.9%) isolates were *Salmonella* Enteritidis and 2 (11.1%) isolates as *Salmonella* Paratyphi A. All isolates were sensitive to ceftazidime, and high resistance was seen with nalidixic acid with 14 (77.8%) isolates.

Conclusion: Increasing antibiotic resistance in many bacterial pathogens such as *Salmonella* has been a major threat for human health. Therefore, identifying the antibiotic resistance patterns of *Salmonella* serovars may help in treatment of the associated infections.

Received February 19, 2020; Revised May 15, 2020; Accepted August 15, 2020

Background

Today, antibiotic resistance is growing rapidly; the excessive or inappropriate use of antibiotics led to the emergence of resistant strains of *Salmonella*.^{1,2} Extending resistance to antimicrobials may be created through direct or indirect consumption of antibiotic-containing foods. Animal foods can act as reservoirs of resistance genes. The prohibition of unnecessary use of antibiotics not only helps to prevent the spread of resistant strains but also prevents losing therapeutic effect of them.³ Hence, the community health systems should also consider these issues in animal husbandry and agriculture.¹ However, with increasing awareness, more countries have banned the misuse of antibiotics, but the resistance of pathogens has not diminished so far, and now the main concern for

food safety is inhibitory activity against such resistant bacteria.⁴ *Salmonella* is a gram-negative rod-shaped bacterium in the *Enterobacteriaceae* family with more than 2500 serovars. This bacterium causes salmonellosis, one of the most common food-borne diseases transmitted to human by contaminated foods such as poultry, beef, pork, egg, milk, cheese, seafood, fruits, juice, and vegetables.⁵ It is estimated that about 94 million cases of gastroenteritis and 155 000 deaths occur because of *Salmonella* each year worldwide, of which 85% are food related (6). Nontyphoid *Salmonella* spp., such as *S. enterica* serovar Typhimurium and *S. enterica* serovar Enteritidis, are important agents of food-borne outbreaks worldwide.⁷ Unfortunately, the outbreak of *Salmonella* even in developed countries has increased recently. In addition, the emergence of

antibiotic resistance of *S. enterica* serovars (Typhimurium, Enteritidis, and Paratyphi A) to common antibiotics and their spread has been an issue for the health system.⁸ The overuse of antibiotics in animals for prevention or treatment of diseases and growth promotion has increased the antibiotic resistance in zoonotic food-borne pathogens such as *Salmonella*.⁹ Of particular concern is development of resistance to effectual drugs such as fluoroquinolones and β -lactams. Today, the number of *Salmonella* isolates with high level of resistance to several antibiotics (multiple drug resistance [MDR]) is growing. MDR *Salmonella* may be transmitted to human throughout the production chain and fecal-oral route.^{8,10} This study aimed to isolate and determine the *Salmonella* serotypes in foodborne outbreaks by evaluating phenotypic characteristics and sequencing of ITS region of 16S-23S rRNA gene followed by the antimicrobial susceptibility testing (AST).

Materials and Methods

Sample Collection, Isolation, and Phenotypic Identification

In this descriptive cross-sectional study, from October 2013 to September 2014, a total of 614 diarrheal stool samples were collected from 173 foodborne outbreaks from different provinces of Iran. Exclusion criteria were history of gastrointestinal disease in the past one month, diarrhea after medication, and viral diarrhea. According to the standard methods, stool samples were transferred to the Kerry Blair transfer medium (Merck, Germany), then transferred to the Selenite F (Merck, Germany) and cultured on Hektoen Enteric Agar medium (Merck, Germany) for 8 to 12 hours in the microbiology laboratory of the Department of Pathobiology of Tehran University of Medical Sciences, Tehran, Iran. After incubation for 24 hours at 37°C, the black colonies suspected to *Salmonella* were chosen and then tested for phenotypic characteristics and biochemical tests such as indole, lactose, H₂S, lysine, citrate, Methyl Red-Voges Proskauer (MR-VP), Urea, and motility.¹¹

Antimicrobial Susceptibility Testing

AST was performed using disc diffusion method on Mueller-Hinton agar as recommended by the Clinical and Laboratory Standards Institute (CLSI).⁶ The used antibiotic discs (MAST, UK) for AST were amoxicillin (20 μ g), chloramphenicol (30 μ g), tetracycline (30 μ g), ampicillin (10 μ g), trimethoprim-sulfamethoxazole (23.75/1.25 μ g), nalidixic acid (30 μ g), ciprofloxacin (5 μ g), cefotaxime (30 μ g), ceftazidime (30 μ g), and meropenem (10 μ g). After incubation for 18 hours, based on the diameter of created inhibition zone around the antibiotic discs, *Salmonella* isolates were considered as sensitive (S) or intermediate (I) or resistant (R).¹²

Genotypic Identification

Identification and determination of the *Salmonella* strains was performed using sequencing of the 16S-23S rRNA gene internal transcribed spacer region. The primer sequences were as follows: reverse primer 5'-TATAGCCCCATCGTGTAGTCAGAAC-3' and forward primer 5'-TGC GGC TGG ATC ACC TCC TT-3'.¹³ Genomic DNA of the *Salmonella* isolates was extracted using boiling method from purified cultures.¹⁴ In short, using sterile loop, a few colonies of bacteria were suspended in 50 μ L of Sodium Chloride-Tris-EDTA (STE) Buffer in 1.5 mL microtube and then boiled for 10 minutes. After centrifugation at 13300 rpm for 3 minutes, the supernatant was removed and finally concentration of DNA was measured by NanoDrop 3300. The polymerase chain reaction (PCR) reaction was carried out in a total volume of 20 μ L containing 15.1 μ L of sterile distilled water, 2 μ L of PCR-buffer (10 \times), 0.6 μ L of MgCl₂ (50 mM) 0.4 μ L of dntp-mix (10 mM), 0.3 μ L of Primer F, 0.3 μ L of Primer R, 0.3 μ L of Taq polymerase (5 u/ μ L) (Pishgam Co., Iran), and 2 μ L of DNA. The PCR temperature conditions included initial denaturation at 94°C for 2 minutes, followed by 35 cycles of denaturation at 94°C for 30 seconds, annealing at 71°C for 30 seconds and extension at 72°C for 50 seconds. Finally, the final extension was allowed at 72°C for 10 minutes. After amplification, to analyze the PCR products, electrophoresis was done using a 1% agarose gel containing safe stain and a TBE 0.5X for electrophoresis buffer, as well as a 100-bp ladder and a voltage of 80-100 V.

Results

Phenotypic Identification and Antimicrobial Susceptibility Testing

Out of 614 samples, 18 (2.93%) isolates were identified as *Salmonella*. The most abundant serotype was *Salmonella* Enteritidis with a frequency of 16 (88.89%), while the remaining 2 (11.11%) isolates were *Salmonella* Paratyphi A.

Of the total 18 isolates of *Salmonella* collected from different provinces, Alborz province with 5 (27.78%) isolates and Mazandaran with 4 (22.22%) isolates had the highest rate of *Salmonella* outbreak. Moreover, 15 (83.33%) isolates belonged to the samples collected in the spring and summer seasons. Women were the major origin of *Salmonella* with the frequency of 15 (83.33%) isolates, while only 3 (16.67%) isolates were obtained from men. The dominant age groups were 7-12-year-old and 25-35-year-old subjects, each with 4 (22.22%) isolates.

According to the results of AST, the highest rate of resistance was related to nalidixic acid (14 (77.77%)) and the highest susceptibility was seen in ceftazidime (18 (100%)), followed by amoxicillin and chloramphenicol (17 (94.44%)) (Table 1).

Table 1. Antimicrobial Susceptibility Pattern of 18 *Salmonella* Isolates

Antibiotic	Abbreviation	Sensitive No. (%)	Intermediate No. (%)	Resistant No. (%)
meropenem	MEM	2 (11.11)	13 (72.22)	3 (16.66)
Cefotaxime	CTX	14 (77.77)	4 (22.22)	0 (0)
Chloramphenicol	C	17 (94.4)	1 (5.55)	0 (0)
Co-trimoxazole	TS	13 (72.22)	1 (5.55)	4 (22.22)
Ciprofloxacin	CIP	16 (88.88)	2 (11.11)	0 (0)
Nalidixic Acid	NA	3 (16.66)	1 (5.55)	14 (77.77)
Amoxicillin	A	17 (94.4)	0 (0)	1 (5.55%)
Tetracycline	T	14 (77.77)	0 (0)	4 (22/22)
Ampicillin	AP	14 (77.77)	4 (22.22)	0 (0)
Ceftazidime	CAZ	18 (100)	0 (0)	0 (0)

Genotypic Identification

After amplification of target gene and electrophoresis of the PCR products, 312 kb bands were observed in gel, which confirmed *Salmonella* genus (Figure 1). After confirmation, PCR products were sent to Takapoozist company to sequence it for determination of the serotypes of *Salmonella*. After sequencing and alignment by BLAST (NCBI tool), it was found that 16 isolates belonged to *Salmonella* Enteritidis and two isolates belonged to *Salmonella* Paratyphi A.

Discussion

The emergence of foodborne pathogens, along with their antibiotic resistance and effects on the general health of the community is one of the most important challenges to the healthcare system.¹⁵ Kozak et al showed that Canada, with the highest consumption of vegetables and fresh fruits, had the highest rates for the most common causes of food-borne outbreak and *Salmonella* had the highest morbidity rate (about 50%).¹⁶ This study showed that vegetables and fruits were the most frequent sources of *Salmonella* isolates causing foodborne outbreak. Of the

total 18 isolates, 6 (33%) isolates belonged to the patients with consumption of fruits and vegetables, and 5 (27%) isolates were related to vegetables and meat, which are similar to the study by Kozak et al.

A study in Turkey on the antimicrobial resistance of non-typhoid *Salmonella* in 2000-2002 showed that there is a decrease in sensitivity to ciprofloxacin especially in *Salmonella* C-type, which is slightly different with the results of our research because isolated serotypes in our study were all sensitive to ciprofloxacin, except for 2 (11%) cases that were semi-sensitive.¹⁷ In a study conducted in Palestine between 1997 and 2002, 53.3% non-typhoid *Salmonella* was isolated, and the highest resistance rate was related to nalidixic acid (79%), which was similar to our study (77%).¹⁸ In a study by Hur et al in Korea in 2011, 46 isolates of *Salmonella* obtained from different sources were investigated to detect antibiotic resistance genes, and the highest resistance rate was related to nalidixic acid and sulfamethoxazole. Of the 46 isolates, 21 isolates were resistant to ampicillin, piperacillin, and ticarcillin, of which 19 isolates had the *bla*-TEM gene and only one had the *bla*-CTX-M gene.¹⁹ In our study, only 5 (27%) and 1 (5%) isolates of all isolates were semi-sensitive to ampicillin and chloramphenicol, respectively; in addition, no resistance was observed in the mentioned antibiotics.

Although the antibiotic resistance of *Salmonella* enteritidis is significantly lower than *Salmonella* typhimurium,²⁰ this is important since *Salmonella* enteritidis is the most important cause of salmonellosis in food-borne outbreaks.^{15,21,22} Therefore, it is anticipated that the information obtained from this study could be helpful in detection of *Salmonella* and its antibiotic resistance pattern in order to reduce the therapeutic costs and take the necessary control and prevention measurements.

Authors' Contributions

MMSD, Conceived and designed the experiments; MK, ZR, carried out all the experiments; MA, writing the first draft of manuscript; SY, RB, contributed in literature survey and help in writing the manuscript; SMAL, designed preparation of experiments; MKSY, writing the final draft of manuscript.




Figure 1. PCR Agarose Gel Electrophoresis Results for Detection of the 16S-23S rRNA Gene in *Salmonella* Isolates (Product Size = 312 bp). 1, 2, and 3: *Salmonella* Isolates. 4: Negative Control.

Ethical Approval

Not applicable.

Conflict of Interest Disclosures

The authors declare that they have no conflict of interests.

Financial Support

The present study is financially supported by Tehran University of Medical Sciences, Tehran, Iran.

Acknowledgments

This study was part of a research project approved by the Food Microbiology Research Center, Tehran University of Medical Sciences, Tehran, Iran (contract no: 23115).

References

1. Kalily E, Hollander A, Korin B, Cymerman I, Yaron S. Adaptation of *Salmonella enterica* serovar Senftenberg to linalool and its association with antibiotic resistance and environmental persistence. *Appl Environ Microbiol.* 2017;83(10):e03398-16. doi:10.1128/aem.03398-16
2. Kisluk G, Kalily E, Yaron S. Resistance to essential oils affects survival of *Salmonella enterica* serovars in growing and harvested basil. *Environ Microbiol.* 2013;15(10):2787-2798. doi:10.1111/1462-2920.12139
3. Capalonga R, Ramos RC, Both JM, et al. *Salmonella* serotypes, resistance patterns, and food vehicles of salmonellosis in southern Brazil between 2007 and 2012. *J Infect Dev Ctries.* 2014;8(7):811-817. doi:10.3855/jidc.3791
4. Koluman A, Dikici A. Antimicrobial resistance of emerging foodborne pathogens: status quo and global trends. *Crit Rev Microbiol.* 2013;39(1):57-69. doi:10.3109/1040841x.2012.691458
5. Zaghoul M, Al-Ebshahy E, Khaliel S, Ellakany H. Molecular identification of beta-lactamase-producing *Salmonella enteritidis* from broiler chickens. *Alex J Vet Sci.* 2017;55(1):204-209. doi:10.5455/ajvs.246419
6. Yang X, Huang J, Wu Q, et al. Prevalence, antimicrobial resistance and genetic diversity of *Salmonella* isolated from retail ready-to-eat foods in China. *Food Control.* 2016;60:50-56. doi:10.1016/j.foodcont.2015.07.019
7. Nakaya H, Yasuhara A, Yoshimura K, Oshihoi Y, Izumiya H, Watanabe H. Life-threatening infantile diarrhea from fluoroquinolone-resistant *Salmonella enterica* typhimurium with mutations in both gyrA and parC. *Emerg Infect Dis.* 2003;9(2):255-257. doi:10.3201/eid0902.020185
8. Afshari A, Baratpour A, Khanzade S, Jamshidi A. *Salmonella enteritidis* and *Salmonella typhimurium* identification in poultry carcasses. *Iran J Microbiol.* 2018;10(1):45-50.
9. Harichandran D, Dinesh KR. Antimicrobial susceptibility profile, treatment outcome and serotype distribution of clinical isolates of *Salmonella enterica* subspecies enterica: a 2-year study from Kerala, South India. *Infect Drug Resist.* 2017;10:97-101. doi:10.2147/idr.s126209
10. Fey PD, Safranek TJ, Rupp ME, et al. Ceftriaxone-resistant salmonella infection acquired by a child from cattle. *N Engl J Med.* 2000;342(17):1242-1249. doi:10.1056/nejm200004273421703
11. Ge Z, Xue S, Jianmei Z, et al. Isolation, identification, and characterization of foodborne pathogens isolated from egg internal contents in China. *J Food Prot.* 2016;79(12):2107-2112. doi:10.4315/0362-028x.jfp-16-168
12. Cockerill F, Patel J, Alder J, Bradford P, Dudley M, Eliopoulos G. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Third Informational Supplement (M100-S23). Wayne, PA: CLSI; 2013.
13. Chiu TH, Chen TR, Hwang WZ, Tsen HY. Sequencing of an internal transcribed spacer region of 16S-23S rRNA gene and designing of PCR primers for the detection of *Salmonella* spp. in food. *Int J Food Microbiol.* 2005;97(3):259-265. doi:10.1016/j.ijfoodmicro.2004.04.005
14. Agersborg A, Dahl R, Martinez I. Sample preparation and DNA extraction procedures for polymerase chain reaction identification of *Listeria monocytogenes* in seafoods. *Int J Food Microbiol.* 1997;35(3):275-280. doi:10.1016/s0168-1605(97)01245-2
15. Fardsanei F, Soltan Dallal MM, Douraghi M, et al. Genetic diversity and virulence genes of *Salmonella enterica* subspecies enterica serotype enteritidis isolated from meats and eggs. *Microb Pathog.* 2017;107:451-456. doi:10.1016/j.micpath.2017.04.026
16. Kozak GK, MacDonald D, Landry L, Farber JM. Foodborne outbreaks in Canada linked to produce: 2001 through 2009. *J Food Prot.* 2013;76(1):173-183. doi:10.4315/0362-028x.jfp-12-126
17. Erdem B, Ercis S, Hascelik G, Gur D, Aysev AD. Antimicrobial resistance of *Salmonella enterica* group C strains isolated from humans in Turkey, 2000-2002. *Int J Antimicrob Agents.* 2005;26(1):33-37. doi:10.1016/j.ijantimicag.2005.03.007
18. Weinberger M, Solnik-Isaac H, Shachar D, et al. *Salmonella enterica* serotype Virchow: epidemiology, resistance patterns and molecular characterisation of an invasive *Salmonella* serotype in Israel. *Clin Microbiol Infect.* 2006;12(10):999-1005. doi:10.1111/j.1469-0691.2006.01466.x
19. Hur J, Kim JH, Park JH, Lee YJ, Lee JH. Molecular and virulence characteristics of multi-drug resistant *Salmonella enteritidis* strains isolated from poultry. *Vet J.* 2011;189(3):306-311. doi:10.1016/j.tvjl.2010.07.017
20. Ahmed AM, Younis EE, Ishida Y, Shimamoto T. Genetic basis of multidrug resistance in *Salmonella enterica* serovars enteritidis and Typhimurium isolated from diarrheic calves in Egypt. *Acta Trop.* 2009;111(2):144-149. doi:10.1016/j.actatropica.2009.04.004
21. Fardsanei F, Nikkhahi F, Bakhshi B, Salehi TZ, Tamai IA, Soltan Dallal MM. Molecular characterization of *Salmonella enterica* serotype enteritidis isolates from food and human samples by serotyping, antimicrobial resistance, plasmid profiling, (GTG)5-PCR and ERIC-PCR. *New Microbes New Infect.* 2016;14:24-30. doi:10.1016/j.nmni.2016.07.016
22. Soltan Dallal MM, Khalilian M, Masoumi Asl H, et al. Molecular epidemiology and antimicrobial resistance of *Salmonella* spp. Isolated from resident patients in Mazandaran province, Northern Iran. *J Food Qual Hazards Control.* 2016;3(4):146-151.