ijer.skums.ac.ir

The prevalence and related factors of low birth weight

Marziye Safari¹, Azam Samiee¹, Fatemeh Salehi², SeyedeNafiseh Ahmadi³, SeyedeSoghra Ahmadi^{1*}

¹Student Research Committee, Semnan University of Medical Sciences, Semnan, I.R. Iran;²Epidemiology and Biostatistics Dept., Tehran University of Medical Sciences, Tehran, I.R. Iran; ³Public Health Dept.,Shahrekord University of Medical Sciences, Shahrekord, I.R. Iran. Received: 3/Oct/2015 Accepted: 28/Nov/2015

ABSTRACT

Background and aims: The most common reason in neonatal mortality rate in the world is low birth weight (LBW). The objective of our study was to determine the prevalence and related factors associated in birth weight in Garmsar, Iran in 2013.

Methods: This cross sectional study was carried out on 683 live births which delivered in Garmsar in 2013. The required information was collected by examining the health records of pregnant women and completion of the data registration forms. Data collection was controlled by using SPSSand analyzed by using an Independent T-test, Chi-square test and logistic regression.

Results: According to the results, 32 infants out of 683 infants born in 2013 had low birth weight, i.e. the weight less than 2500g. LBW prevalence estimated in this study was 4.7%. There were significant statistical relations between low birth weight and the variables including number of births, mothers' occupation and interval of less than 3 years between pregnancies and systolic and diastolic blood pressures.

Conclusion: Although our results regarding LBW prevalence and risk factors confirm the results of other studies, it was recommended perspective studies to reconfirm the LBW risk factors.

Keywords: Birth Weight, Low Birth Weight, Prevalence, Related factors, Garmsar.

INTRODUCTION

Infant mortality is one of the serious worries in the developing and the developed countries. Despite the fact that infant mortality rate (IMR) is gradually reducing in the world, changing neonatal mortality rate (NMP) is much slower. The common reasons in neonatal mortality rate in the world include preterm childbirth and low birth weight (LBW).¹⁻⁴ Normal weight on birth is considered 2500-3999g, lower than which is considered as low weight, and higher than that is high birth weight or macrosomia.⁵

Low birth weight is related to increasing risks of cardiac ischemic diseases, hypertension and diabetes in the later ages in life.⁶⁻¹⁰ LBW

*Corresponding author: Seyede Soghra Ahmadi, Public Health Dept., Semnan University of Medical Sciences, Semnan, I.R. Iran, Tel: 00989136743820, E-mail:ahmadi2744@yahoo.com

prevalence is different in the developing countries (5-7%) and the developed countries (19%).¹¹ According to a reviewing study, LBW prevalence in Iran is 7%, showing an increasing trend from 1991 to 2010.¹²

Birth weight in infants is affected by different factors, including maternal and environmental factors.^{13,14} By identifying and controlling the risk factors that are mainly depending on living and social conditions, the low birth weight of infants could be prevented.^{15,16} According to various evidences, maternal factors, including genetic, social/cultural elements, population factors, medical and behavioral conditions of mothers, are effective on the birth weight, both directly and indirectly.^{13,17}

Importance of the above factors and incompatibility of the results in the performed studies with each other as well as lack of adequate studies in this regard in the city of Garmsar led us towards analyzing the prevalence of low birth weight in that city in 2013. The relations of maternal and infantile factors with the birth weight in infants were also investigated.

METHODS

The present study is a cross sectional study. The total infants under Garmsar urban centers who were born in 2013 (683 infants) formed our statistical population. Collecting the required data has done by census. The criterion to enter the study included all the infants who were born in 2013, and the criteria regarding lack of admission into the study included twins or multiple births and inadequate information about the infant or the mother. The case group included infants with low weight, i.e. the infants who had birth weight less than 2500g. The control group consisted of the infants with birth weight of above 2500g. The required information was collected by considering the health file of the families, and completion of the checklists of registering the information with variables related to the infants (sex, weight) and mothers (BMI, age, education, occupation, nationality, i.e. Iranian or Afghani). The other collected information included the number of child births, the interval of less than 3 years between the last and the previous childbirths, systolic and diastolic blood pressures and urinary tract infection. To observe moralities in this study, the required permissions were obtained from the research department of the university, the province and the city health centers, and the personal information of the participants were considered confidential, and the data were analyzed in general.

After collection and controlling again, the data were placed in SPSS19 software. The descriptive statistics (average and standard deviation) were used for describing the data, and the analytical statistics (independent t-test, chi-square test and logistic regression) were used for analyzing them.

RESULTS

According to the results, 32 infants out of 683 infants born in 2013 have low birth weight, i.e. the weight less than 2500g. LBW prevalence estimated in this study was 4.7%.

The average age of mothers who gave birth to infants with less than 2500g of weight was 27.8+1.3, and the average age of mothers in the group bearing infants with over 2500g of weight was 27.6+0.2. This indicates that their age difference was not statistically significant (P=0.877). The average systolic blood pressure in the mothers having low-weighed infants was 10.91, and it was 10.31 in mothers having infants with the weight of over 2500g (P=0.001). The average diastolic blood pressure in mothers having infants with low weight was 7.03, and 6.62 in mothers having infants with the weight of over 2500g (P=0.004).

Sixty two point five percent of the women, who gave birth to infants with the weight less than 2500g, and 46.7% of the women, who had infants with the weight of over 2500g, had normal BMI.

According to Table 1, there were no significant relations between low birth weight

and the variables including infants' sex, mothers' education and urinary infections during pregnancy, nationality, mothers' ages and mothers' BMI rates.

Table 1:Distribution of variables related to maternal and neonatal variables of infan	s with	low
birth weight		

Variable		Low birth weight	Р	
	Thin	5(7.4%)		
BMI	Normal	20(6.1%)	0.099	
	Over weight	5(2.8%)		
	Obesity	2(1.9%)		
	1	25(6.8%)		
Number of births	2	5(2.2%)	0.015	
	>2	2(2.3%)		
	Housewife	19(3.5%)		
Mother occupation	Practitioner	13(10.1%)	0.027	
	Student	0(0%)		
	Illiterate	2(3.2%)		
Mother education	The diploma and diploma	15(4%)	0.475	
	Collegiate	15(6.1%)		
The interval of less than 3 years	Yes	27(5.9%)		
between the last and the previous	No	5(2.2%)	0.024	
childbirths				
Mother urinary tract infection	Yes	7(4.6%)		
Nationality	No	25(4.7%)	0.211	
	Iranian	30(5%)		
	Afghani	2(2.3%)	0.414	
K Z	Male	15(4.4%)		
Infant sex	Female	17(5%)	0.698	

There were significant statistical relations between low birth weight and the variables including number of births, mothers' occupation and interval of less than 3 years between pregnancies and systolic and diastolic blood pressures.

The predicting rates by logistic regression analysis of different variables in

emergence of low weight can be seen in Table 2. The variables relating to low weight (number of births, mothers' occupation and interval of less than 3 years between pregnancies and systolic and diastolic blood pressures), or the ones close to being significant (BMI) were used in the logistic model. And, then we observed that LBW had significant relations only with systolic blood pressure, occupation of mothers and

BMI, when the effects of other variables are under the control.

Table 2: Distribution	of predictive variables in low birth we	eight infants using logistic regression
	analysis	

Variable		Low birth weight	Р	OR	95% confidence
					interval for OR
	Thin	5(7.4%)			
BMI	Normal	20(6.1%)	0.031	1.677	1.047-2.684
	Over weight	5(2.8%)			
	Obesity	2(1.9%)			
	1	25(6.8%)			
Number of births	2	5(2.2%)	0.198	1.678	0.72-4.898
	>2	2(2.3%)			
	Housewife	19(3.5%)			
Mother occupation	Practitioner	13(10.1%)	0.037	0.490	0.250-0.959
	Student	0(0%)			
The interval of less than 3 years between	Yes	27(5.9%)			
the last and the previous childbirths	No	5(2.2%)	0.702	1.313	0.326-5.282
Systolic pressure	-	10.91	0.050	0.620	0.384-1.001
Diastolic pressure	-	7.03	0.512	0.817	0.445-1.497

DISCUSSION

According to the results, 32 infants out of 683 infants born in 2013 had low birth weight, i.e. the weight less than 2500g. LBW prevalence estimated in this study was 4.7%. There were significant statistical relations between low birth weight and the variables including number of births, mothers' occupation and interval of less than 3 years between pregnancies and systolic and diastolic blood pressures.

LBW prevalence is 9.9% in Pakistan.¹⁸ and 7.4% in Japan.¹⁹ This rate is reported in Ardabil to be 6.3%, 7.2% in Shahroud, and 7% in Rafsanjan.^{20,21} According to the results of this study, LBW prevalence was 4.7%, indicating

less prevalence in comparison with similar studies.

Increasing the number of childbirths reduces lower birth weight, significantly.²² In other words, there is a significant positive relation between the numbers of childbirths and increasing birth weight.²³ By increasing the number of childbirths, the birth weight shows also an increase in this study, but this relation was not significant.

The results of some epidemiologic studies showed that the rate of emergence of low birth weight at the beginning and the end of fertility, i.e. between 15 to 19 years of age and between 35-40 years of age, is higher.^{24,25} It is believed that the possibility of LBW emergence is higher in women over 35 years of age. However, there is a lot of doubt about it. Many researchers have suggested that this risk is not related to mothers' ages, but it is related to problems like many chronic diseases (diabetes and blood pressure).²⁶ Anyhow, no significant relation was observed in this study between the average age of mothers and low birth weight. These results were in conformity with the results of studies by Lio et al., Yogo et al., and Delbaere et al.²⁷⁻²⁹

In many studies significant relation was observed between LBW and the interval between pregnancies. By decreasing this interval, the rate of LBW increases.^{30,31} The intervals between pregnancies, varied in the developing countries (3-6 months) and in the developed countries (1-2 years), could increase LBW due to the pregnancy.³² In this study, a significant relation was also observed between the intervals in pregnancies of less than 3 years and low birth weight.

The results of some studies showed the relation between hypertension and LBW.³³⁻³⁵ Hypertension in mothers during pregnancy reduces blood flow in embryos, leading to reducing embryo's growth and low birth weight, but this is not approved epidemiologically.³⁴ Significant relation was observed in this study between systolic and diastolic blood pressures and LBW.

The results of the study showed that low weight in girls is more common than in boys, but the relation did not become significant. Thus, the results in this study were not in conformity with the results in the studies by Khalil et al., Orescov et al., and Hashemiannejad et al., who showed relations existing between the sex and birth weight.³⁶⁻³⁸

No significant relation was observed in this study between mothers' education and LBW. The results were in conformity with the results obtained by Chaman et al., Zeighami et al., and Delaram, but they were not in conformity with the results of Roudbari studies.^{21,31,39,40}

Mothers' weights and BMI are important factors before pregnancy, since there is a close relation between mothers' and embrvos' nutritional conditions and the birth weight, and embryos' growth is a function of mothers' nutritional conditions, especially in pre-pregnancy period. Thus, mothers' BMI rates, before or at the beginning of pregnancy, could be used for evaluating the infants' weights.41,42 Significant relation is observed in Yodav et al., and Hassan et al. studies, between BMI and low birth weight.^{43,44} By controlling altering variables in this study, a significant relation was observed between mother's MMI snf LBW of the infant.

The results of the study showed relations between LBW and the mother's occupation. It was in conformity with the results obtained by Fallah et al., and Toutounchi el al., but had no conformity with study results of Esmaeeli et al.^{35,45,46}

The results showed no relations between LBW and nationality (Iranian and Afghani). It had no conformity with the result obtained by Rezaian et al.⁴⁷

The difference in the results about existence or lack of relations between LBW risk factors in different studies could be due to differences in methods, population under study and the sample volume used in the studies. A disorder in birth weight is one of the important factors in emergence of many short-term or long-term consequences in life, and is a main determining factor in health aspects. Thus, regarding the approaches that prevent the emergence of this consequence, the pre-pregnancy cares should be highly considered, so that by such cares, women could be ready for a safe and secure pregnancy.

CONCLUSION

Although our results regarding LBW prevalence and risk factors confirm the results of other studies, it was recommended perspective studies to reconfirm the LBW risk factors.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENT

Hereby we would like to express our especial thanks to the Research Council of Semnan University of Medical Sciences, co-vice president of health care in Garmsar and specially Garmsar health center staffs for their unwavering support in this study.

REFERENCES

1. Kramer MS. Determinants of low birth weight: methodological assessment and meta-analysis. Bull World Health Organ. 1987; 65(5): 663-737.

2. Rich-Edwards JW, Buka SL, Brennan RT, Earls F. Diverging associations of maternal age with low birthweight for black and white mothers. Int J Epidemiol. 2003; 32(1): 83-90.

3. Kumar P, Seshadri R. Neonatal morbidity and growth in very low birth-weight infants after multiple courses of antenatal steroids. J Perinatol. 2005; 25(11): 698-702.

4. Basu S, Rathore P, Bhatia BD. Predictors of mortality in very low birth weight neonates in India. Singapore med J. 2008; 49(7): 556-60.

5. Kang BH, Moon JY, Chung SH, Choi YS, Lee KS, Chang JY, et al. Birth statistics of high birth weight infants (macrosomia) in Korea. Korean J Pediatr. 2012; 55(8): 280-5.

6. Forsen T, Eriksson J, Tuomilehto J, Reunanen A, Osmond C, Barker D. The fetal and childhood growth of persons who develop type 2 diabetes. Ann Intern Med. 2000; 133(3): 176-82.

7. Frankel S, Elwood P, Sweetnam P, Yarnell J, Smith GD. Birthweight, body-mass index in middle age, and incident coronary heart disease. Lancet. 1996; 348(9040): 1478-80. 8. Whincup P, Cook D, Papacosta O, Walker M. Birth weight and blood pressure: cross sectional and longitudinal relations in childhood. Bmj. 1995; 311(7008): 773-6.

9. Akre O, Ekbom A, Hsieh CC, Trichopoulos D, Adami HO. Testicular nonseminoma and seminoma in relation to perinatal characteristics. J Natl Cancer Inst. 1996; 88(13): 883-9.

10. Tibblin G, Eriksson M, Cnattingius S, Ekbom A. High birthweight as a predictor of prostate cancer risk. Epidemiology. 1995; 6(4): 423-4.

11. Valero De Bernabe J, Soriano T, Albaladejo R, Juarranz M, Calle ME, Martinez D, et al. Risk factors for low birth weight: a review. Eur J Obstet Gynecol Reprod Biol. 2004; 116(1): 3-15.

12. Nazari F, Vaisi Z, Sayehmiri K, et al. Prevalence and trends of low birth weight in Iran: A systematic review. J Nurs MIdwifery. 2013; 22(79):1-8.

13. Emanuel I, Kimpo C, Moceri V. The association of maternal growth and socioeconomic measures with infant birthweight in four ethnic groups. Int J Epidemiol. 2004; 33(6): 1236-42.

14. Astone NM, Misra D, Lynch C. The effect of maternal socio-economic status throughout the lifespan on infant birthweight. Paediatr Perinat Epidemiol. 2007; 21(4): 310-8.

15. Tootoonchi P. Low birth weight among newborn infants at Tehran hospitals. J Pediatr. 2007; 17(2): 186-92.

16. Zahedi R, Managheb SE., karimi MS. Investigation of effective factors on low birth weight in Jahrom. J Jahrom UnivMed Sci. 2009; 6(2): 56-65.

17. Shin YH, Choi SJ, Kim KW, Yu J, Ahn KM, Kim HY, et al. Association between maternal characteristics and neonatal birth weight in a Korean population living in the Seoul metropolitan area, Korea: a birth cohort study (COCOA). J Korean Med Sci. 2013; 28(4): 580-5.

18. Badshah S, Mason L, McKelvie K, Payne R, Lisboa PJ. Risk factors for low birthweight in the public-hospitals at Peshawar, NWFP-Pakistan. BMC Public Health. 2008; 8: 197.

19. Suzuki K, Tanaka T, Kondo N, Minai J, Sato M, Yamagata Z. Is maternal smoking during early pregnancy a risk factor for all low birth weight infants? J Epidemiol. 2007; 18(3): 89-96.

20. Mirzarahimi M, Hazrati S, Ahmadi P, Alijahan R. Prevalence and risk factors for low birth weight in Ardabil, Iran. Iran J Neonatol. 2013: 4(1): 18-23.

21. Chaman R, et al. Low Birth Weight and Its Related Risk Factors. HK J Paediatr. 2012; 17: 191-2.

22. Rezaeian M, Goujani R, Sheikh Fathollahi M, Vaziri Nejad R, Manshori A, Razi S. A Comparative Study on Prevalence of Preterm Birth and Low Birth Weight in Iranians and Afghans Races in Rafsanjan Nik-Nafs Hospital in 2011-2012. J Rafsanjan Univ Med Sci. 2014; 13(1): 67-82.

23. Figueras F, Meler E, Iraola A, Eixarch E, Coll O, Figueras J, et al. Customized birthweight standards for a Spanish population. Eur J Obstet Gynecol Reprod Biol. 2008; 136(1): 20-4.

24. Xue F, Willett WC, Rosner BA, Forman MR, Michels KB. Parental characteristics as predictors of birthweight. Hum Reprod. 2008; 23(1): 168-77.

25. Ziadeh S. Obstetric outcome of teenage pregnancies in North Jordan. Arch Gynecol Obstet. 2001; 265(1): 26-9.

26. Rebollo AG, Montero CM. [Perinatal variables and health inequalities in a health-care district in Caceres, Spain]. Gac Sanit. 2000; 14(1): 31-8.

27. Cnattingius S, Forman MR, Berendes HW, Isotalo L. Delayed childbearing and risk of adverse perinatal outcome. A population-based study. JAMA. 1992; 268(7): 886-90.

28. Hsieh TT, Liou JD, Hsu JJ, Lo LM, Chen SF, Hung TH. Advanced maternal age and adverse perinatal outcomes in an Asian population. Eur J Obstet Gynecol Reprod Biol. 2010; 148(1): 21-6.

29. Yogev Y, Melamed N, Bardin R, Tenenbaum-Gavish K, Ben-Shitrit G, Ben-Haroush A. Pregnancy outcome at extremely advanced maternal age. Am J Obstet Gynecol. 2010; 203(6): 558 e1-7.

30. Delbaere I, Verstraelen H, Goetgeluk S, Martens G, De Backer G, Temmerman M. Pregnancy outcome in primiparae of advanced maternal age. Eur J Obstet Gynecol Reprod Biol. 2007; 135(1): 41-6.

31. Jafari F, Eftekhar H, Pourreza A, Mousavi J. Socio-economic and medical determinants of low birth weight in Iran: 20 years after establishment of a primary healthcare network. Public Health. 2010; 124(3): 153-8.

32. Roudbari M, Yaghmaei M, Soheili M. Prevalence and risk factors of low-birthweight infants in Zahedan, Islamic Republic of Iran. East Mediterr Health J. 2007; 13(4): 838-45.

33. Ochoa Sangrador C, Luque Benlloch C, Carrascal Tejado A. [Prematurity, low birth weight and the interval between pregnancies]. An Esp Pediatr. 1996; 45(1): 67-70.

34. Ugboma H, Onyearugha C. Low birth weight delivery: Prevalence and associated factors as seen at a tertiary health facility. Niger J Clin Pract. 2013; 16(2): 184-7.

35. Xiong X, Demianczuk NN, Saunders LD, Wang FL, Fraser WD. Impact of preeclampsia and gestational hypertension on birth weight by gestational age. Am J Epidemiol. 2002; 155(3): 203-9.

36. Fallah M, Afrouz G, Heidari G. Examining the factors effective on birth weight among babies of Yazd Province in 2007. Toloo-e-behdasht. 2008; 7(3): 57-64. 37. Khalil MM, Alzahra E. Fetal gender and pregnancy outcomes in Libya: A retrospective study. Libyan J Med. 2013; 8: 1-4.

38. Orskou J, Henriksen TB, Kesmodel U, Secher NJ. Maternal characteristics and lifestyle factors and the risk of delivering high birth weight infants. Obstet Gynecol. 2003; 102(1): 115-20.

39. Hashemian Nejad N, Pejhan A, Rakhshani MH, Hoseini BL. The Incidence of Low Birth Weight (LBW) and Small-for-Gestational Age (SGA) and its Related Factors in Neonates, Sabzevar, Iran. Int J Pediatr. 2014; 2(4): 73-8.

40. Zeyghami B, Parisay Z.A Study on Correlation of Mother's Risk Factors with Low Birth Weight of Newborns at a Multiple Regression Model in Kohghiloyeh and Boyerahmad Province in 2004-2005. ArmaghaneDanesh. 2006; 10(4): 37-45.

41. Delaram M. The incidence and related factors of low birth weight. Iran J Nurs. 2010; 23(64): 29-36.

42. Goshtasebi A, Banaem LM, Rodbary MA, Bakouei S. The association between

rch

preconception body mass index and pregnancy weight gain on birth weight. J Mazandaran Univ Med Sci. 2011; 21(84): 81-5.

43. Nemmati A, Refahi S, Barak M, Jafari M, Ettehad G. The relationship between some of maternal anthropometric measurments with neonatal birth weight of Alavi Hospital, Ardebil. J Ardabil Unvi Med Sci. 2007; 7(1): 84-9.

44. Yadav H, Lee N. Maternal factors in predicting low birth weight babies. Med J Malaysia. 2013; 68(1): 44-7.

45. Hassan NE, Shalaan AH, El-Masry SA. Relationship between maternal characteristics and neonatal birth size in Egypt. East Mediterr Health J. 2011; 17(4): 281-9.

46. Tootonchi P. Low birth weight among newborn infants at Tehran hospitals. Iran J Pediatr. 2007; 17: 186-92.

47. Esmaeili H. The relationship between BMI at the beginning of pregnancy With birth weight and other pregnancy outcomes. J Obst Gynecol Infertil. 2014; 16: 1-10.

How to cite the article: Safari M, Samiee A, Salehi F, Ahmadi SN, Ahmadi SS. The prevalence and related factors of low birth weight. Int J Epidemiol Res. 2016; 3(3): 214-221.