

Epidemiology and trend of stomach cancer mortality in Iran (2006-10)

Neda Amoori¹, Sepideh Mahdavi², Mostafa Enayatrad^{3*}

¹*Biostatic and Epidemiology Dept., Abadan University, Abadan, I.R. Iran;*

²*Epidemiology and Biostatistics Dept., Yazd ShahidSadoughi University of Medical Sciences, Yazd, I.R. Iran;*

³*Epidemiology Dept., School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, I.R. Iran.*

Received: 8/Dec/2015 Accepted: 4/Apr/2016

ABSTRACT

Background and aims: Stomach cancer is one of the most common cancers and the second cause of deaths among cancers worldwide cancer in the world. This study aimed to estimate the rate and trend of death from stomach cancer in Iranian population between 2006 and 2010.

Methods: This study was a cross-sectional conducted using mortality data in Iranian population between 2006 and 2010. The number of deaths, death rate, and sex ratio of mortality, mortality rates by age and gender groups, and trends of mortality from stomach cancer were calculated and reported. Cochran Armitage test was used for linear trend by Winpepi software to study the trends of Stomach cancer mortality.

Results: The results showed that between 2006 and 2010, 34950 cases of death have been reported from stomach cancer. The mortality rate per 100,000 reached from 11.22 in 2006 to 11.06 in 2010. All years studied indicated that death due to stomach cancer were higher in males than females. Also, with increasing age, the mortality rate of stomach cancer was increased.

Conclusion: Our findings showed the decreasing trend of mortality due to stomach cancer may be due to increasing in lifespan and survival, enjoying the better diagnostic and therapeutic procedures for patients, and underestimation of the number of stomach cancer deaths. Since the cancer is the most frequent cause of death among cancers in Iran, additional investigation is necessary to determine the risk factors of the cancer to decline the mortality rate.

Keywords: Stomach cancer, Trend, Mortality, Iran.

Original article

INTRODUCTION

Cancer is considered as one of the most common causes of death worldwide because it's incidence and prevalence is increasing.¹⁻⁴

It is estimated that cancer prevalence increases up to 45% in developing countries by 2025.⁵⁻⁹ Cancers are important in

***Corresponding author:** Mostafa Enayatrad, Epidemiology Dept., School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran, Tel: 00989163442067, E-mail: mostafaenayatrad@gmail.com

designing and planning health system programs due to irreparable consequences and staggering costs of diagnosis and treatment. More than half of the cancers and 60% of cancer deaths occur in less developed countries, and along with changing lifestyle, similar to the Western countries, cancer incidence is increasing in developing countries.¹⁰

Among the cancers, gastrointestinal cancer is prevalent in Iranian men while breast cancer is the most common cancer in Iranian women. Based on cancer registry reports in Iran stomach cancer is the most frequent, then the colon and rectum cancers are the most common gastrointestinal cancers.¹⁰ Latest reports indicated that the incidence of stomach cancer is nearly 934,000 cases per year, and the cancer is considered the fourth most common cancer and after lung cancer, the second cause of deaths from cancer in the world.⁵ In contrast with the declining trend of stomach cancer worldwide, new cases of the cancer are increasing every year.¹¹ It also has remained as a public health problem in 21 countries. This could be due to increasing in the human lifespan.¹²

Stomach cancer is one of the most important gastrointestinal tract cancers.¹³ According to the estimations of the International Agency for Research on Cancer (IARC) in 2012; stomach cancer is the fourth most common cancer in the world. Half of the cases occur yearly in East Asia and mostly in China. Stomach cancer is third cancer mortality of both sexes in the worldwide. Moreover, in 2012 in Iran Stomach cancer accounted for 11.4% of all cases of cancer occurrence and was considered as the second most prevalent cancer. In the same year, Stomach cancer which accounted for 15.5 percent of all mortalities caused by cancers was in fact the most deadly cancer in Iran.¹⁴⁻¹⁷ The current global trend of

stomach cancer incidence has shown a substantial decline since 1975, when stomach cancer was the world's most common cancer.¹⁸ Stomach cancer is much more common in certain Asian, Central European, Central American, and South American countries, especially Japan, Chile, Costa Rica, Hungary, and Poland. The highest incidence of stomach cancer occurs in Japan and Eastern Asia; by contrast, its incidence is relatively low in Western Europe.¹⁹

Smoking and chronic infections are the most important risk factors for all cancers, especially stomach cancer so that they cause 30% and 10% of all cancers, respectively.¹ Some studies expressed that *Helicobacter pylori* and poor diet are the most important determinant factor for stomach cancer. *Helicobacter pylorus*, as one of the most common human infections worldwide, has prevalence of 50% in the world and 90% in developing countries. The prevalence of bacterium is also high in Iran.²⁰ High-salt diet is another risk factor but a healthy diet, including fresh fruits and vegetables, may reduce the risk of disease.²¹

One of the most important steps in controlling cancer is collecting data on cancer mortality in any country. To evaluate activities for cancer control, monitoring trends in cancer mortality by demographic characteristics and time is necessary. This study aimed to estimate the rate and trend of death from stomach cancer in Iranian population between 2006 and 2010.

METHODS

The present study was a cross-sectional using data on mortality in Iran during 2006-10. Data were collected by Technology and Applied Research and Information Management Center of the Ministry of Health and Medical Education from various sources including National Organization for

Civil Registration, cemeteries, hospitals, and care homes. The necessary data integration has been performed in the Ministry of Health and Medical Education.²²

In the present study, the data on the number of deaths from Stomach cancer during 2006-10 in 29 provinces of Iran were obtained based on mortality in Iran after removing empty codes, and were entered on a pre-designed form. Meanwhile, cancers were coded based on The International Classification of Diseases for Oncology ICD-O (second edition). The C16 code was given to Stomach cancer. Number of deaths, mortality rate, male/female mortality ratio, and mortality rates in the age and sex groups were calculated and reported. Moreover, a picture of the trend of changes in mortality due to Stomach cancer in Iran was drawn. Charts with excel 2010 software was performed.

Cochrane Armitage test was used for linear trend by Winpepi software 2.1 to study the trends in cancer mortality.

RESULTS

In this study, all data on mortality from stomach cancer was investigated in all sex and age groups between 2006 and 2010. Of 34, 950 deaths from stomach cancer reported, 22,695 cases occurred in men and 12,255 in women. The highest number of deaths in males (6105 cases) and females (4050 cases) was seen in 2010. Overall, the number of deaths was more in men than women and the sex ratio was 1.85.

The results obtained from calculation of mortality rates per 100,000 people showed that the death rates from stomach cancer reached from 11.22 in 2006 to 11.06 in 2010. Mortality rates in all years studied indicated stomach cancer prevalence was higher in males than females. As shown in Table1 and Figure1, the mortality rate in males reached from 14.7 per 100,000 people in 2006 to 15.51 per 100,000 people in 2010, while the rate was 7.64 per 100,000 people in 2006 and 7.53 per 100,000 people in 2010 among females.

Table 1: Mortality from stomach cancer per hundred thousand people in the study group sex

Year of study	Mortality per hundred thousand			Crude mortality rate			Sex ratio
	Woman	Men	Total	Woman	Men	Total	
2006	7.64	14.70	11.20	2147	4260	6407	1.98
2007	7.92	15.47	11.75	2075	4159	6234	2.00
2008	7.43	15.09	11.31	1971	4103	6074	2.08
2009	7.49	14.78	11.17	2012	4068	6080	2.02
2010	7.53	15.51	11.06	4050	6105	10155	1.50
Total				12255	22695	34950	1.85




Figure 1: The rate of death from stomach cancer per hundred thousand people in the study group sex

As seen in Table 2 and Figure 2, results in all sex and age groups studied showed with increase in age, the mortality rate of stomach cancer also increases. The lowest mortality rate due to the cancer was observed in the age group of less than 5 years and between 5 to 14 years, but as age increases, the rate

increased in people over 70 years. In other words, in men older than 70 years, the mortality rate was increased from 244.85 per 100,000 people in 2006 to 243.44 per 100,000 people in 2010, and in women, the rate was 118.71 per 100,000 in 2006 and 123.84 per 100,000 in 2010.

Table 2: The rate of death from stomach cancer per hundred thousand in the year under study, by sex and age group

Year of Study	Under 5 years			5-14			15-49			50-69			Above 70 years		
	W	M	T	W	M	T	W	M	T	W	M	T	W	M	T
2006	0	0	0	0	0	0	1.47	2.41	1.95	26.62	48.94	37.68	118.71	224.85	175.44
2007	0	0	0	0	0	0	1.66	2.79	2.23	25.14	49.81	37.30	129.49	241.78	189.22
2008	0	0	0	0	0	0	1.43	2.10	1.76	24.74	49.22	36.79	118.92	246.58	186.32
2009	0	0	0	0	0	0	1.20	2.19	1.70	23.15	47.81	35.28	129.07	242.14	188.30
2010	0	0	0	0	0	0	1.45	2.38	1.92	23.38	44.49	33.77	123.84	243.44	185.98




Figure 2: The rate of death from stomach cancer per hundred thousand in the year under study, by sex and age

DISCUSSION

Studies determined that stomach cancer is the most frequent cause of death among cancer from tumor mass in the world and the most frequent cause of death from cancer in Iran. The

mortality rate from the cancer was 8.4 per100,000 in East Azerbaijan and north of Iran compared with 3.6 per 100,000 in south of Iran.^{23,24} In comparison during the last decades,

the mortality due to the cancer has significantly decreased in most developed countries, in some Asian countries (China, Korea and Japan), East Europe, and South America there is a high prevalence.²⁵ According to the estimations of the International Agency for Research on Cancer (IARC) in 2012, stomach cancer is the fourth most common cancer in the world. Half of the cases occur yearly in East Asia and mostly in China. Stomach cancer is third cancer mortality of both sexes worldwide. Moreover, in 2012 in Iran Stomach cancer accounted for 11.4% of all cases of cancer occurrence and was considered as the second most prevalent cancer. In the same year, stomach cancer which accounted for 15.5 percent of all mortalities caused by cancers was in fact the most deadly cancer in Iran.^{16,17,26} In Iran, while the northern and northwestern regions are high risk areas for gastric cancer, there are several intermediate and low risk populations in other geographical areas.² Stomach cancer is the second common cancer in Iranian men and the fourth in Iranian women. Its incidence is more in males than females.²⁷ In Iran, most northern and north western areas are at high risk for stomach cancer, whereas the central and western provinces are at medium risk and the southern regions are at a low risk.¹⁵ The highest incidence of Stomach cancer in the country was seen in Mazandaran, Golestan, and Ardabil provinces but the lowest was reported from the Kerman province.²⁸ Another study showed that the greatest rate was reported from the Ardebil province and the lowest was observed in Khuzestan and Chaharmahal and Bakhtiari provinces.²⁹

Findings indicated that the mortality rate due to stomach cancer has been constant during years of the study. That is, the mortality rate reached from 11.22 to 11.06. Another study revealed that stomach cancer mortality rate has increased from 1.64 to 9.67 per 100,000 between 1995 and 2003.³⁰ A study reported that in 2004 deaths from the cancer were 12.02 per 100,000.³¹ It is also shown that the mortality rate was 15 and 8.1 per 100,000 in men and women, respectively.³²

In our study, deaths from the cancer were more in patients over 70 years. In other words, as age increases, the mortality rate increases. A study conducted in the Mazandaran province,

North of Iran, also emphasized that the mortality rate was more in people over 60 than other age groups. As well, the mean age at death was higher for men than women.³³ Our results; similar to some studies showed that sex ratio (male/female) for stomach cancer mortality was 1.85, unlike some studies that reported the sex ratio of males to females as doubled.³²⁻³⁵

Etiologic factors related to stomach cancer are unknown but many environmental factors, such as age, sex, blood group, heredity, poor diet, geographical area, smoking, and alcohol, are possible risk factors for developing the cancer.^{35,36} Given the significant geographic and ethnic differences in the incidence of stomach cancer in various areas, it can be concluded that the cancer risk is specified to a large extent by environmental factors such as dietary factors and *Helicobacter pylori* infection.²⁸ *Helicobacter pylori* is considered as the most infectious agent for developing stomach cancer.^{37,38} The prevalence of this infection is high in Iran.³⁹ The prevalence of *Helicobacter pylori* in the developing countries, including Iran, is more than 80% and in developed countries less than 30%.³⁸ In a population-based study, conducted in Ardebil (a province in West of Iran), 90% of the population at risk of the infection was known and the lowest prevalence of the bacterium was seen in the Sistan and Baluchistan province, south East of Iran, where there is the lowest incidence of stomach cancer.^{40,41}

The results obtained from this study is different from other studies regarding mortality from stomach cancer because the findings showed that the mortality rate of the cancer is decreasing, while another study on the data of cancer registry reported an increasing trend related to deaths from the disease.^{2,42-44} However, the decrease may be due to increasing lifespan and survival, enjoying the better diagnostic and therapeutic procedures for patients and underestimation of the number of death due to stomach cancer. Since the cancer has the most frequent deaths from cancers in Iran, additional investigation is necessary to determine risk factors of the cancer to reduce the mortality rate. Diagnostic interventions are required to reduce death from this cancer.

Data on cancer mortality can be used to guide policy makers in order to setup cancer

prevention programs. But this aim needs reliable death registry systems which reports death statistics annually. On the other hand, the analysis of death statistic subject to misclassification is a major problem in epidemiological analysis leading to biases estimates, and can therefore cause one to underestimate health risks.⁴⁵

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENT

The authors also express their gratitude to all who has helped the researchers of this study.

REFERENCES

1. Stewart B, Kleihues P. World cancer report. Lyon: IARC press; 2003.
2. Enayatrads M, Salehiniya H. Trends in gastric cancer incidence in Iran. *J Mazandaran Univ Med Sci.* 2014; 24(114): 8-16.
3. Enayatrads M, Amoori N, Salehiniya H. Epidemiology and trends in breast cancer mortality in Iran. *Iran J Public Health.* 2015; 44(3): 430-1.
4. Enayatrads M, Salehiniya H. An investigation of changing patterns in breast cancer incidence trends among Iranian women. *Journal of Sabzevar University of Medical Sciences.* 2014; 22(1): 27-35.
5. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin.* 2005; 55(2): 74-108.
6. Koochi F, Enayatrads M, Salehiniya H. A Study of the Epidemiology and Trends in Cancer Incidence in Iranian Elderly 2003-2009. *Arak Med University J.* 2015; 18: 57-66.
7. Rafiemanesh H, Enayatrads M, Salehiniya H. Epidemiology and Trends of Mortality from prostate cancer in Iran. *J Isfahan Med School.* 2015; 33(220): 1-7.
8. Razi S, Enayatrads M, Mohammadian-Hafshejani A, Salehiniya H, Fathali-Loy-Dizaji M, Soltani S. The epidemiology of skin cancer and its trend in Iran. *Int J Prev Med.* 2015; 6: 64.
9. Almasi Z, Rafiemanesh H, Salehiniya H. Epidemiology characteristics and trends of incidence and morphology of stomach cancer in Iran. *Asian Pac J Cancer Prev.* 2015; 16(7): 2757-61.
10. Moghimi-Dehkordi B, Safaee A, Zali MR. Comparison of colorectal and gastric cancer: Survival and prognostic factors. *Saudi J Gastroenterol.* 2009; 15(1): 18-23.
11. Sadjadi A, Zahedi M, Nouraei M, Alimohammadian M, Ghorbani A, Bahmanyar S, et al. The first population-based cancer survey in Kerman Province of Iran. *Iran J Public Health.* 2007; 36(4): 26-34.
12. Pourfarzi F, Whelan A, Kaldor J, Malekzadeh R. The role of diet and other environmental factors in the causation of gastric cancer in Iran: A population based study. *Int J Cancer.* 2009; 125(8): 1953-60.
13. Haidari M, Nikbakht MR, Padsar Y, Najaf F. Trend analysis of gastric cancer incidence in Iran and its six geographical areas during 2000-2005. *Asian Pac J Cancer Prev.* 2012; 13(7): 3335-41.
14. IAFRo C. GLOBOCAN 2012: Estimated cancer incidence, mortality and prevalence worldwide in 2012: World Health Organization; 2014 [Accessed on, 9]. Available from: http://globocan.iarc.fr/Pages/fact_sheets_cancer.aspx.
15. Enayatrads M, Salehiniya H. Trends in gastric cancer incidence in Iran. *J Mazandaran Univ Med Sci.* 2014; 24(114): 8-16.
16. Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, Rosso S, Coebergh JW, Comber H, et al. Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. *Eur J Cancer.* 2013; 49(6): 1374-403.
17. Ferro A, Peleteiro B, Malvezzi M, Bosetti C, Bertuccio P, Levi F, et al. Worldwide trends in gastric cancer mortality

- (1980-2011), with predictions to 2015, and incidence by subtype. *Eur J Cancer*. 2014; 50(7): 1330-44.
18. Kim Y, Park J, Nam BH, Ki M. Stomach cancer incidence rates among Americans, Asian Americans and Native Asians from 1988 to 2011. *Epidemiol Health*. 2015; 37: e2015006.
 19. Sunny L, Yeole BB, Hakama M, Shiri R, Mathews S, FalahHassani K, et al. Decreasing trend in the incidence of stomach cancer in Mumbai, India, during 1988 to 1999. *Asian Pac J Cancer Prev*. 2004; 5(2): 169-74.
 20. Yazdizadeh B, Jarrahi AM, Mortazavi H, Mohagheghi MA, Tahmasebi S, Nahvijou A. Time trends in the occurrence of major GI cancers in Iran. *Asian Pac J Cancer Prev*. 2005; 6(2): 130-4.
 21. Boyle P, Levin B. World cancer report 2008: IARC Press, International Agency for Research on Cancer; 2008.
 22. Khosravi A, Aghamohamadi S, Kazemi E, Pour Malek F, Shariati M. Mortality profile in Iran (29 provinces) over the years 2006 to 2010. Tehran: Ministry of Health and Medical Education Pub; 2013.
 23. Pera M, Cameron AJ, Trastek VF, Carpenter HA, Zinsmeister AR. Increasing incidence of adenocarcinoma of the esophagus and esophagogastric junction. *Gastroenterology*. 1993; 104(2): 510-3.
 24. Malekzadeh R, Sajjadi R, Derakhshan MH. Ardabil cancer registry data population-based cancer. Second abstracts, National Congress of Medical Science Research, Ardabil Club Young. 2003; 105.
 25. Brenner H, Rothenbacher D, Arndt V. Epidemiology of stomach cancer. *Methods Mol Biol*. 2009; 472: 467-77.
 26. WHO. Fact sheet N297; [cited 2012 Accessed September 10]. Available from: http://www.who.int/mediacentre/factsheets/fs_297/en/index.html.
 27. Hamta A, Solimani M, Rasoli Z. Determination of chromosomal changes in DMBA-induced skin cancer in SD rat strains. *J Arak Univ MedSci*. 2009; 12(2): 73-87.
 28. Malekzadeh R, Derakhshan MH, Malekzadeh Z. Gastric cancer in Iran: epidemiology and risk factors. *Arch Iran Med*. 2009; 12(6): 576-83.
 29. Sadjadi A, Malekzadeh R, Derakhshan MH, Sepehr A, Nourai M, Sotoudeh M, et al. Cancer occurrence in Ardabil: Results of a population-based cancer registry from Iran. *Int J Cancer*. 2003; 107(1): 113-8.
 30. Pourhoseingholi M, Faghihzadeh S, Hajizadeh E, Gatta G, Zali M, Abadi A. Trend analysis of gastric cancer and colorectal cancer mortality in Iran, 1995-2003. *Iran J Cancer Prev*. 2011; 4(1): 38-43.
 31. Mousavi SM, Somi MH. Gastric cancer in Iran 1966-2006. *Asian Pac J Cancer Prev*. 2009; 10(3): 407-12.
 32. Zendehtdel K, Marzban M, Nahvijou A, Jafari N. Six-fold difference in the stomach cancer mortality rate between northern and southern Iran. *Arch Iran Med*. 2012; 15(12): 741-6.
 33. Charati JY, Zare S, Ghorbanpour E, Shabankhani B. Demographic and geographical pattern of mortality rate from stomach cancer and related factors in Mazandaran province from 2001 to 2005. *J Mazandaran Unive Med Sci*. 2010; 20(79): 1-7.
 34. Price P, Sikore K. Treatment of cancer. 4th ed. London: Arnold Press; 2002: 583-9935.
 35. De Stefani E, Boffetta P, Carzoglio J, Mendilaharsu S, Deneo-Pellegrini H. Tobacco smoking and alcohol drinking as risk factors for stomach cancer: A case-control study in Uruguay. *Cancer Causes Control*. 1998; 9(3): 321-9.
 36. Stael von Holstein C, Eriksson S, Huldt B, Hammar E. Endoscopic screening during 17 years for gastric stump carcinoma. *Scand J Gastroenterol*. 1991; 26(10): 1020-6.
 37. Helicobacter, Group CC. Gastric cancer and Helicobacter pylori: a combined analysis of 12 case control studies nested

within prospective cohorts. *Gut*. 2001 September 1, 2001; 49(3): 347-53.

38. Persson C, Jia Y, Pettersson H, Dillner J, Nyren O, Ye W. H. pylori seropositivity before age 40 and subsequent risk of stomach cancer: A glimpse of the true relationship? *PLoS One*. 2011; 6(3): e17404.

39. Samadi F, Babaei M, Yazdanbod A, Fallah M, Nouraei M, Nasrollahzadeh D, et al. Survival rate of gastric and esophageal cancers in Ardabil province, North-West of Iran. *Arch Iran Med*. 2007; 10(1): 32-7.

40. Malekzadeh R, Sotoudeh M, Derakhshan MH, Mikaeli J, Yazdanbod A, Merat S, et al. Prevalence of gastric precancerous lesions in Ardabil, a high incidence province for gastric adenocarcinoma in the northwest of Iran. *J ClinPathol*. 2004; 57(1): 37-42.

41. Metanat M, Sharifi-Mood B, Izadi S. Prevalence of *Helicobacter pylori* infection

in healthcare workers. *Turkish J Med Sci*. 2010; 40(6): 965-9.

42. Amoori N, Mirzaei M, Cheraghi M. Incidence of cancers in Kuzestan province of Iran: Trend from 2004 to 2008. *Asian Pac J Cancer Prev*. 2014; 15(19): 8345-9.

43. Karami K, Cheraghi M, Amori N, Pedram M, Sobhani A. Common cancers in Khuzestan province, south west of Iran, during 2005-2011. *Asian Pac J Cancer Prev*. 2014; 15(21): 9475-8.

44. Keyghobadi N, Rafiemanesh H, Mohammadian-Hafshejani A, Enayatrad M, Salehiniya H. Epidemiology and trend of cancers in the province of Kerman: southeast of Iran. *Asian Pac J Cancer Prev*. 2015; 16(4): 1409-13.

45. Stamey JD, Young DM, Seaman JW, Jr. A Bayesian approach to adjust for diagnostic misclassification between two mortality causes in Poisson regression. *Stat Med*. 2008; 27(13): 2440-52.

How to cite the article: Amoori N, Mahdavi S, Enayatrad M. Epidemiology and trend of stomach cancer mortality in Iran (2006-10). *Int J Epidemiol Res*. 2016; 3(3): 265-275.