

Association of Serum 25-Hydroxyvitamin D Levels With Markers of Metabolic Syndrome in Adult Women in Ramsar, Iran

Zamzam Paknahad^{1,2,*}; Azam Ahmadi Vasmehjani²; Mohammad Reza Maracy³

¹Food Security Research Center, Isfahan University of Medical Sciences, Isfahan, IR Iran

²Department of Clinical Nutrition, Faculty of Nutrition and Food Sciences, Isfahan University of Medical Sciences, Isfahan, IR Iran

³Department of Epidemiology and Biostatistics, Faculty of Health, Isfahan University of Medical Sciences, Isfahan, IR Iran

*Corresponding author: Zamzam Paknahad, Department of Clinical Nutrition, Faculty of Nutrition and Food Sciences, Isfahan University of Medical Sciences, Isfahan, IR Iran. Tel: +98-3117922784, Fax: 98-3116682509, E-mail: paknahad@hlth.mui.ac.ir

Received: April 15, 2014; Revised: May 17, 2014; Accepted: May 25, 2014

Background: Epidemiological studies showed that vitamin D deficiency is associated with components of metabolic syndrome.

Objectives: The aim of the present study was to determine the association between serum 25 hydroxyvitamin D concentration and components of metabolic syndrome in Iranian adult women.

Patients and Methods: This study was comprised of 156 women aged ≥ 30 years with at least three of the five criteria of metabolic syndromes. Serum 25-hydroxyvitamin D (25 (OH)D) levels and components of the metabolic syndrome were determined. Metabolic syndrome was defined according to NCEP/ATP III criteria. The subjects were classified into three groups according to their serum concentration of 25 (OH)D.

Results: Mean of serum concentration of 25-hydroxyvitamin D was 20.5 ± 10.8 ng/mL with %54.5, 23.1% and 22.4% of subjects were deficient, insufficient and sufficient in vitamin D, respectively. After adjustment for age, BMI, physical activity, and ANCOVA, fasting blood sugar concentration was shown to be inversely associated with serum 25 (OH)D ($P = 0.004$). On the other hand HDL-C showed significant correlation across different groups of vitamin D status ($P = 0.014$). Waist Circumference had favorable changes, without any statistically significant correlation. Also no significant association was observed between other components of metabolic syndrome and 25 (OH)D in different groups.

Conclusions: The components of metabolic syndrome are influenced by serum 25 (OH)D concentrations. The finding of this investigation revealed that FBS and HDL-C concentrations related to serum 25 (OH)D. Therefore, further longitudinal studies and randomized clinical trials are necessary to determine the possible role of vitamin D in prevention of diabetes and cardiovascular disease.

Keywords: Blood Fasting Sugar; Women; Vitamin D; Metabolic Syndrome

1. Background

Vitamin D deficiency is a common health problem (1). The prevalence of hypovitaminosis D is increasing worldwide (2, 3), and is more dramatic in Asian countries (4). According to different studies the prevalence of vitamin D deficiency in Iran varied from 44.8% to 79.6% (5). Studies suggest that nearly 30–50% of adult populations are at risk of vitamin D deficiency (6). However, it is increasing with aging as, the prevalence of vitamin D deficiency in elderly has been reported about 50% worldwide (7). The natural food sources of vitamin D are limited. Much of vitamin D is produced through direct exposure to sunlight. On the other hand, the latitude and season affect the cutaneous synthesis of vitamin D, the extent of which is less known (8). Moreover, vitamin D deficiency can be caused by inadequate vitamin D intake and absorption, obesity, dark skin, aging and decline in physical activity (9, 10). Although, the main role of vitamin D is regulating calcium homeostasis and bone health, it is now known that vitamin D deficiency

is associated with insulin resistance, metabolic syndrome, hypertension, diabetes, cancer and autoimmune diseases (11). Several recent reviews have reported associations between low levels of 25 (OH)D and components of metabolic syndrome (MetS) (12-14). The guidelines of the National Cholesterol Education Program/Adult Treatment Panel III (NCEP/ATP III) MetS indicate the presence of at least three of the following five criteria. These include fasting blood sugar ≥ 110 mg/dL, TG ≥ 150 mg/dL, low serum HDL-C (< 40 mg/dL (males) or < 50 mg/dL (females), waist circumference > 102 cm in males and > 88 cm in females, blood pressure $\geq 130/85$ mmHg or using antihypertensive drugs (15, 16).

Because of the differences in definitions and populations studied, precise prevalence of MetS is unknown (17). However, data indicate that the prevalence of this syndrome among adult Iranian population aged ≥ 20 years is higher than those in most developed countries

Implication for health policy/practice/research/medical education

This paper discusses about Vitamin D role in the development of metabolic syndrome

Copyright © 2014, Health Policy Research Center, Shiraz University of Medical Sciences This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

(5). Individuals with MetS are at greater risk of coronary vascular disease (CVD), coronary heart disease (CHD) and diabetes (18). Physical inactivity, aging, obesity especially the central type and unhealthy diet influence the development of MetS and related diseases. Vitamin D status may be regarded as a novel possible risk factor for MetS and relevant diseases (19, 20). Acceptable biochemical criteria for vitamin D are still a matter of debate. However, most experts have defined vitamin D deficiency as 25 (OH)D level lower than 20 ng/mL, and insufficiency as it is 21-29.99 ng/mL, the optimal concentration of 25 (OH) D is at least 30 ng/mL (21, 22). Epidemiological studies have shown different results. A cross-sectional study on 45 year old subjects, reported that serum 25 (OH)D was associated with all features of MetS (23). But, other studies showed a relationship between 25 (OH)D and fasting glucose or blood pressure and triglyceride (7, 12) Third National Health and Nutrition Examination Survey have reported an inverse association between 25 (OH)D and hyperglycemia, and abdominal obesity (24). Despite the increasing prevalence of metabolic syndrome and vitamin D deficiency among Iranian population, especially women, limited studies have been conducted to explore this condition. Thus, it is seemed necessary to carry out an investigation to establish the association between metabolic syndrome and vitamin D deficiency in the foregoing populations and to delineate this relationship. Our hypothesis is that serum 25 (OH)D levels are associated with components of MetS.

2. Objectives

Therefore, in this study we evaluated the relationship between serum 25 (OH)D and component of the metabolic syndrome among Iranian women with MetS.

3. Patients and Methods

3.1. Subjects

Based on the underlying sample size Equation 1:

$$\text{Equation 1. } \frac{\left(z_{1-\frac{\alpha}{2}} + z_{1-\beta}\right)^2 \times 4}{\left(\ln \frac{1+r}{1-r}\right)^2} + 3 = 156$$

$Z_{1-\alpha/2}$, Z for Type I error = 1.96; $Z_{1-\beta}$, Z for Type II error = 0.84; r: correlation coefficient = 0.22. 156 women with MetS were needed for adequate power. Thus, in this cross-sectional study 156 women aged ≥ 30 years were selected by consecutive random sampling. Obese women with high blood pressure referred to us by physicians in health centers. Women consuming vitamin D and/or calcium supplements, as well as drugs affecting the metabolism of vitamin D during past 3 to 4 months, or having a history of renal or hepatic disease, overt diabetes and malignancies and those with at least three criteria for metabolic syn-

dromes, as defined by NCEP/ATP III, were excluded from the study. Subjects were categorized into three groups based on serum concentrations of 25 (OH)D. The first category included: women with a serum concentration of 25 (OH)D < 20 ng/mL which were considered as vitamin D deficient. The second group had 25 (OH)D equal to 20- 29.99 ng/mL regarded as insufficient, and third group with 25 (OH)D concentration of ≥ 30 ng/mL as vitamin D sufficient one (18). Sampling was conducted in the summer after filling demographic questionnaire.

3.2. Laboratory Measurement

Blood samples were collected after an overnight fasting of 12-h, and serum was promptly separated by centrifugation and kept at -20°C until used. After all the blood samples were taken, fasting glucose and lipid profile were measured by routine enzymatic methods (Pars Azmoon kits; Tehran, Iran) using an autoanalyzer (Tokyo Boeki Prestige 24, Japan). Serum 25 (OH)D concentration was determined by radioimmunoassay ELISA (Euroimmune, Germany).

3.3. Assessment of Dietary Intake

Usual dietary intake was assessed using a validated 168-item semi quantitative food-frequency questionnaire (FFQ) (25). Which evaluated dietary intake during the past year, based on their consumption frequency, weekly and monthly? That questionnaire included a list of foods with serving (26). Each food and beverage was then coded according to the prescribed protocol and analyzed for content of energy and the other nutrients using N4 nutritional software designed for Iranian foods. The mean dietary vitamin D was obtained. All FFQs were processed by a trained dietitian.

3.4. Assessment of Other Variables

Waist circumference (WC) was measured at the narrowest level between the lowest rib and the iliac crest by a nonstretch tape at the end of a normal expiration. Body weight was measured by Seca digital scale with light clothes and without shoes to the nearest 100 gr. Height was measured by non-stretch tape to the nearest 0.1 cm with bare feet while the shoulders were in a normal position body mass index (BMI) was calculated as weight in kilograms divided by the square of height in meters. Data were obtained by an interview based on International Physical Activity Questionnaire and expressed as MET hours per day (MET-h/day) (27). Body weight was measured by Seca digital scale with light clothes with bare feet to the nearest 100 g. Height was measured by non nonstretch tape to the nearest 0.1 cm while not wearing shoes and the shoulders were in a normal position body mass index (BMI) was calculated as weight (in kilograms) divided on the square of height (in meters). Blood pressure was measured two times waiting 15-minutes between measurements using a mercury sphygmoma-

nometer on the right hand in a sitting position. MAP was calculated as $[(2 \times \text{diastolic}) + \text{systolic}] / 3$ (28).

3.5. Statistical Analysis

We analyzed all data by SPSS software 16. We used ANOVA for comparing variables between groups. However, because of significant correlation between HDL-C and triglyceride as two components of metabolic syndrome, we introduced ANCOVA for comparing their relationship to Vitamin D status by adjusting confounding factors. Kolmogorov-smirnov was used to evaluate normal distribution of data. All variables reported as Means \pm SD and values less than 0.05 were considered statistically significant.

4. Results

Of 156 women participating in this study, those with at least three of the five specific criteria for metabolic syndrome, which was prepared according to NCEP/ATP III, were included in this investigation. The general characteristics of subjects are shown in Table 1.

Subjects were categorized into three groups based on serum concentration of 25 (OH)D. Eighty five subjects (54.5%) with 25 (OH)D < 20 ng/mL were vitamin D deficient. Thirty-six (23.1%) had 25 (OH)D = 20-29.99 ng/mL that were insufficient and and thirty-five (22.4%) women had sufficient 25 (OH)D (≥ 30 ng/mL). The Kolmogorov-Smirnov test was used for testing the normality of all variables. We compared metabolic syndrome components between different groups of is vitamin D status by using ANCOVA (Table 2).

ANCOVA showed an inverse association between fasting blood sugar with serum 25 (OH)D ($P = 0.004$). On the other hand, MANCOVA revealed significant correlation between HDL-C across different groups of vitamin D status ($P = 0.014$). Waist circumference had favorable changes, but with no statistically significant correlation. Regarding components of metabolic syndrome, no significant differences were observed across 25 (OH) D groups.

5. Discussion

In this study, more than fifty percent of the participants had serum 25 (OH)D levels below 20 ng/mL. Our findings indicated that some components of MetS are significantly associated with serum 25 (OH)D concentrations. FBS concentrations was inversely associated with increasing serum 25 (OH)D concentration ($P = 0.004$). Also HDL-C was positively correlated with 25 (OH)D concentrations ($P = 0.014$, a relationship independent from confounders). Also, WC had favorable changes across different vitamin D. status groups, but with no significant correlation. Furthermore, no significant relationship was observed between the mean blood pressure and triglyceride across 25 (OH)D concentrations. The results of several studies,

Table 1. Anthropometric, Biochemical, and Dietary Characteristics of Subjects (n = 156)^{a,b}

Variables	Results
Age, y	46.1 \pm 9.5
BMI, kg/m ²	34 \pm 5.3
WC, cm	106.9 \pm 9.9
MAP, mmHg	10.2 \pm 0.9
Physical activity index, MET-h/day	33 \pm 2.7
Diet	-
Total energy, kcal/d	2444 \pm 830
Protein, g/d	84 \pm 31.4
Carbohydrate, g/d	369 \pm 122
Fat, g/d	74.1 \pm 47.4
Dietary Vitamin D, μ g/d	1.1 \pm 1.3
Biochemical	-
Triglyceride, mg/dL	176 \pm 66.1
Fasting plasma glucose, mg/dL	115.7 \pm 30.2
HDL-C, mg/dL	50.4 \pm 7.7
25 (OH)D, ng/mL	20.5 \pm 10.8

^a Abbreviations: BMI, body mass index; HDL, high density lipoprotein; MAP, mean arterial pressure; MET, metabolic equivalent; WC, waist circumference.

^b All variables reported as Means \pm SD.

Table 2. Comparison of the Metabolic Syndrome Components Across Different Groups of Vitamin D Status by Analysis of Variance^{a,b,c}

	Deficient (12.25 \pm 3.79) (n = 85)	Insufficient (24.55 \pm 3.39) (n = 36)	Sufficient (36.71 \pm 5.6) (n = 35)	P Value
Triglyceride, mg/dL	188.27 \pm 87.63	182.25 \pm 67.10	162.57 \pm 48.063	0.121 ^d
WC, cm	107.65 \pm 10.266	107.63 \pm 10.62	104.31 \pm 7.64	0.142 ^e
FBS, mg/dL	127.28 \pm 33.72	113.22 \pm 47.08	97.22 \pm 14.79	0.004 ^e
HDL-C, mg/dL	49.35 \pm 8.67	50.44 \pm 7.11	54.228 \pm 7.48	0.014 ^d
MAP, mmHg	10.231 \pm 1.144	10.184 \pm 0.89	10.247 \pm 0.79	0.66 ^e

^a Abbreviations: FBS, fastin blood sugar; HDL-C, high density lipoprotein.

^b Deficient, 25 (OH)D < 20 ng/mL; Insufficient, 20-29.99 ng/mL; Sufficient, ≥ 30 ng/ml.

^c All variables reported as Mean \pm SD. Controlled by age, BMI, PAL.

^d P is obtained by analysis of covariance (ANCOVA).

^e P is obtained by multiple analysis of covariance (MANCOVA).

although inconsistent, have reported increasing evidence about relationships between serum 25 (OH)D and components of MetS (4, 21, 23). Our findings of the relationship between 25 (OH)D and glucose homeostasis are in agreement with those of Brock et al. (12). The relationship between 25 (OH)D and MetS may be approached mechanisms including: 1) increasing the secretion and insulin sensitivity; 2) stimulated expression of insulin receptor; 3) enhancement of glucose transport into the cells of target tissues 4) and regulation of intracellular calcium (20, 23). Respecting the relationship of blood glucose to 25 (OH)D indicated in most epidemiological studies, it is necessary to design randomized clinical trials in order to determine the therapeutic dose of vitamin D for glycemic control, particularly in patients with type 2 diabetes. The association between waist circumference and vitamin D might be due to: 1) solubility of vitamin D in adipose tissue, 2) slow bioavailability of circulating vitamin D and 3) inadequate intake of vitamin D due to low physical activity in obese subjects and lesser exposure to sunlight (12).

In contrast with most studies, our study showed an inverse and non-significant relationship between WC and 25 (OH)D concentrations. This may be due to our small sample size in insufficient and sufficient categories of 25 (OH)D vitamin groups. However, HDL-C concentrations showed a positive and significant positive relationship with 25 (OH)D levels. Our findings were in agreement with those of Brock et al. study, in that vitamin D deficiency was not associated with central obesity (12), but this relationship was significant in studies reported by Maki et al. who used a larger sample size (21). We found no relationship between dietary vitamin D and markers of the Mets, and also observed no association between 25 (OH)D and dietary vitamin D. Such negative relationships may be explained by the absence of correlation, the limited food sources of vitamin D and measuring error in dietary assessment of vitamin D by FFQ. However, fewer studies have been conducted on the effect of sunlight on the synthesis of vitamin D in skin. Thus, the questionnaire used was designed depending on the latitude of different areas.

The present study showed that sun exposure was not associated with levels of 25 (OH)D, a finding in agreement with Hashemipour et al. study on Tehranian adults (4). Possible reasons might be either due to imprecise oral answers to questions regarding exposure period to sunlight or the type of clothes worn by the subjects. The cutaneous synthesis of vitamin D levels are influenced by latitude, aging, season, the different intensity of sunlight at different times of the day. It is therefore necessary to consider such entities in preparing a valid questionnaire. Studies have shown that, vitamin D deficiency may be associated with negligible physical activity (10). Approximately, the strategies for activity should be considered for at least 55% of subjects. We found no significant relationship between blood pressure and vitamin D status. Similarly, Forouhi et al. (27) did not find any such association after 10 years

follow up and pointed out that association of blood markers of vitamin D with blood pressure was complex. Several studies have shown that vitamin D may regulate blood pressure by regulating the renin-angiotensin system and inhibits the renin mRNA expression (29, 30). But, most of the results in this area are conflicting. Our results are in contrast with those of some studies (7, 13) that found statistically significant inverse association between 25 (OH) D with blood pressure. According to these studies, significant relationship could be attributed to using predicted vitamin D levels, self-reported hypertension and using repeated measure analyses. There is little evidence to suggest that vitamin D status can affect the development of dyslipidemia, an area deserving further studies. Our study have some limitations including small sample size, cross sectional design, single sex, using questionnaire lacking accurate exposure time to sunlight and precise assessment of vitamin D intake. In conclusion, vitamin D deficiency may have an important role in MetS components. Therefore, further longitudinal studies and randomized clinical trials are necessary to examine the association between vitamin D deficiency and features of MetS.

Authors' Contribution

Zamzam Paknahad has designed the study, written and edited manuscript; Azam Ahmadi Vasmehjani has done data collection, references search; Mohammad Reza Maracy has contributed in statistical analysis. All authors read and approved the final manuscript. Thanks to all authors for their support and help in this study

Funding/Support

This work was fully financially supported by Isfahan University of Medical Sciences, IR Iran

References

1. Kositsawat J, Freeman VL, Gerber BS, Geraci S. Association of A1C levels with vitamin D status in U.S. adults: data from the National Health and Nutrition Examination Survey. *Diabetes Care*. 2010;**33**(6):1236-8.
2. Chiu KC, Chu A, Go VL, Saad MF. Hypovitaminosis D is associated with insulin resistance and beta cell dysfunction. *Am J Clin Nutr*. 2004;**79**(5):820-5.
3. Makariou S, Liberopoulos E, Florentin M, Lagos K, Gazi I, Challa A, et al. The relationship of vitamin D with non-traditional risk factors for cardiovascular disease in subjects with metabolic syndrome. *Arch Med Sci*. 2012;**8**(3):437-43.
4. Hashemipour S, Larijani B, Adibi H, Javadi E, Sedaghat M, Pajouhi M, et al. Vitamin D deficiency and causative factors in the population of Tehran. *BMC Public Health*. 2004;**4**:38.
5. Hashemipour S, Sarokhani M, Asefzadeh S, Mehrtash B, Sheikholeslami H, Ghoddosi A. Effect of Different Dosage of Intramuscular Vitamin D3 on Serum 25 (OH) D. *Iran J Endocrinol Metab*. 2008;**10**(2):115-20.
6. Parker J, Hashmi O, Dutton D, Mavrodaris A, Stranges S, Kandala NB, et al. Levels of vitamin D and cardiometabolic disorders: systematic review and meta-analysis. *Maturitas*. 2010;**65**(3):225-36.
7. Park HY, Lim YH, Kim JH, Bae S, Oh SY, Hong YC. Association of serum 25-hydroxyvitamin D levels with markers for metabolic syndrome in the elderly: a repeated measure analysis. *J Korean Med Sci*. 2012;**27**(6):653-60.

8. Webb AR, Kline L, Holick MF. Influence of season and latitude on the cutaneous synthesis of vitamin D3: exposure to winter sunlight in Boston and Edmonton will not promote vitamin D3 synthesis in human skin. *J Clin Endocrinol Metab*. 1988;**67**(2):373-8.
9. Liu E, Meigs JB, Pittas AG, McKeown NM, Economos CD, Booth SL, et al. Plasma 25-hydroxyvitamin d is associated with markers of the insulin resistant phenotype in nondiabetic adults. *J Nutr*. 2009;**139**(2):329-34.
10. Wicherts IS, van Schoor NM, Boeke AJ, Visser M, Deeg DJ, Smit J, et al. Vitamin D status predicts physical performance and its decline in older persons. *J Clin Endocrinol Metab*. 2007;**92**(6):2058-65.
11. Riachy R, Vandewalle B, Kerr Conte J, Moerman E, Sacchetti P, Lukowiak B, et al. 1,25-dihydroxyvitamin D3 protects RINm5F and human islet cells against cytokine-induced apoptosis: implication of the antiapoptotic protein A20. *Endocrinology*. 2002;**143**(12):4809-19.
12. Brock K, Ke L, Koo F, Jang H, Clemson L. Vitamin D and Metabolic Syndrome in Immigrant East Asian Women Living in Sydney, Australia: A Pilot. *J Metab Syndr*. 2012.
13. Burgaz A, Orsini N, Larsson SC, Wolk A. Blood 25-hydroxyvitamin D concentration and hypertension: a meta-analysis. *J Hypertens*. 2011;**29**(4):636-45.
14. Gagnon C, Lu ZX, Magliano DJ, Dunstan DW, Shaw JE, Zimmet PZ, et al. Serum 25-hydroxyvitamin D, calcium intake, and risk of type 2 diabetes after 5 years: results from a national, population-based prospective study (the Australian Diabetes, Obesity and Lifestyle study). *Diabetes Care*. 2011;**34**(5):1133-8.
15. Gulseth HL, Gjelstad IM, Tierney AC, Lovegrove JA, Defoort C, Blaak EE, et al. Serum vitamin D concentration does not predict insulin action or secretion in European subjects with the metabolic syndrome. *Diabetes Care*. 2010;**33**(4):923-5.
16. Pereira MA, Jacobs DR Jr, Van Horn L, Slattery ML, Kartashov AI, Ludwig DS. Dairy consumption, obesity, and the insulin resistance syndrome in young adults: the CARDIA Study. *JAMA*. 2002;**287**(16):2081-9.
17. Ford ES, Giles WH, Dietz WH. Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA*. 2002;**287**(3):356-9.
18. Wannamethee SG, Shaper AG, Lennon L, Morris RW. Metabolic syndrome vs Framingham Risk Score for prediction of coronary heart disease, stroke, and type 2 diabetes mellitus. *Arch Intern Med*. 2005;**165**(22):2644-50.
19. Bartke A. Insulin and aging. *Cell Cycle*. 2008;**7**(21):3338-43.
20. Anderssen SA, Carroll S, Urdal P, Holme I. Combined diet and exercise intervention reverses the metabolic syndrome in middle-aged males: results from the Oslo Diet and Exercise Study. *Scand J Med Sci Sports*. 2007;**17**(6):687-95.
21. Maki KC, Rubin MR, Wong LG, McManus JF, Jensen CD, Marshall JW, et al. Serum 25-hydroxyvitamin D is independently associated with high-density lipoprotein cholesterol and the metabolic syndrome in men and women. *J Clin Lipidol*. 2009;**3**(4):289-96.
22. Dobnig H, Pilz S, Scharnagl H, Renner W, Seelhorst U, Wellnitz B, et al. Independent association of low serum 25-hydroxyvitamin d and 1,25-dihydroxyvitamin d levels with all-cause and cardiovascular mortality. *Arch Intern Med*. 2008;**168**(12):1340-9.
23. Hypponen E, Boucher BJ, Berry DJ, Power C. 25-hydroxyvitamin D, IGF-1, and metabolic syndrome at 45 years of age: a cross-sectional study in the 1958 British Birth Cohort. *Diabetes*. 2008;**57**(2):298-305.
24. Hosseini Nezhad A, Khoshniat Nikoo M, Maghbooli Z, Karimi F, Mirzaei K, Hosseini A, et al. Relationship between serum vitamin D concentration and metabolic syndrome among Iranian adults population. *Daru*. 2009;**17**(Suppl 1):1-5.
25. Esfahani FH, Asghari G, Mirmiran P, Azizi F. Reproducibility and relative validity of food group intake in a food frequency questionnaire developed for the Tehran Lipid and Glucose Study. *J Epidemiol*. 2010;**20**(2):150-8.
26. Ghafarpour M, Houshiar Rad A, Kianfar H. [The manual for household measures, cooking yields factors and edible portion of food]. Tehran: Nashre Olume Keshavarzy; 1999.
27. Forouhi NG, Luan J, Cooper A, Boucher BJ, Wareham NJ. Baseline serum 25-hydroxy vitamin d is predictive of future glycemic status and insulin resistance: the Medical Research Council Ely Prospective Study 1990-2000. *Diabetes*. 2008;**57**(10):2619-25.
28. Mallion JM, Hamici L, Chatellier G, Lang T, Plouin PF, De Gaudemaris R. Isolated systolic hypertension: data on a cohort of young subjects from a French working population (IHPAF). *J Hum Hypertens*. 2003;**17**(2):93-100.
29. Ford ES, Ajani UA, McGuire LC, Liu S. Concentrations of serum vitamin D and the metabolic syndrome among U.S. adults. *Diabetes Care*. 2005;**28**(5):1228-30.
30. Pfeifer M, Begerow B, Minne HW, Nachtigall D, Hansen C. Effects of a short-term vitamin D(3) and calcium supplementation on blood pressure and parathyroid hormone levels in elderly women. *J Clin Endocrinol Metab*. 2001;**86**(4):1633-7.