

Chemical composition and antifungal effect of hydroalcoholic extract of *Allium tripedale* (Tvautv.) against *Candida* species

Shirani M¹, Samimi A^{1*}, Kalantari H¹, Madani M², Kord Zanganeh A³

¹ Department of Pharmacology and Toxicology, School of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

² Department of Microbiology, Falavarjan Branch, Islamic Azad University, Isfahan, Iran

³ Department of Pharmaceutics and Nanotechnology Research Center, School of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Article Info

Article type:
Original article

Article History:

Received: 07 June 2017

Revised: 31 July 2017

Accepted: 05 August 2017

* Corresponding author:

Azin Samimi

Department of Pharmacology and Toxicology, Pharmacy School, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

Email: Azin.samimi831@gmail.com

ABSTRACT

Background and Purpose: Treatment of life-threatening fungal infections caused by *Candida* species has become a major problem. *Candida* spp. are the most important causative agents of candidiasis. *Allium tripedale* is a medicinal plant that has been traditionally used to treat infections. In the present study, we aimed to determine the chemical compounds and antimicrobial activity of hydroalcoholic extract of *A. tripedale* against different species of *Candida*.

Materials and Methods: Phytochemical analysis was performed to identify the possible bioactive components of this extract by using gas chromatography and mass spectroscopy (GC-MS). The hydroalcoholic extract of *A. tripedale* were collected. Different concentrations of *A. tripedale* (50, 25, 12.5, and 6.25 mg/ml) were used to evaluate its antifungal activity against *Candida* species (*C. albicans*, *C. parapsilosis*, and *C. krusei*) using disk diffusion assay.

Results: The GC-MS analysis revealed the presence of 40 different phytoconstituents with peak area; the major compounds were tetracosane, hexadecanoic acid, 1-eicosanol, 1,2-dihydro-pyrido[3,2,1-kl]phenothiazin-3-one, 2-hexadecen-1-ol, and 3,7,11,15-tetramethyl. Hydroalcoholic extract showed strong antimicrobial activity (inhibition zone ≥ 20 mm), moderate antimicrobial activity (inhibition zone < 12 -20 mm), and no inhibition (zone < 12 mm). In addition, the hydroalcoholic extract exhibited the highest antimicrobial properties against *C. albicans* strains.

Conclusion: *A. tripedale* extract had a considerable inhibitory effect against various *Candida* species, but its highest inhibitory effect was against *Candida albicans*. Further investigations are required to detect the performance of this plant in the treatment of *Candida* infection.

Keywords: *Allium tripedale*, *Candida* species, Candidiasis, GC-MS

➤ How to cite this paper

Shirani M, Samimi A, Kalantari H, Madani M, Kord Zanganeh A. Chemical composition and antifungal effect of hydroalcoholic extract of *Allium tripedale* (Tvautv.) against *Candida* isolates. *Curr Med Mycol*. 2017; 3(1): 6-12. DOI: [10.18869/acadpub.cmm.3.1.6](https://doi.org/10.18869/acadpub.cmm.3.1.6)

Introduction

Plants are a great source of useful phytochemicals, which have inhibitory effects against some microorganisms in vitro and are effective in the treatment of various conditions [1]. Generally, 1-10% of plants (out of approximately 250,000-500,000 species) on earth are used by humans [2]. In recent years, there has been a growing global interest in the use of medicinal plants for disease prevention and treatment, especially in Iran [3]. Limited success in the treatment of human diseases, undesirable side effects of chemical drugs, and growing emergence of drug resistance, particularly to antibiotics, have led to increased use of medicinal plants [4].

Medicinal plants are a widespread source of biologically active compounds including alkaloids,

tannins, flavonoids, and phenolic compounds. Accordingly, they are of marked significance to the health of individuals and communities and are widely used for disease treatment [2].

A. tripedale belonging to the Liliaceae family, is a wild *Allium* species native to the Caucasus (North + South), Iraq, Turkey, and Iran. This plant has long and strong stems (50-90 cm in length) and some what unpleasant taste [5, 6]. *A. tripedale* has been extensively used by locals as a spicy vegetable and for the treatment of infections. Given the presence of saponins in the structure of this plant, it is expected to have inhibitory effect against pathogenic fungi [7].

Since the early 1990s, the increase in the number of infections caused by pathogenic and opportunistic fungi has been introduced as the leading cause of

mortality among hospitalized patients [8]. In other words, a large number of people are suffering from fungal infections, and these infections are posing a great threat to mankind [9]. In addition, the increased use of antifungal agents has led to the development of resistance to the available drugs.

Candida albicans as an opportunistic pathogen plays an important role in the infection and is the most common cause of cutaneous, oral, and systemic diseases in immunodeficiency patients [10]. Although *Candida albicans* is still the major species isolated from clinical samples in the majority of individuals, it is well known that some other non-*albicans* *Candida* spp. such as *Candida glabrata*, *Candida krusei*, *Candida parapsilosis*, and *Candida tropicalis* infections are significantly widespread. Candidiasis associated with this kind of non-*albicans* *Candida* spp. pose a clinical challenge because they are resistant to common antifungal agents such as fluconazole and amphotericin B [11, 12].

Regarding the increase in the use of antifungal agents and resistance to some types of *Candida* spp. and the undesirable side effects of chemical drugs, it is essential to explore new sources of treatment, particularly among herbal plants [8, 13]. To the best of our knowledge, no has yet explored the antifungal activity of *A. tripedale* against *Candida* isolates. The purpose of this study was to evaluate chemical composition and antimicrobial activity of hydroalcoholic extract of *A. tripedale* against different *Candida* spp.

Materials and Methods

Plant collection

Tripedale was collected from the highlands of Shahrekord in southeast of Iran (Isfahan Province). The collected samples were identified in Ahvaz Agricultural and Natural Research Centre (Herbarium No. A151640100AP). Extraction and laboratory examinations were carried out in Ahvaz University of Medical Sciences, Ahvaz, Iran. The aerial parts of the plants were aired indoors at room temperature and then finely powdered using an electric grinder (Busch, MKM6003, Slovenia). It took two days to extract 20 g of plant materials by soxhlet with 120 ml ethanol 80%. The extract was filtered using Whatman qualitative filter paper, Grade 1. The extract was preserved in sterilized airtight bottles at 4°C, and then to prepare the dried extracts, the solution was placed in a bain-marie at 40°C for 24 h prior to use [14].

Gas chromatography and mass spectroscopy (GC-MS) analysis

GC-MS analysis of ethanolic extract of the whole *A. tripedale* was performed on GC 7890A equipped with MS 5975C detector and HP-5ms capillary column (30 × 0.25 m, 0.25 μm; Agilent Co., USA). The initial column temperature was set at 60°C, then increased from 60°C to 190°C (heating rate: 5°C per

minute), from 190°C to 270°C for 30 min, and finally kept at 270°C for approximately 5 min; the total analysis time was about 34 min.

Compound identification

Interpretation of GC-MS was performed via the National Institute Standard and Technology (NIST) database. The spectra of the unknown components were compared with those of the known ones registered in the NIST library. The name, molecular weight, and structure of the components of the test materials were determined.

Preparation of organisms

Standard strains of *C. albicans* (ATCC 3153), *C. parapsilosis* (ATCC 2195), and *C. krusei* (ATCC 573) were obtained from the Department of Mycology, Faculty of Veterinary Medicine, Tehran University, Tehran, Iran. The strains were cultured on Sabouraud Dextrose Agar (SDA) (Merck, Germany) medium. Fungal suspension was prepared with concentration adjusted to 1.5×10^6 CFU/ml in sterile distilled water as described by Forbes et al. [15].

Well diffusion assay

Agar well diffusion method is extensively used to evaluate the antimicrobial activity of plants or microbial extracts. To determine the effective concentration, inhibition zones of hydroalcoholic extract of *A. tripedale* were examined against *C. albicans* (ATCC 3153), *C. parapsilosis* (ATCC 2195), and *C. krusei* (ATCC 573) strains by using well assay technique. The agar plate surface was inoculated overnight by spreading inoculum of *Candida* spp. over the entire SDA surface. A hole 6 to 8 mm in diameter was punched with a sterile tip. Then, the extract was added to the pits in the agar medium and incubated under suitable conditions at 27°C for 24 h [16]. The diameter of the inhibitory zone was measured, and the corresponding effective concentration was chosen for subsequent experiments [17].

Disk diffusion method

The fungal broth culture aliquots were added to SDA. Sterile paper disks (Merck, Germany) were impregnated with 50 μl of extract solution and placed on the culture plates. The plates were incubated at 37°C for 24 h. Antifungal activity was evaluated by measuring the inhibition zone diameter [18]. Fluconazole was used as positive control [19], whereas paper disks loaded with solvents (ethanol and distilled water) were used as negative controls.

Statistical analysis

Statistical analysis was performed using SPSS, version 10.0. The inhibition diameters of the test substances were expressed as mean and standard deviation. Group comparisons were performed using One-way analysis of variance (ANOVA) followed by Waller-Duncan Post Hoc test. *P*-value less than 0.05

was considered statistically significant.

Results

The phytochemical analysis

The phytochemicals present in the hydroalcoholic extract of *A. tripedale* were identified by GC-MS analysis; GC-MS running time is 34 min. The active compounds in the hydroalcoholic extract of the plant, their retention time (RT), molecular formula, and molecular weight are provided in Table 1, and GC-MS chromatograms are presented in Figure 1.

The gas chromatogram is used to help identify a mixture of compounds by separating compounds according to each compound's retention time. The heights of the peaks indicate the relative concentrations of the components present in the plant. GC-MS

analysis revealed the presence of 40 compounds by dichloromethane solvent; the major compounds included tetracosane, hexadecanoic acid, 1-eicosanol, 1,2-dihydro-pyrido[3,2,1-kl]phenothiazin-3-one, 2-hexadecen-1-ol, and 3,7,11,15-tetramethyl.

Disk and well diffusion assay

Preliminary screening of the antifungal activity of hydroalcoholic extracts of *A. tripedale* was performed against *Candida* spp. using the disk and well diffusion assay. The results showed variation in the antifungal properties of hydroalcoholic extract of *A. tripedale* (Table 2). The extract showed strong activity (inhibition zone ≥ 20 mm), moderate activity (inhibition zone < 12 -20 mm), and no inhibition (zone < 12 mm). Fluconazole, a known antifungal antibiotic, as a positive

Table 1. Phytochemicals identified in the hydroalcoholic extract of *A. tripedale* by gas chromatography and mass spectroscopy

S.NO.	ID	RT	Area%	CAS	Molecular formula	Molecular weight g/mol
1	2-Pentene, 2-methyl	3.133	0.2	625-27-4	CH ₃ CH ₂ CH=C(CH ₃) ₂	84.16
2	Furan, 2,4-dimethyl-	7.624	0.18	3710-43-8	C ₆ H ₈ O	96.1271
3	Ethane, 1,1,2,2-tetrachloro-	9.324	0.72	79-34-5	C ₂ H ₂ Cl ₄	167.8493
4	Benzaldehyde	11.435	0.19	100-52-7	C ₇ H ₆ O	106.1219
5	2,4 HEPTADIENAL	11.801	1.70	4313-03-5	C ₇ H ₁₀ O	110.1537
6	Nonanal	13.924	0.5	124-19-6	C ₉ H ₁₈ O	142.2386
7	Benzeneacetaldehyde	14.193	0.42	122-78-1	C ₈ H ₈ O	120.1485
8	2-Fluorophenylhydrazine	14.908	0.13	2368-80-1	C ₆ H ₇ FN ₂	126.1316
9	Octanoic Acid	15.441	0.35	124-07-2	C ₁₆ H ₃₀ O ₄ Sn	405.117
10	Benzoic acid	1.654	0.24	65-85-0	C ₇ H ₆ O ₂	122.1224
11	Pyridine, 3-(phenylazo)-	17.581	0.18	2569-55-3	C ₁₁ H ₉ N ₃	183.21
12	4-Pyridinamine, N-methyl-N,3-dinitro-	18.73	0.14	104503-82-4	C ₆ H ₆ N ₄ O ₄	198.13624
13	3-Methyl-2,3-dihydro-benzofuran	18.347	0.34	13524-73-7	C ₉ H ₁₀ O	134.1751
14	1-Carboxymethyl-2(1H)-pyridone	19.440	0.29	56546-36-2	C ₇ H ₇ NO ₃	153.1354
15	.alpha.-(Aminomethylene)glutaconicanhydride	20.12	0.15	67598-07-6	C ₆ H ₅ NO ₃	139.1088
16	2,4-Decadienal	20.093	0.43	2363-88-4	C ₁₀ H ₁₆ O	152.23344
17	2-Bromo-5-(hydroxymethyl)pyridine	20.178	0.12	122306-01-8	C ₆ H ₆ BrNO	188.02194
18	(E,Z,Z)-2,4,7-Tridecatrinal	20.310	0.67	1000314-35-6	C ₁₃ H ₂₀ O	192.3016
19	Hexadecane, 7,9-dimethyl-	20.836	0.74	21164-95-4	C ₁₈ H ₃₈	254.50
20	2,4-Decadienal	20.934	0.64	2363-88-4	C ₁₀ H ₁₆ O	152.2334
21	2-Methoxy-4-vinylphenol	21.763	0.6	7786-61-0	C ₉ H ₁₀ O ₂	150.1745
22	2-Propenoic acid, 3-phenyl-	24.910	0.41	621-82-9	C ₉ H ₈ O ₂	148.1586
23	2(4H)-Benzofuranone, 5,6,7,7a-tetrahydro-4,4,7a-trimethyl-, (R)-	28.687	0.25	17092-92-1	C ₁₁ H ₁₆ O ₂	180.24
24	Tetradecanal	29.265	0.26	124-25-4	C ₁₁ H ₂₀ O ₂	184.2753
25	1,1-Difluoro-2-methyl-3-ethyl cyclopropane	30.032	0.22	1000144-82-1	C ₆ H ₁₀ F ₂	120.140406
26	Cyclodecane	30.140	0.19	293-96-9	C ₁₀ H ₂₀	140.27
27	Tetradecanoic acid	30.243	0.65	544-63-8	C ₁₄ H ₂₈ O ₂	228.37
28	2-Hexadecen-1-ol, 3,7,11,15-tetramethyl-, [R-[R*,R*-(E)]]-	30.306	4.67	150-86-7	C ₂₀ H ₄₀ O	296.531
29	Methyl .beta.-d-galactopyranoside	30.524	0.48	1000126-04-6	C ₇ H ₁₄ O ₆	194.18246
30	3-Pyridinamine, N-methyl-2-nitro-	30.655	0.15	32605-06-4	C ₆ H ₇ N ₃ O ₂	153.1387
31	2-Pentadecanone, 6,10,14-trimethyl	31.565	0.85	502-69-2	C ₁₈ H ₃₆ O	268.4778
32	Hexadecanoic acid	340380	6.91	57-10-3	C ₁₆ H ₃₂ O ₂	256.42
33	Phthalic acid, butyl undecyl ester	34529	0.92	1000308-91-2	C ₂₃ H ₃₆ O ₄	376.52954
34	Hexadecanoic acid, ethyl ester	34.586	0.41	628-97-7	C ₁₈ H ₃₆ O ₂	284.47724
35	Phthalic acid, propyl nonyl ester	36.641	0.41	1000309-06-4	C ₂₀ H ₃₀ O ₄	334.4498
36	Cyclohexanol, 1-methyl-4-(1-methylethyl)-	36.829	0.89	21129-27-1	C ₁₀ H ₂₀ O	156.27
37	Phthalic acid, isobutyl pent-2-en-4-yn-1-yl ester	45.761	1.56	1000315-45-6	C ₁₇ H ₁₈ O ₄	286.32242
38	1-Eicosanol	49.338	6.79	629-96-9	C ₂₀ H ₄₂ O	298.54688
39	Tetracosane	50.751	34.17	646-31-1	C ₂₄ H ₅₀	338.6538
40	1,2-Dihydropyrido(3,2,1-kl)phenothiazin-3-one	54.236	4.94	69513-42-4	C ₁₅ H ₁₁ NOS	253.31894




Figure 1. Gas chromatography and mass spectroscopy chromatogram of hydroalcoholic extract of *A. tripedale*

Table 2. Antifungal activity of the *A. tripedale* in disk and well diffusion assay

Extract concentration (mg/mL)	Zone of inhibition (mm)												
	<i>C. albicans</i>				<i>C. parapsilosis</i>				<i>C. krusei</i>				
	Disk diffusion assay		Well diffusion assay		Disk diffusion assay		Well diffusion assay		Disk diffusion assay		Well diffusion assay		
	48	72	48	72	48	72	48	72	48	72	48	72	
Hydro-alcoholic extract	50	21±0.17	10±0.0	30±0.17	10±0.28	-	-	28±0.28	28±0.34	17±0.11	10±0.25	-	-
	25	6±0.36	-	21±0.57	-	-	-	20±0.11	19±1.04	5±0.28	-	-	-
	12.5	3±.28	-	22±28	-	-	-	18±0.0	10±0.28	5±0.28	-	-	-
	6.25	-	-	19±0.20	-	-	-	-	-	-	-	-	-

control significantly inhibited the growth of *Candida* spp. (Figure 2). Based on the available evidence, the major effective antifungal activity by the hydroalcoholic extract was achieved against *C. albicans*

(Figure 3). The hydroalcoholic extract of *A. tripedale* inhibited the growth of *C. parapsilosis* by well diffusion assay and *C. krusei* by disk diffusion assay in a dose-dependent manner (Figures 4, 5).




Figure 2. Anti-fungal activity of fluconazole (50 mg/ml) against *C. albicans* by disc (A) and well (B) diffusion assay after 48 h




Figure 3. Anti-fungal activity of hydroalcoholic extract of *A. tripedale* (50 mg/ml) against *C. albicans* by disc (A) and well (B) diffusion assay after 48 h



Figure 4. Anti-fungal activity of hydroalcoholic extract of *A. tripedale* (50 mg/ml) against *C. parapsilosis* well diffusion assay after 48 h



Figure 5. Anti-fungal activity of hydroalcoholic extract of *A. tripedale* (50 mg/ml) against *C. krusei* by well diffusion assay after 48 h

Discussion

Evidence suggests that fungal infections annually affect more than a billion people, and this rate is ever increasing. *Candida* spp. can be systemic or infect different parts of the body such as skin, nails, respiratory tract, urogenital system, and alimentary canal [20]. Although several species of *Candida* are potentially pathogenic in humans, *Candida albicans* is the most important cause of severe candidiasis [21]. Trizoles initially appear to be highly effective against fungal infections, but nowadays, increased resistance is being reported and azole resistant has instigated extensive research to evaluate the effect of antifungal agents from different sources, especially medicinal plants [19]. The most famous antifungal medicinal plants belong to Liliaceae family, where more reports are found on antifungal activity of *Allium* genus [22]. The antifungal properties of the *Allium* genus were mentioned in some studies. Shams-Ghahfarokhi et al. (2007) reported that aqueous extracts of *Allium cepa* and *Allium sativum* had antifungal activity against *Malassezia furfur*, *Candida* spp. and several strains of various dermatophyte species in a dose-dependent

manner with the maximum of 100% at defined concentrations [23]. Another study by Amin and Kapadnis proved the antifungal activity of *Allium ascalonicum* against 23 fungal strains [24].

In this study, we examined the antifungal effect of hydroalcoholic extract of *A. tripedale* against different strains of *Candida* by disk and well diffusion assay. Our results revealed that the hydroalcoholic extract (50 mg/ml) had the greatest effect on *C. albicans*. However, it also had inhibitory effect against *C. parapsilosis* and *C. krusei*.

Based on the analysis conducted on the hydroalcoholic extract components using GC-MS method, 40 compounds were identified in this plant that had different properties. We found that tetracosane and other higher alkenes had antioxidant, antitumor, and antifungal properties, particularly against fungal spores and germination [25]. Tetradecanoic acid and eicosane had antioxidant and antimicrobial activities [26]. Hexadecanoic acid is known to have antioxidant and hypocholesterolemic properties and is a constituent of nematicides, pesticides, lubricants, antiandrogens, flavoring agents, hemolytics 5-alpha reductase inhibitors,

antifeedants, and insect-repellents [27].

Benzoic acid derivatives possess antibacterial and antifungal properties. Phenazopyridine hydrochloride is a topical analgesic that relieves the irritative symptoms associated with urinary tract infection through acting on the mucosal lining of the urinary tract. This agent is compatible with antibiotics and relieves pain before the antibiotic begins to control the infection. Propionic acid is an important chemical commonly used as a raw material in different industries [28]. Propionic acid, the biopreservative produced by *Propionibacterium* spp., is capable of inhibiting the growth of molds, bacteria, and dairy-spoilage yeasts such as *Zygosaccharomyces bailii* and *Candida* spp. [29]. Phenolic compounds, esters, alkanes, aldehydes, alkenes, and ketones are the major volatile compounds, which have anti-inflammatory, antiarthritic, antidiabetic, antiulcer, hypolipidemic, antiatherosclerotic, anti-HIV, and cytotoxic activities [30]. Based on the results of the present study, hydroalcoholic extract of *A. tripedale* had a significant inhibitory effect against the growth of various strains of *Candida*. In sum, it seems that *A. tripedale* is a major source of anti-fungal compounds, which can be applied for the treatment of infectious diseases.

Conclusion

This is the first report on the GC-MS analysis of *A. tripedale*. It can be concluded that *A. tripedale* contains various important bioactive compounds. Therefore, it is recommended as a plant of phytochemical and pharmaceutical importance. Further studies are required to isolate the active ingredients of the extract and elucidate its mechanism of action in various diseases.

Acknowledgments

This study was labeled as a Student Research Project (No. 94S85) and was financially supported by the Student Research Committee of Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

Author's contribution

A. S and M. S designed and managed the study and contributed to data analysis and interpretation. A. S wrote the main manuscript. M. S. revised the first draft of the manuscript. M. M and A. KZ set up the test and managed the research. All the authors reviewed the manuscript.

Conflicts of interest

None declared.

Financial disclosure

There was no financial interest related to the materials of the manuscript.

References

- Ramesh P, Okigbo RN. Effects of plants and medicinal plant combinations as anti-infectives. *Afr J Pharm Pharmacol*. 2008; 2(7):130-5.
- Arif T, Bhosale JD, Kumar N, Mandal TK, Bendre RS, Lavekar GS, et al. Natural products-antifungal agents derived from plants. *J Asian Nat Prod Res*. 2009; 11(7):621-38.
- Hashemi SJ, Asgarpanah J, Alaei Z, Sadeghian S, Hasani H, Azimi A. In vitro antifungal activity of four medicinal plants used in Iranian Traditional Medicine. *Res J Pharm*. 2014; 1(1):39-43.
- Rios JL, Recio MC. Medicinal plants and antimicrobial activity. *J Ethnopharmacol*. 2005; 100(1-2):80-4.
- Kazemi S, Asgary S, Moshtaghian J, Rafieian M, Adelnia A, Shamsi F. Liver-protective effects of hydroalcoholic extract of *allium hirtifolium* boiss. In rats with alloxan-induced diabetes mellitus. *ARYA Atheroscler*. 2010; 6(1):11-5.
- Mnayer D, Fabiano-Tixier AS, Petitcolas E, Hamieh T, Nehme N, Ferrant C, et al. Chemical composition, antibacterial and antioxidant activities of six essential oils from the Alliaceae family. *Molecules*. 2014; 19(12):20034-53.
- Lazarević JS, Đorđević AS, Zlatković BK, Radulović NS, Palić RM. Chemical composition and antioxidant and antimicrobial activities of essential oil of *Allium sphaerocephalon* L. subsp. *sphaerocephalon* (Liliaceae) inflorescences. *J Sci Food Agric*. 2011; 91(2):322-9.
- Pfaller MA, Diekema DJ. Epidemiology of invasive Candidiasis: a persistent public health problem. *Clin Microbiol Rev*. 2007; 20(1):133-63.
- Rehman A, Rehman A, Ahmad I. Antibacterial, antifungal, and insecticidal potentials of *Oxalis corniculata* and its isolated compounds. *Int J Anal Chem*. 2015; 2015:842468.
- Corsello S, Spinillo A, Osnengo G, Penna C, Guaschino S, Beltrame A, et al. An epidemiological survey of vulvovaginal candidiasis in Italy. *Eur J Obstet Gynaecol Reprod Biol*. 2003; 110(1):66-72.
- Richter SS, Galask RP, Messer SA, Hollis RJ, Diekema DJ, Pfaller MA. Antifungal susceptibilities of *Candida* species causing vulvovaginitis and epidemiology of recurrent cases. *J Clin Microbiol*. 2005; 43(5):2155-62.
- Nasrollahi Z, Yadegari MH, Roudbar Mohammadi S, Roudbary M, Hosseini Poor M, Nikoomanesh F, et al. Fluconazole resistance *Candida albicans* in females with recurrent Vaginitis and Pirl1 overexpression. *Jundishapur J Microbiol*. 2015; 8(9):e21468.
- Dabur R, Singh H, Chhillar AK, Ali M, Sharma GL. Antifungal potential of Indian medicinal plants. *Fitoterapia*. 2004; 75(3-4):389-91.
- Das K, Tiwari RK, Shrivastava DK. Techniques for evaluation of medicinal plant products as antimicrobial agents: current methods and future trends. *J Med Plants Res*. 2010; 4(2):104-11.
- Forbes BA, Sahm DF, Weissfeld AS. Diagnostic microbiology. *Bailey Scott Diagn Microbiol*. 2002; 11(1):11-4.
- Balouiri M, Sadiki M, Ibsouda SK. Methods for in vitro evaluating antimicrobial activity: a review. *J Pharm Biomed Anal*. 2016; 6(2):71-9.
- Kim HJ, Suh HJ, Lee CH, Kim JH, Kang SC, Park S, et al. Antifungal activity of glyceollins isolated from soybean elicited with *Aspergillus sojae*. *J Agric Food Chem*. 2010; 58(17):9483-7.
- Sojakova M, Liptajova D, Borovsky M, Subik J. Fluconazole and itraconazole susceptibility of vaginal yeast isolates from Slovakia. *Mycopathologia*. 2004; 157(2):163-9.
- Kirkpatrick WR, Turner TM, Fothergill AW, McCarthy DI, Redding SW, Rinaldi MG, et al. Fluconazole disk diffusion susceptibility testing of *Candida* species. *J Clin*

- Microbiol. 1998; 36(11):3429-32 .
20. Brown GD, Denning DW, Gow NA, Levitz SM, Netea MG, White TC. Hidden killers: human fungal infections. *Sci Transl Med*. 2012; 4(165):165rv13.
 21. Amjad L, Rezvani Z, Madani M. The effect of methanolic extract of *anthesis gayana* on *Candida* Spp. *Intl J Agric Crop Sci*. 2013; 5(10):1140.
 22. Abad MJ, Ansuategui M, Bermejo P. Active antifungal substances from natural sources. *Arkivoc*. 2007; 7(11):116-45.
 23. Shams-Ghahfarokhi M, Shokoohamiri MR, Amirrajab N, Moghadasi B, Ghajari A, Zeini F, et al. In vitro antifungal activities of *Allium cepa*, *Allium sativum* and ketoconazole against some pathogenic yeasts and dermatophytes. *Fitoterapia*. 2006; 77(4):321-3.
 24. Amin M, Kapadnis BP. Heat stable antimicrobial activity of *Allium ascalonicum* against bacteria and fungi. *Indian J Exp Biol*. 2005; 43(8):751-4 .
 25. Dandekar R, Fegade B, Bhaskar VH. GC-MS analysis of phytoconstituents in alcohol extract of *Epiphyllum oxypetalum* leaves. *J Pharm Phytochem*. 2015; 4(1):149-54.
 26. Theng KB, Korpenwar AN. Phytochemical analysis of ethanol extract of *ampelocissus latifolia* (Roxb.) planch tuberous root using UV-vis, ftir and GC-MS. *Int J Pharm Sci Res*. 2015; 6(9):3936.
 27. Anjusha S, Gangaprasad A, Radhamany PM. GC-MS analysis of leaf extract of *gynochthodes umbellata* (L.) razafim. and b. bremer (Rubiaceae). *Int J Pharm Sci Res*. 2015; 6(11):4826.
 28. Zelenitsky SA, Zhanel GG. Phenazopyridine in urinary tract infections. *Ann Pharmacother*. 1996; 30(7-8):866-8.
 29. Choojun S, Yoonprayong P. Improvement of propionic acid production for antifungal activity from whey by calcium alginate immobilization of *propionibacterium acidipropionici* TISTR 442. *J Agric Sci Tech*. 2012; 2(7A):863.
 30. Chandrasekar T, Rao MR, Kumar RV, Prabhu K, Nandha Kumar S, Divya D. GC-MS analysis, antimicrobial, antioxidant activity of an Ayurvedic medicine, Nimbapatradi Choomam. *J Chem Pharm Res*. 2015; 7(8):124-36.

Archive of SID