



# Radiographic and Surgical Outcomes After Stand-Alone Lateral Lumbar Interbody Fusion

Amrit S. Khalsa,<sup>1</sup> Gregory M. Mundis,<sup>2</sup> Justin B. Ledesma,<sup>1</sup> Pooria Hosseini,<sup>1</sup> James D. Bruffey,<sup>2</sup> Stacie Nguyen,<sup>1</sup> Behrooz A. Akbarnia,<sup>1</sup> and Robert K. Eastlack<sup>2,\*</sup>

<sup>1</sup>San Diego Spine Foundation, San Diego, California, United States

<sup>2</sup>Division of Orthopaedic Surgery, Scripps Clinic, La Jolla, California, United States

\*Corresponding author: Division of Orthopaedic Surgery, Scripps Clinic 10666 North Torrey Pines Road, MS116 La Jolla, 92037 California, USA. Tel: +1-858-554-7988, Email: eastlack.robert@scrippshealth.org

Received 2018 June 27; Accepted 2018 July 15.

## Abstract

**Objectives:** Lateral lumbar interbody fusion (LLIF) is increasingly being utilized in isolation to achieve a large surface-area interbody fusion with an indirect decompression for spinal stenosis. This retrospective chart review was done to determine the viability of performing stand-alone (SA) LLIF.

**Methods:** Forty-nine patients at least 18 years of age with minimum one-year follow-up at a single institution underwent SA-LLIF using minimally invasive surgery (MIS) approach without further posterior surgery between 2011 and 2015. One to five-level fusions were included. Retrospective review of surgical outcomes and radiographic parameters were examined preoperatively, acutely post-operatively and at 1 year postoperatively.

**Results:** Forty-nine patients (102 spinal segments) underwent SA-LLIF. Fusion levels ranged from one to five with a mean of  $2.1 \pm 2.1$ . Mean blood loss was  $68 \pm 63.2$ cc and mean surgical time was  $143.4 \pm 66.5$  minutes. Fifty-seven percent had undergone prior spine surgery unrelated to their index procedure. Complication rate was 38.9% and reoperation rate was 20.4%. No difference in complication rates was noted between constructs with three or more levels fused versus less than three levels fused. At one-year, significant improvement was noted with pelvic tilt, pelvic incidence, and lumbar lordosis.

**Conclusions:** SA-LLIF is an optional MIS treatment of stable degenerative disc disease and spinal stenosis, with good one-year correction and maintenance of radiographic parameters. With complication rate of 38.9% and reoperation rate of 20.4%, true benefit of forgoing posterior supplemental fixation may be questioned.

**Keywords:** Lateral Lumbar Interbody Fusion, Minimally Invasive Surgery, Tand-Alone Interbody Fusion, Degenerative Disc Disease, Spinal Stenosis

## 1. Background

Degenerative disc disease (DDD) is almost universal in the aging population (1). Long-term sequela of this condition can lead to loss of disc height, neuroforaminal and central canal stenosis, and eventual endplate sclerosis as advanced arthritic disease begins (2, 3). Treatment for DDD is often a spectrum ranging from conservative management to decompression and fusion.

Posterior decompression procedures can put spinal levels, already biomechanically abnormal due to underlying disease, at further risk of instability (4, 5). Spondylolisthesis as well as sagittal and coronal plane deformities can develop, subsequently worsening stenosis and neurologic compromise.

Lateral lumbar interbody fusion (LLIF) through a minimally-invasive surgery (MIS) approach is increasingly being utilized in isolation to achieve a large surface-area interbody fusion with an indirect decompression for spinal stenosis (6-10). This technique forgoes posterior fixation with a pedicle screw-rod construct in the appropriately selected patient with a stable spinal segment. The obvious benefits include decreased blood loss, shorter operative time, and absence of posterior soft-tissue disruption (11-13). Scant data exists regarding surgical and radiographic outcomes following this treatment algorithm that intuitively may have the benefits of decreasing adjacent-segment forces and the potential disadvantage of a less-stable fusion construct.

## 2. Methods

To identify the viability of stand-alone (SA)-LLIF, a retrospective case series review of patients undergoing the procedure for DDD, spinal stenosis, and low-grade spondylolisthesis between T12 and L5 was conducted at a single institution after approval from an institutional review board.

Forty-nine patients at least 18 years of age underwent SA-LLIF via a standard MIS retroperitoneal, trans-psoas approach as previously described in the literature. Interbody cages varied from 8 to 12 mm in height, 45 to 60 mm in width, 18 to 26 mm in depth, and 8 to 10 degrees of lordosis. No further posterior surgery was performed. Index procedures occurred between 2011 and 2015 with a minimum one-year follow-up. One to five-level SA-LLIF were included. A retrospective review of surgical outcomes and radiographic parameters were employed preoperatively, acutely postoperatively and at one year postoperatively.

Clinical data was assessed through chart review of physical examination and clinical history in both the inpatient and outpatient settings. Two independent surgeons that were not involved with direct care of the patients analyzed digital x-ray images. The following radiographic parameters were assessed utilizing Surgimap Spine software (Nemaris Inc, New York, NY, USA): pelvic tilt (PT), sacral slope (SS), pelvic incidence (PI), lumbar lordosis (LL), L4 to S1 LL (PI-LL), motion segment angle (MSA), and intradiscal angle (IDA). X-rays were reviewed preoperatively, postoperatively, and at one year. Computed tomography (CT) and x-rays images were reviewed to assess fusion.

### 2.1. Statistical Analysis

All statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS Version 24, Chicago, Ill, USA). Pre- to postoperative comparisons were assessed by non-parametric Wilcoxon signed-rank test. Between group comparisons were assessed by non-parametric Mann-Whitney U test. Categorical variables were compared using chi-squared test. Statistical significance was set at alpha level of 0.05.

## 3. Results

Forty-nine patients (102 spinal segments) underwent SA-LLIF. The mean age, fusion levels, estimated blood loss and surgical time are provided in [Table 1](#).

Prior spine surgery unrelated to their index procedure had been undergone by 57.1%. Overall complication rate was 38.9% ([Table 2](#)).

**Table 1.** Patient Demographics

| Descriptor (N = 49)            | Mean  | Range        | Standard Deviation |
|--------------------------------|-------|--------------|--------------------|
| Age, y                         | 63.2  | (32, 84)     | 9.8                |
| BMI                            | 27.4  | (18.4, 46.3) | 5.3                |
| Number of LLIFs                | 2.1   | (1, 5)       | 1.3                |
| Total estimated blood loss, ml | 68    | (0, 300)     | 63.2               |
| Total surgical time, minutes   | 143.4 | (72, 328)    | 66.5               |

Abbreviations: BMI, body mass index; LLIF, lateral lumbar interbody fusion.

**Table 2.** Complications Experienced

| Complications                   | No. (%)   |
|---------------------------------|-----------|
| <b>Overall complication</b>     | 19 (38.9) |
| <b>Reoperation</b>              | 10 (20.4) |
| Major                           | 2 (4.1)   |
| Minor                           | 17 (34.7) |
| <b>Surgical</b>                 | 18 (36.7) |
| Infection                       | 0 (0.0)   |
| Implant                         | 2 (4.1)   |
| Radiographic                    | 10 (20.4) |
| Neurologic                      | 8 (16.3)  |
| <b>Cardiopulmonary</b>          | 0 (0.0%)  |
| Vascular                        | 0 (0.0)   |
| <b>Gastrointestinal</b>         | 0 (0.0)   |
| <b>Renal</b>                    | 0 (0.0)   |
| <b>Operative</b>                | 1 (2.0)   |
| <b>Wound problems</b>           | 0 (0.0)   |
| <b>Adjacent segment disease</b> | 5 (10.2)  |
| <b>Pseudoarthrosis</b>          | 3 (6.1)   |

Eight cases were adjacent segment disease (ASD), seven being symptomatic, five undergoing reoperation at >1 year, 2 at < 1 year. Four cases were pseudoarthrosis, one undergoing reoperation at >1 year, two at < 1 year, and one non-operatively managed. One case was an endplate fracture at 6 weeks postoperative requiring the addition of posterior fixation. Others included three cases of ongoing radiculopathy (2 L5, 1 S1) and two cases of residual motor deficit (one foot dorsiflexion and one hip abductor) at 1 year. Overall reoperation rate was 20.4%. No difference in complication rates were noted between constructs with three or more levels fused versus less than three levels fused or constructs with prior fusions ([Table 3](#)).

At 1-year, significant improvement was seen with PT, PI,

**Table 3.** Complication Rates Were Noted Between Constructs With and Without Prior Fusions

| Descriptor          | No. Prior Lumbar Fusion (N = 36), No. (%) | Prior Lumbar Fusion (N = 13), No. (%) | P     |
|---------------------|---|---------------------------------------|-------|
| <b>Complication</b> | 12 (33.3)                                 | 7 (53.8)                              | 0.193 |
| <b>Reoperation</b>  | 6 (16.7)                                  | 4 (30.8)                              | 0.280 |
| <b>Major</b>        | 1 (2.8)                                   | 1 (7.7)                               | 0.443 |
| <b>Minor</b>        | 11 (30.6)                                 | 6 (46.2)                              | 0.311 |
| <b>Surgical</b>     | 11 (30.6)                                 | 7 (53.8)                              | 0.135 |
| <b>Implant</b>      | 1 (2.8)                                   | 1 (7.7)                               | 0.443 |
| <b>Radiographic</b> | 7 (19.4)                                  | 3 (23.1)                              | 0.781 |
| <b>Neurologic</b>   | 5 (13.9)                                  | 3 (23.1)                              | 0.442 |
| <b>Operative</b>    | 1 (2.8)                                   | 0 (0.0)                               | 0.544 |

and LL. PI-LL were both significantly improved acutely postoperative and at 1 year (Table 4). At both acutely postoperative and 1-year follow-up, mean IDA and MSA were significantly improved when compared with preoperative values (Table 5).

Sub-analysis showed no difference in postoperative radiographic parameters for patients that underwent reoperation (Table 6).

Further sub-analysis was performed to evaluate the effect of deformity, defined as preoperative sagittal PI-LL > 10 and PT > 20, on SA-LLIF. There were 22 (44.9%) deformity patients (DEF) and 27 (55.1%) non-deformity patients (NDEF) (Table 7).

Mean age was 65.2 and 61.6 in DEF and NDEF, respectively. No statistical differences between groups was seen in demographics. An average of 1.9 levels were fused in NDEF vs. 2.5 (P = 0.135) in DEF. Complication (DEF 40.7% and NDEF 36.4%, P = 0.754) and reoperation rates (22.7% DEF and 22.1% NDEF, P = 0.966) were similar between groups. Reoperation at < 1 year for DEF included one pseudoarthrosis and for NDEF one endplate fracture, one symptomatic ASD, and two pseudoarthrosis (PSA). At > 1-year reoperations included for DEF one PSA and three ASD; for NDEF two ASD. Other complications included for DEF was one residual hip abductor weakness, one persistent S1 radiculopathy, one asymptomatic ASD, and one symptomatic PSA treated non-operatively. NDEF had one residual foot dorsiflexion weakness and two persistent L5 radiculopathies. Overall neurologic risk was 13.6% for DEF and 18.5% for NDEF, which did not reach significance (p = 0.646).

#### 4. Discussion

Much discussion exists about how to manage the degenerative milieu of DDD, spinal stenosis, low-grade spondylolisthesis and mild degenerative deformities. When the decision to proceed with fusion is made, debate again exists about the method to use. MIS techniques attempt to provide a stable fusion construct while minimizing collateral damage that may further progress the underlying disease process (14, 15). SA-LLIF is a relatively new technique that attempts to minimize the deleterious effects of posterior instrumentation while still providing a reliable segmental fusion.

Several biomechanical studies have shown SA-interbody fusion provide sufficient stability while reducing stress at adjacent levels (16, 17). By restoring tensile strain on intact ligamentous structures, insertion of an interbody device can provide adequate stability while restoring disc height and correcting anterior and middle column alignment (18).

Marchi et al. (12), reported a revision rate of 13.5% after SA-LLIF with the majority related to implant subsidence. Previous studies show that subsidence plays no role in reported outcome measures but is a source of at least one revision in this case series. The majority of revisions reported were due to pseudoarthrosis and adjacent segment failure in this case series. Adjacent segment failure remains controversial as possibly the natural course of the degenerative process. In the case series by Ahmadian et al. (14), only two patients out of 59 underwent reoperation for continued symptomatic stenosis and cage migration. Watkins et al. (19) found a 19% pseudoarthrosis rate in a case series of 23 consecutive patients and 37 levels treated. Another study identified a surgical revision rate of 10.3% in 117 patients undergoing SA-LLIF (20). Tempel et al. (21), had two vertebral body fractures in a series of 335 patients after SA-LLIF.

Several previously published studies indicate that SA-LLIF consistently adds approximately 2° - 3° of segmental lordosis per level. This is consistent with the current data. Malham et al. (22), reported a significant overall lumbar lordosis improvement from 48.8° to 55.2° at 1 year. Several other studies have shown mixed correction of global lordosis similar to the current study (23, 24).

No previous studies have examined the use of SA-LLIF in adult deformity surgery despite the increasing trend for MIS lateral spine surgery in this population. Though SA-LLIF remains controversial, advocates can see based on

**Table 4.** Results Preoperative, Acutely Postoperative and One-Year Postoperative

| Descriptor                | Preoperative | Postoperative           | 1-Year Postoperative     |
|---------------------------|--------------|-------------------------|--------------------------|
| Pelvic tilt, degrees      | 20.7 ± 9.4   | 20.8 ± 9.0              | 18.4 ± 7.8 <sup>a</sup>  |
| Sacral slope, degrees     | 34.6 ± 8.4   | 33.4 ± 8.5              | 35.4 ± 8.5               |
| Pelvic incidence, degrees | 55.3 ± 11.4  | 54.1 ± 11.1             | 53.7 ± 11.6 <sup>a</sup> |
| PI-LL, degrees            | 13.3 ± 21.4  | 8.0 ± 12.7 <sup>a</sup> | 6.9 ± 11.8 <sup>a</sup>  |
| Lumbar lordosis, degrees  | 42.0 ± 20.3  | 46.1 ± 11.4             | 46.8 ± 12.5 <sup>a</sup> |
| L4-S1 LL, degrees         | 33.1 ± 14.3  | 33.3 ± 8.9              | 35.0 ± 9.8               |

Abbreviations: PI-LL, pelvic incidence-lumbar lordosis; LL, lumbar lordosis.

<sup>a</sup> Significantly different from preoperative.**Table 5.** Comparison of Preoperative and Postoperative Angles

| Descriptor                     | N   | Preoperative               | Acute Postoperative | P       |
|--------------------------------|-----|----------------------------|---------------------|---------|
| Intradiscal angles, Degrees    | 102 | 3.0 ± 3.4                  | 7.2 ± 3.9           | < 0.001 |
| Motion-segment angles, Degrees | 104 | 9.7 ± 9.6                  | 12.6 ± 9.1          | < 0.001 |
|                                |     | <b>Preoperative</b>        | <b>Latest</b>       |         |
| Intradiscal angles, Degrees    | 104 | 3.0 ± 3.4                  | 6.6 ± 3.5           | < 0.001 |
| Motion-segment angles, Degrees | 101 | 9.9 ± 9.5                  | 12.6 ± 9.4          | < 0.001 |
|                                |     | <b>Acute Postoperative</b> | <b>Latest</b>       |         |
| Intradiscal angles, Degrees    | 102 | 7.2 ± 3.9                  | 6.6 ± 3.6           | 0.117   |
| Motion-segment angles, Degrees | 101 | 12.7 ± 9.1                 | 12.6 ± 9.4          | 0.789   |

**Table 6.** Sub-Analysis Showed No Difference in Postoperative Radiographic Parameters for Patients Who Underwent Reoperation

| Descriptor  | No. Reoperation (N = 39) | Reoperation (N = 10) | P     |
|---|--------------------------|----------------------|-------|
| Postoperative pelvic tilt, Degrees                      | 21.5                     | 19.3                 | 0.705 |
| Postoperative sacral slope, Degrees                     | 33.9                     | 31.4                 | 0.549 |
| Postoperative pelvic incidence, Degrees                 | 55                       | 50.7                 | 0.285 |
| Postoperative pelvic incidence-lumbar lordosis, Degrees | 8.5                      | 6.1                  | 0.517 |
| Postoperative lumbar lordosis, Degrees                  | 46.5                     | 44.6                 | 0.817 |
| Postoperative L4-S1 lumbar lordosis, Degrees            | 33.5                     | 32.6                 | 0.913 |

the current study that SA-LLIF is a reasonable alternative to more traditional reconstruction approaches used in an appropriately selected patient population.

Limitations of this study include the lack of patient reported outcome measures. This study focused on surgical complications and radiographic outcomes. The retrospective case series cohort is a limitation. More long-term follow-up beyond 1-year radiographic data would be helpful in further delineating the nature of adjacent segment disease and complication rates. We further acknowledge that full-length scoliosis radiographs would have been ideal for measuring global sagittal parameters in the deformity population.

SA-LLIF is a viable option for minimally invasive treat-

ment of stable DDD and spinal stenosis, with good one-year correction and maintenance of radiographic sagittal parameters. With an overall complication rate approaching 40%, the true benefit of forgoing posterior supplemental fixation may be questioned and the appropriate candidate may need to be defined for this less invasive procedure to minimize complications.

### Acknowledgments

We thank the scripps clinic medical group for the internal grant to support this research and the Shiley center for orthopaedic research and education at scripps clinic for their expertise and support.

**Table 7.** Sub-Analysis Performed to Evaluate the Effect of Deformity, Defined as Preoperative Sagittal PI-LL > 10 and Pelvic Tilt > 20, on Stand-Alone Lateral Interbody Fusion

| Descriptor                         | Non-Deformity Patients | Deformity Patients | P                  |
|------------------------------------|------------------------|--------------------|--------------------|
| N                                  | 27                     | 22                 | 0.329              |
| Age, y                             | 61.6                   | 65.2               | 0.944              |
| Female                             | 12 (44.4%)             | 10 (45.5%)         | 0.406              |
| Body mass index                    | 27.9                   | 26.8               | 0.136              |
| Prior spine surgery, N             | 18 (66.7%)             | 10 (45.5%)         | 0.135              |
| Total LLIFs, N                     | 50                     | 54                 | 0.967              |
| Preoperative hemoglobin            | 14.1                   | 14.2               | 0.7                |
| Total estimated blood loss, mL     | 68                     | 68.1               | 0.132              |
| Total operating room time, minutes | 132.4                  | 156.9              | 0.004 <sup>a</sup> |
| Total length of surgery, hours     | 1.9                    | 2.9                | 0.329              |
| Complication, N (%)                | 11 (40.7%)             | 8 (36.4%)          | 0.754              |
| Reoperation, N (%)                 | 5 (18.5%)              | 5 (22.7%)          | 0.716              |
| Major complications, N (%)         | 2 (7.4%)               | 0 (0.0%)           | 0.192              |
| Minor complications, N (%)         | 9 (33.3%)              | 8 (36.4%)          | 0.825              |
| Surgical complications, N (%)      | 11 (40.7%)             | 7 (31.8%)          | 0.519              |
| Implant complications, N (%)       | 2 (7.4%)               | 0 (0.0%)           | 0.192              |
| Radiographic, N (%)                | 4 (14.8%)              | 6 (27.3%)          | 0.282              |
| Neurologic, N (%)                  | 5 (18.5%)              | 3 (13.6%)          | 0.646              |
| Operative, N (%)                   | 1 (3.7%)               | 0 (0.0%)           | 0.362              |
| Pre to 1-y PI, degrees             | 0.6 <sup>a</sup>       | -6 <sup>a</sup>    | 0.003 <sup>a</sup> |
| Pre to 1-y SS, degrees             | -1.6                   | 3.7 <sup>a</sup>   | 0.03 <sup>a</sup>  |
| Pre to 1-y PI, degrees             | -1                     | -2.3               | 0.351              |
| Pre to 1-y PI-LL, degrees          | -6 <sup>a</sup>        | -6.9 <sup>a</sup>  | 0.01 <sup>a</sup>  |
| Pre to 1-y LL, degrees             | 4.9                    | 4.6                | 0.116              |
| Pre to 1-y L4-S1 LL, degrees       | 2                      | 1.9                | 0.475              |
| Pre to 1-y IDA, degrees            | 4.1                    | 4                  | 0.608              |
| Pre to 1-y MSA, degrees            | 4.1                    | 1.8                | 0.183              |

Abbreviations: IDA = intradiscal angles; LL = lumbar lordosis; MSA = motion-segment angles; PI = pelvic incidence; PI-LL = pelvic incidence-lumbar lordosis; PT = pelvic tilt; SS = sacral slope.

<sup>a</sup> Indicates statistical significance.

## Footnote

**Funding/Support:** An internal grant from scripps clinic medical group supported this research.

## References

1. Teraguchi M, Yoshimura N, Hashizume H, Yamada H, Oka H, Minamide A, et al. Progression, incidence, and risk factors for intervertebral disc degeneration in a longitudinal population-based cohort: the Wakayama Spine Study. *Osteoarthritis Cartilage*. 2017;**25**(7):1122–31. doi: [10.1016/j.joca.2017.01.001](https://doi.org/10.1016/j.joca.2017.01.001). [PubMed: [28089899](https://pubmed.ncbi.nlm.nih.gov/28089899/)].
2. Farshad-Amacker NA, Farshad M, Winklehner A, Andreisek G. MR imaging of degenerative disc disease. *Eur J Radiol*. 2015;**84**(9):1768–76. doi: [10.1016/j.ejrad.2015.04.002](https://doi.org/10.1016/j.ejrad.2015.04.002). [PubMed: [26094867](https://pubmed.ncbi.nlm.nih.gov/26094867/)].
3. Kanna RM, Shetty AP, Rajasekaran S. Patterns of lumbar disc degeneration are different in degenerative disc disease and disc prolapse magnetic resonance imaging analysis of 224 patients. *Spine J*. 2014;**14**(2):300–7. doi: [10.1016/j.spinee.2013.10.042](https://doi.org/10.1016/j.spinee.2013.10.042). [PubMed: [24231779](https://pubmed.ncbi.nlm.nih.gov/24231779/)].
4. Bisschop A, Kingma I, Bleys RL, van der Veen AJ, Paul CP, van Dieen JH, et al. Which factors prognosticate rotational instability following lumbar laminectomy? *Eur Spine J*. 2013;**22**(12):2897–903. doi: [10.1007/s00586-013-3002-3](https://doi.org/10.1007/s00586-013-3002-3). [PubMed: [24043337](https://pubmed.ncbi.nlm.nih.gov/24043337/)]. [PubMed Central: [PMC3843774](https://pubmed.ncbi.nlm.nih.gov/PMC3843774/)].
5. Guha D, Heary RF, Shamji MF. Iatrogenic spondylolisthesis following laminectomy for degenerative lumbar stenosis: systematic review and current concepts. *Neurosurg Focus*. 2015;**39**(4). E9. doi: [10.3171/2015.7.FOCUSI5259](https://doi.org/10.3171/2015.7.FOCUSI5259). [PubMed: [26424349](https://pubmed.ncbi.nlm.nih.gov/26424349/)].
6. Ozgur BM, Hughes SA, Baird LC, Taylor WR. Minimally disruptive decompression and transforaminal lumbar interbody fusion. *Spine J*. 2006;**6**(1):27–33. doi: [10.1016/j.spinee.2005.08.019](https://doi.org/10.1016/j.spinee.2005.08.019). [PubMed: [16413444](https://pubmed.ncbi.nlm.nih.gov/16413444/)].
7. Johnson RD, Valore A, Villaminar A, Comisso M, Balsano M. Pelvic parameters of sagittal balance in extreme lateral interbody fusion for degenerative lumbar disc disease. *J Clin Neurosci*. 2013;**20**(4):576–81. doi: [10.1016/j.jocn.2012.05.032](https://doi.org/10.1016/j.jocn.2012.05.032). [PubMed: [23375396](https://pubmed.ncbi.nlm.nih.gov/23375396/)].
8. Palejwala SK, Sheen WA, Walter CM, Dunn JH, Baaj AA. Minimally invasive lateral transpoas interbody fusion using a stand-alone construct for the treatment of adjacent segment disease of the lumbar spine: review of the literature and report of three cases. *Clin*

- Neurol Neurosurg.* 2014;**124**:90–6. doi: [10.1016/j.clineuro.2014.06.031](https://doi.org/10.1016/j.clineuro.2014.06.031). [PubMed: [25019458](https://pubmed.ncbi.nlm.nih.gov/25019458/)].
9. Tohmeh AG, Khorsand D, Watson B, Zielinski X. Radiographical and clinical evaluation of extreme lateral interbody fusion: effects of cage size and instrumentation type with a minimum of 1-year follow-up. *Spine (Phila Pa 1976)*. 2014;**39**(26):E1582–91. doi: [10.1097/BRS.0000000000000645](https://doi.org/10.1097/BRS.0000000000000645). [PubMed: [25341985](https://pubmed.ncbi.nlm.nih.gov/25341985/)].
  10. Karikari IO, Nimjee SM, Hardin CA, Hughes BD, Hodges TR, Mehta AI, et al. Extreme lateral interbody fusion approach for isolated thoracic and thoracolumbar spine diseases: initial clinical experience and early outcomes. *J Spinal Disord Tech.* 2011;**24**(6):368–75. doi: [10.1097/BSD.0b013e3181ffefd2](https://doi.org/10.1097/BSD.0b013e3181ffefd2). [PubMed: [21150667](https://pubmed.ncbi.nlm.nih.gov/21150667/)].
  11. von Keudell A, Alimi M, Gebhard H, Hartl R. Adult Degenerative Scoliosis with Spinal Stenosis Treated with Stand-Alone Cage via an Extreme Lateral Transposoas Approach; a Case Report and Literature Review. *Arch Bone Jt Surg.* 2015;**3**(2):124–9. [PubMed: [26110180](https://pubmed.ncbi.nlm.nih.gov/26110180/)]. [PubMed Central: [PMC4468624](https://pubmed.ncbi.nlm.nih.gov/PMC4468624/)].
  12. Marchi L, Abdala N, Oliveira L, Amaral R, Coutinho E, Pimenta L. Stand-alone lateral interbody fusion for the treatment of low-grade degenerative spondylolisthesis. *ScientificWorldJournal.* 2012;**2012**:456346. doi: [10.1100/2012/456346](https://doi.org/10.1100/2012/456346). [PubMed: [22545019](https://pubmed.ncbi.nlm.nih.gov/22545019/)]. [PubMed Central: [PMC3324177](https://pubmed.ncbi.nlm.nih.gov/PMC3324177/)].
  13. Udby PM, Bech-Azeddine R. Clinical outcome of stand-alone ALIF compared to posterior instrumentation for degenerative disc disease: A pilot study and a literature review. *Clin Neurol Neurosurg.* 2015;**133**:64–9. doi: [10.1016/j.clineuro.2015.03.008](https://doi.org/10.1016/j.clineuro.2015.03.008). [PubMed: [25855474](https://pubmed.ncbi.nlm.nih.gov/25855474/)].
  14. Ahmadian A, Bach K, Bolinger B, Malham GM, Okonkwo DO, Kanter AS, et al. Stand-alone minimally invasive lateral lumbar interbody fusion: multicenter clinical outcomes. *J Clin Neurosci.* 2015;**22**(4):740–6. doi: [10.1016/j.jocn.2014.08.036](https://doi.org/10.1016/j.jocn.2014.08.036). [PubMed: [25684343](https://pubmed.ncbi.nlm.nih.gov/25684343/)].
  15. Berjano P, Langella F, Damilano M, Pejrona M, Buric J, Ismael M, et al. Fusion rate following extreme lateral lumbar interbody fusion. *Eur Spine J.* 2015;**24 Suppl 3**:369–71. doi: [10.1007/s00586-015-3929-7](https://doi.org/10.1007/s00586-015-3929-7). [PubMed: [25893332](https://pubmed.ncbi.nlm.nih.gov/25893332/)].
  16. Kretzer RM, Molina C, Hu N, Umekoji H, Baaj AA, Serhan H, et al. A Comparative Biomechanical Analysis of Stand Alone Versus Facet Screw and Pedicle Screw Augmented Lateral Interbody Arthrodesis: An In Vitro Human Cadaveric Model. *Clin Spine Surg.* 2016;**29**(7):E336–43. doi: [10.1097/BSD.0b013e3182868ef9](https://doi.org/10.1097/BSD.0b013e3182868ef9). [PubMed: [27137151](https://pubmed.ncbi.nlm.nih.gov/27137151/)].
  17. Choi KC, Ryu KS, Lee SH, Kim YH, Lee SJ, Park CK. Biomechanical comparison of anterior lumbar interbody fusion: stand-alone interbody cage versus interbody cage with pedicle screw fixation – a finite element analysis. *BMC Musculoskelet Disord.* 2013;**14**:220. doi: [10.1186/1471-2474-14-220](https://doi.org/10.1186/1471-2474-14-220). [PubMed: [23890389](https://pubmed.ncbi.nlm.nih.gov/23890389/)]. [PubMed Central: [PMC3726285](https://pubmed.ncbi.nlm.nih.gov/PMC3726285/)].
  18. Wang SH, Celic I, Choi SY, Riccomagno M, Wang Q, Sun LO, et al. Dlg5 regulates dendritic spine formation and synaptogenesis by controlling subcellular N-cadherin localization. *J Neurosci.* 2014;**34**(38):12745–61. doi: [10.1523/JNEUROSCI.1280-14.2014](https://doi.org/10.1523/JNEUROSCI.1280-14.2014). [PubMed: [25232112](https://pubmed.ncbi.nlm.nih.gov/25232112/)]. [PubMed Central: [PMC4166160](https://pubmed.ncbi.nlm.nih.gov/PMC4166160/)].
  19. Watkins R IV, Watkins R III, Hanna R. Non-union rate with stand-alone lateral lumbar interbody fusion. *Medicine (Baltimore)*. 2014;**93**(29). e275. doi: [10.1097/MD.0000000000000275](https://doi.org/10.1097/MD.0000000000000275). [PubMed: [25546670](https://pubmed.ncbi.nlm.nih.gov/25546670/)]. [PubMed Central: [PMC4602616](https://pubmed.ncbi.nlm.nih.gov/PMC4602616/)].
  20. Nemani VM, Aichmair A, Taher F, Lebl DR, Hughes AP, Sama AA, et al. Rate of revision surgery after stand-alone lateral lumbar interbody fusion for lumbar spinal stenosis. *Spine (Phila Pa 1976)*. 2014;**39**(5):E326–31. doi: [10.1097/BRS.0000000000000141](https://doi.org/10.1097/BRS.0000000000000141). [PubMed: [24299718](https://pubmed.ncbi.nlm.nih.gov/24299718/)].
  21. Tempel ZJ, Gandhoke GS, Okonkwo DO, Kanter AS. Impaired bone mineral density as a predictor of graft subsidence following minimally invasive transposoas lateral lumbar interbody fusion. *Eur Spine J.* 2015;**24 Suppl 3**:414–9. doi: [10.1007/s00586-015-3844-y](https://doi.org/10.1007/s00586-015-3844-y). [PubMed: [25739988](https://pubmed.ncbi.nlm.nih.gov/25739988/)].
  22. Malham GM, Ellis NJ, Parker RM, Blecher CM, White R, Goss B, et al. Maintenance of Segmental Lordosis and Disk Height in Stand-alone and Instrumented Extreme Lateral Interbody Fusion (XLIF). *Clin Spine Surg.* 2017;**30**(2):E90–8. doi: [10.1097/BSD.0b013e3182aa4c94](https://doi.org/10.1097/BSD.0b013e3182aa4c94). [PubMed: [28207620](https://pubmed.ncbi.nlm.nih.gov/28207620/)].
  23. Lykissas MG, Jain VV, Nathan ST, Pawar V, Eismann EA, Sturm PF, et al. Mid- to long-term outcomes in adolescent idiopathic scoliosis after instrumented posterior spinal fusion: a meta-analysis. *Spine (Phila Pa 1976)*. 2013;**38**(2):E113–9. doi: [10.1097/BRS.0b013e31827ae3d0](https://doi.org/10.1097/BRS.0b013e31827ae3d0). [PubMed: [23124268](https://pubmed.ncbi.nlm.nih.gov/23124268/)].
  24. Marchi L, Abdala N, Oliveira L, Amaral R, Coutinho E, Pimenta L. Radiographic and clinical evaluation of cage subsidence after stand-alone lateral interbody fusion. *J Neurosurg Spine.* 2013;**19**(1):110–8. doi: [10.3171/2013.4.SPINE12319](https://doi.org/10.3171/2013.4.SPINE12319). [PubMed: [23662890](https://pubmed.ncbi.nlm.nih.gov/23662890/)].