

The Effect of Gluteus Medius Kinesio Taping on Pain in Athletes with Patellofemoral Pain Syndrome during Functional Tasks

Shahabeddin Bagheri^{1*}, Aynollah Naderi², Ali Taherinia³, Mohammad Hosain Rezvani⁴

¹ School of Humanities, University of Nahavand, Nahavand, Iran.

² School of Physical Education and Sport Sciences, Shahrood University of Technology, Shahrood, Iran.

³ School of Medicine, Alborz University of Medical Sciences, Karaj, Iran.

⁴ School of Sport Sciences, Shahrood University of Technology, Shahrood, Iran.

Received: 16 August 2019

Accepted: 17 November 2019

Abstract

Background: The weakness of hip muscles, abductors, and external rotators has been confirmed in individuals with patellofemoral pain syndrome in comparison with healthy people. This study investigates the effects of gluteus medius Kinesio taping in athletes with patellofemoral pain syndrome during functional tasks.

Methods: A total of 30 females with PFPS were randomly assigned into Kinesio taping group (n = 15) and the placebo group (n = 15). Pain intensity was measured using the numerical pain rating scale in 4 functional tasks; walking, squatting, ascending, and descending stairs, at baseline plus immediately as well as 24 following Kinesio taping. The data were then statistically analyzed.

Results: Mixed-model ANOVAs, with repeated measures on time, indicated statistically significant differences between groups in pain over time for walking (P = 0.000), squatting (P = 0.007), and descending stairs tasks (P = 0.000). However, there were no significant differences between groups in the ascending stair task (P = 0.651).

Conclusions: The results showed that compared to placebo in reducing pain in athletes with PFPS, gluteus Medius Kinesio taping was effective in functional tasks of walking, squatting, as well as moving up and down stairs.

Keywords: Gluteus medius, Kinesio taping, Patellofemoral pain syndrome, Functional task.

*Corresponding to: S Bagheri, Email: bagherishahab@yahoo.com

Please cite this paper as: Bagheri S, Naderi A, Taherinia A, Rezvani MH. The effect of gluteus Medius Kinesio taping on pain in athletes with patellofemoral pain syndrome during functional tasks. Int J Health Stud 2019;5(4):1-5

water resistant for 3-5 days.^{10,11} The use of KT affects the skin more than the muscles and articular structure. It is believed that to reduce pain and facilitate circulation, KT influences the interstitial tissue of the usual tension of the skin by providing appropriate local stimuli through the skin.⁹ It has been suggested that KT functions through giving persistent stimulation through the skin to enhance proprioception.¹¹ KT is also used to improve muscle function, increase lymphatic and vascular flow, reduce pain, and help improve joint misalignments.¹²

Various studies have been conducted on the efficacy of KT on injuries and musculoskeletal disorders. Reduction of neck pain and increase in cervical range of motion immediately and 24 hours after use in people with whiplash,¹³ mitigation of pain in shoulder movements in patients with shoulder pain,¹⁴ improvements in disability index and considerable decline of pain after one week KT use, have been reported in patients with chronic low back pain.¹⁵

Previously patellar taping has been used for pain reduction, increasing the strength of quadriceps, enhancing neuromuscular recruitment, and modifying the onset of VMO activation in the patients with PFPS.^{10,16,17} Also, taping creates a mechanical change in the patella, which reduces knee pain and results in the early treatment in people with PFPS. While PFPS can cause symptoms that range from uncomfortable to very painful, it seems that the condition can usually be treated by KT and exercise therapy.^{18,19,20}

In the literature, various studies have examined the effects of KT in patients PFPS.^{16,21-23} Since the imbalance in the EMG activity of the VMO and VL muscles leads to abnormal patellar tracking,²⁴ in these studies, mainly patellar taping has been considered. It is believed that increasing shear and compressive loads associated with abnormal patellar tracking may contribute to the appearance of patellofemoral pain.²⁵ The hypothesis that patella tracking is due to abnormal patellar kinematics on the constant femur is the basis for the design of therapeutic protocols in patients with PFPS. This is the result of the kinematic studies considered in non-weight-bearing conditions, regardless of the motion of the femur.²⁵ Powers et al. observed that the kinematics of the patellofemoral joint are different in bearing weight conditions.²⁶ According to this study, the main cause of the lateral tilt and its lateral displacement in patients with patella instability is the internal rotation of the femur.²⁶ For this reason, in recent studies, the proximal parts of the knee joint have been more widely

Introduction

Patellofemoral pain syndrome (PFPS) is one of the most common knee disorders in athletes. This complication is twice more likely to occur in females than in men.^{1,2} The most salient symptoms are pain when squatting, moving up and down stairs, running, jumping, and cycling.³

Given the disagreement about the etiology of PFPS, different treatments^{4,5} such as Quadriceps,⁴ and hip abductor and external rotator strengthening,⁶ hamstring and iliotibial band stretching, bracing,² patella taping⁷ have been suggested in people with PFPS. Taping is used both for the treatment and prevention of sports injuries. The major function of taping is to provide support while moving. Kinesio taping (KT) was introduced by Kenzo Kase in 1990 and became increasingly common among athletes.⁸ The techniques used by KT differ from traditional tape.⁹ KT is designed to imitate the elasticity of the skin, which provides appropriate local stimuli for articular structuring, rather than muscle strengthening.^{10,11} KT can be stretched from 30 to 140% of its original length and is

considered.²⁷⁻²⁹ This study is of great importance regarding the role and effect of gluteus medius (GMed) in pelvic and femoral biomechanics in PFPS. The reduction of muscle strength of the hip joint and the change in EMG activity of GMed in patients with PFPS have been reported in many studies.³⁰⁻³³ Delayed onset and shorter duration of GMed activity in moving up and down stairs in patients with PFPS have been confirmed in several studies.^{34,35} Delay in the GMed activation may reduce the stability of the hip joint in the frontal plane and increase the spin of the hip and knee joints in weight-bearing activities.³⁶ All these result in excessive femoral adduction and internal rotation during functional tasks and an increase in dynamic knee valgus plus reduction in the contact area of the patellofemoral joint, the potential cause of patellofemoral pain.^{37,38}

According to studies, the use of GMed Kinesio tape in the treatment of patients with PFPS is not well established. Accordingly, the present study investigates the effect of GMed Kinesio taping on pain in athletes with PFPS during functional tasks.

Materials and Methods

Thirty females with PFPS from sports clubs were included in the study. All participants were informed about the research, and each gave her written consent. They were randomly classified into two groups: placebo (15) and KT groups (15). The present study was a part of scientific research conducted in the winter of 2015 in Hamedan, Iran. The inclusion criteria were: being athlete, i.e. participating in athletic activities at least 3 days a week for 30 minutes per day for six months; pain in at least 2 cases; moving up and down stairs, squatting, long-term seating with bending knee, running, jumping, pain score of at least 3 out of 10. Subjects who had any of the cases of patellar dislocation, patellar fracture, knee surgery in the past two years, knee ligament injuries, neurological diseases, pregnancy, tendonitis patella, Osgood-Schlatter disease, and Sinding Larsen disease were excluded.^{23,39,40} Before the treatment, there were no statistically significant differences between the groups in terms of age, height, weight, BMI, and duration of pain (Pvalue > 0.05) (table 1).

To facilitate the activation of the GMed, the KT with the NS-TEX (South Korea) was used lateral to the hip and on the GMed. To employ the KT, the subject lies sideways while the upper leg was on. For implementing the protocol, a KT 1 piece was used in the KT group. In that way, a third of the elemental type began without any traction from the posterior upper thrust. Then, the subject was asked to lie actively in a flexion position, while the femur was in the internal abduction and internal rotation, where a middle third of the tape, which was about 50% tensile, was drawn into the GMed muscle. While the subject's feet were in their original position, the remaining third of the tape was applied without any tensile action in the trochanter area of the femur. The first and second bars on the strip were used in the same way, except for the upper anterolateral throat. The subjects were asked to check the taping position for re-inspection. The KT protocol was based on previous studies in patients with pathologic pain syndrome⁴⁰ and was performed by a practitioner (figure 1).

In the placebo group, a tissue wound adhesive tape was used without tension. The similarity to KT, easy access, lack of elasticity, and its use in similar studies¹⁰ were among the reasons for choosing a scar tissue adhesive tape in this study. The wound adhesive was used precisely in places similar to the KT group. The pain was measured using a numerical pain rating scale in four conditions: walking, squatting, up, and downstairs. In the assessment of walking pain, the subject was asked to walk a distance of 50 meters on a straight path, then the level of pain on the walk was evaluated. To measure pain in the squat, the legs were positioned as wide as the shoulder and fingers, where squat running at a 90-degree angle was evaluated through visual assessment. To coordinate the subjects, ten consecutive repetitions were considered in 15 seconds.⁹ To assess the pain in the up and down stair movements, the subjects were asked to perform 12 steps up and down, and then, the pain was evaluated. In the four situations mentioned, the maximum pain was felt by the subject. The pain assessment protocol was selected based on previous studies.^{9,10,40} On the numerical scale of the pain, the zero means no pain while 4 to 6 signal moderate pain, and 10 represents the maximum possible pain. This scale has been used in previous studies in patients with PFPS.^{4,9,41} All steps of pain measurement were performed before, immediately and 24 hours after KT application. Each measurement was performed three times, and the mean of three measurements was used in the analysis.



Figure 1. GMed taping

SPSS software was employed for statistical analysis. Friedman's statistical test was utilized for comparison before and after the KT application at a significance level of 0.05 in the two groups.

Results

Table 1 reports the demographic characteristic of the subjects. Changes in the pain scores of the two groups in 4 situations; walking, squatting, climbing, moving downstairs,

immediately and 24 hours after the KT application are provided in table 2.

Table 1. Demographic characteristics of subjects

Variable	Placebo group (Mean ± SD)	KT group (Mean ± SD)	Pvalue
Age	23.13 ± 2.61	24.2 ± 1.65	0.193
Height	164.0 ± 0.26	162.53 ± 0.39	0.240
Weight	61.83 ± 3.63	60.66 ± 3.03	0.348
BMI (Kg)	23.00 ± 14.5	22.98 ± 1.31	0.975
Pain(0-10)	6.50 ± 1.45	6.90 ± 1.51	0.601

Table 2. Pain scores before, immediately and 24 after KT in three different situations

Task	Before KT(Mean ± SD)	After KT(Mean ± SD)	24 Hours after KT(Mean ± SD)	Pvalue
Walking				
- KT group	4.44 ± 0.80	2.00 ± 0.84	1.80 ± 0.56	0.000
- Placebo group	4.10 ± 0.87	3.83 ± 0.69	3.73 ± 0.70	*
Squat				
- KT group	6.13 ± 1.50	5.46 ± 1.18	4.40 ± 1.18	0.007
- Placebo group	6.83 ± 1.34	6.40 ± 0.91	6.00 ± 0.92	*
Step-up				
- KT group	6.43 ± 1.21	4.20 ± 1.01	4.06 ± 1.03	0.651
- Placebo group	5.60 ± 0.93	5.10 ± 0.66	5.06 ± 0.88	
Step-down				
- KT group	5.98 ± 1.27	3.60 ± 0.82	3.46 ± 1.24	0.000
- Placebo group	5.78 ± 1.00	5.60 ± 1.05	5.45 ± 0.99	*

The results indicate that time has had a significant effect in the walking task ($P = 0.000$, $F = 40.89$) for both groups (figure 1). Paired comparisons analysis showed a significant difference between before and immediately and 24 hours after KT application ($P = 0.000$), while there was no significant difference between immediately (Pvalue > 0.05) and 24 hours after KT use ($P = 0.345$).




Figure 1. Comparison of pain scores in the walking task in two groups

Mauchly test revealed that the assumed sphericity had been distorted. Thus, the results were corrected using Greenhouse Geisser indicating that over time, a significant effect was observed on squat ($P = 0.000$, $F = 22.66$ (figure 2). Paired comparisons analysis revealed a significant difference between before and immediately ($P = 0.003$), between immediately and 24 hours later, and between before and 24 hours after application of KT ($P = 0.000$).




Figure 2. Comparison of the pain score in a squatting task in two groups

As the results indicate, the time has had a significant effect on pain in the two groups in the stepping up task, ($Pvalue = 0.000$, $F = 34.37$) as shown in Figure. 3. Multiple comparisons showed that there was a significant difference between before and immediately and 24 hours after KT ($Pvalue=0.000$). There was, however, no significant difference between immediate application and 24 after KT ($Pvalue=0.06$).




Figure 3. Comparison of pain scores in the stepping up task in two groups

Based on the results, a significant effect of time on pain was observed in the two groups ($P = 0.000$, $P = 0.66$) (figure. 4). Multiple comparisons indicated that there was a significant difference between before and immediately after KT, before and 24 hours after KT ($P = 0.000$), while there was no significant difference between immediately and 24 after KT using ($P = 0.02$).




Figure 4. Comparison of pain scores in the stepping down task in the two groups

Discussion

In the present study, the effects of KT were investigated on pain in the athlete with PFPS and the following findings were observed: Pain intensity of the patients in walking, squat, and stepping up tasks diminished after the taping application.

Previous studies have confirmed pain reduction and functional activity improvement after the use of KT in patients with PFPS. The results of this study are inconsistent with previous similar studies.

In a study by Campolo et al., there was a significant difference between the KT group and the McConnell type group during a stepping up task in comparison to the control group. Although there was a significant difference between the three groups without taping, KT and McConnell taping, no significant difference was observed during the squat activity.⁴² Miller et al. compared the KT group of GMed, the lumbopelvic region, manual group, and the control group. They found that the KT group had significantly better Y-score and squat scores.⁴⁰ Reduction of pain and improvement in the ratio of VMO / VL EMG activity for people with PFPS in ascending step was reported by Chen et al.⁴³. The results of the study by Fu et al. revealed that the hip abduction improved with abductor KT. However, no increase in the power of GMed was confirmed following the KT in the Fu study.⁴⁴

In contrast, in the study by Aytar et al., despite the positive effects of KT on muscle strength and balance, there was no significant difference between the KT group and the placebo group in the severity of pain in walking up and downstairs.⁴⁵ Jancaitis and colleagues reported improvement in the performance of walking downstairs after the KT application, though no statistically significant difference was observed in the results.⁴⁶ Akbas et al. compared the reinforcement and flexibility exercises with KT and reinforcement exercises and flexibility alone. They observed that KT did not affect the pain reduction and improvement in subjects' performance.²

Based on scientific evidence, KT can have an immediate effect that help to reduce the symptoms of musculoskeletal

disorders.^{9,44} Generally, KT blocks the transfer of pain information to the spinal cord by stimulating the mechanical receptors of the skin, thereby reducing pain.²⁰ The application of KT on the skin surface provides tactile sensory inputs come from cutaneous mechanoreceptors in the skin that interact with the changes in the central nervous system's irritability with motor control. Local stimuli provided by KT replace muscle protections from the joint structure.¹¹ It is also likely that pain relief is a result of improvement in the direction and mechanical displacement of the femur and patella.^{9,43,45} As Powers describes, the weakness of the hip abductor muscles can be the cause of the hip prolapse. This situation would increase the Varus angle of the opposite knee; to compensate for the weakness of the abductor muscles, especially the GMed, the center of mass would change to the support foot, thereby increasing the angle of the knee valgus in the affected foot.⁴⁷ GMed mainly acts to stabilize the femur and pelvis in the frontal plane.⁴⁸ The GMed and gluteus Maximus muscles play an essential role in controlling the hip motions, especially in the subtraction torque on the frontal plate. Changes in the neuromuscular function of the GMed muscle may cause the internal rotation of the femur to rotate under the patella, resulting in abnormal deviation. To adapt to femur adduction, the tibia would be abducted which may cause excessive pronation.³⁷ The hip abductor KT may specifically facilitate the activity of GMed, thereby improving the postural stability and dynamic knee valgus.⁴⁰

Acknowledgement

We appreciate all participants who collaborated with us in this study.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Boling M, Padua D, Marshall S, Guskiewicz K, Pyne S, Beutler A. Gender differences in the incidence and prevalence of patellofemoral pain syndrome. *Scand J Med Sci Sports* 2010;20:725-30. doi:10.1111/j.1600-0838.2009.00996.x
- Petersen W, Ellermann A, Rembitzki IV, Scheffler S, Herbert M, Sprenger FS, et al. The Patella Pro study - effect of a knee brace on patellofemoral pain syndrome: design of a randomized clinical trial. *BMC Musculoskelet Disord* 2014;15:200. doi:10.1186/1471-2474-15-200
- Linschoten R, Middelkoop M, Berger MY, Heintjes EM, Koopmanschap MA, Verhaar JAN, et al. The PEX study - exercise therapy for patellofemoral pain syndrome: design of a randomized clinical trial in general practice and sports medicine. *BMC Musculoskelet Disord* 2006;7:31. doi:10.1186/1471-2474-7-31
- Bagheri S, Bayat MR, Halabchi F. The effect of 8-week exercise program on patellofemoral pain syndrome. *J Res Rehab Sci* 2011;7:357-67. [Persian].
- Bolgla LA, Malone TR, Umberger BR, Uhl TL. Hip strength and hip and knee kinematics during stair descent in females with and without patellofemoral pain syndrome. *J Orthop Sport Phys Ther* 2008;38:12-8. doi:10.2519/jospt.2008.2462
- Regelski CL, Ford BL, Hoch MC. Hip strengthening compared with quadriceps strengthening in conservative treatment of patients with patellofemoral pain: a critically appraised topic. *Int J Athl Ther Train* 2015;20:4-12. doi:10.1123/ijatt.2014-0048
- Kuru T, Yalman A, Dereli EE. Comparison of efficiency of Kinesio® taping and electrical stimulation in patients with patellofemoral pain syndrome. *Acta Orthop Traumatol Turc* 2011;46:385-92. doi:10.3944/aott.2012.2682
- Wen-Dien C, Fu-Chen C, Chia-Lun L, Hung-Yu L, Ping-Tung L. Effects of kinesio taping versus mcconnell taping for patellofemoral pain syndrome: a

- systematic review and meta-analysis. Evidence-Based Complement Altern Med 2015;1-11. doi:10.1155/2015/471208
9. Freedman SR, Brody LT, Rosenthal M, Wise JC. Short-term effects of patellar Kinesio taping on pain and hop function in patients with patellofemoral pain syndrome. Sports Health 2014;6:294-300. doi:10.1177/1941738114537793
 10. Aytar A, Ozunlu N, Surenkok O, Baltaci G, Oztop P, Karatas M. Initial effects of Kinesio® taping in patients with patellofemoral pain syndrome: a randomized, double-blind study. Isokinetic Exerc Sci 2011;19:135-42. doi:10.3233/IES-2011-0413
 11. Kase K, Wallis JKT. Clinical Therapeutic Applications of the Kinesio Taping Method. 3rd ed. Tokyo, Japan: Ken Ikai Co Ltd, 2003.
 12. Callaghan MJ, Selfe J, McHenry A, Oldham JA. Effects of patellar taping on knee joint proprioception in patients with patellofemoral pain syndrome. Man Ther 2008;13:192-9. doi:10.1016/j.math.2006.11.004
 13. González-Iglesias J, Fernández-de-Las-Peñas C, Cleland JA, Huijbregts P, Del Rosario Gutiérrez-Vega M. Short-term effects of cervical Kinesio taping on pain and cervical range of motion in patients with acute whiplash injury: a randomized clinical trial. J Orthop Sports Phys Ther 2009;39:515-21. doi:10.2519/jospt.2009.3072
 14. Thelen MD, Dauber JA, Stoneman PD. The clinical efficacy of Kinesio tape for shoulder pain: a randomized, double-blinded, clinical trial. J Orthop Sports Phys Ther 2008;38:389-95. doi:10.2519/jospt.2008.2791
 15. Castro-Sánchez AM, Lara-Palomo IC, Matarán-Peñarocha GA., Fernández-Sánchez M, Sánchez-Labraca N, Arroyo-Morales M. Kinesio Taping reduces disability and pain slightly in chronic non-specific low back pain: a randomized trial. J Physiother 2012;58:89-95. doi:10.1016/S1836-9553(12)70088-7
 16. Crossley K, Cowan SM, Bennell KL, McConnell J. Patellar taping: is clinical success supported by scientific evidence? Man Ther 2000;5:142-50. doi:10.1054/math.2000.0354
 17. Crossley K, Bennell K, Green S, Cowan S, McConnell J. Physical therapy for patellofemoral pain: a randomized, double-blinded, placebo-controlled trial. Am J Sports Med 2002;30:857-65. doi:10.1177/03635465020300061701
 18. Crossley K, Bennell K, Green S, McConnell J. A systematic review of physical interventions for patellofemoral pain syndrome. Clin J Sport Med 2001;11:103-10. doi:10.1097/00042752-200104000-00007
 19. Baluchi R, Giasi A, Naderi A. A survey of selective movement therapy effectiveness on dynamic postural control of patients with patellofemoral pain syndrome. SJIMU 2011;19:17-23.
 20. Bagheri S, Shojaeddin S, Nazarian A B, Naderi A. The effect of hip abductors and external rotators strengthening in male with patellofemoral pain syndrome. SJIMU 2016;23:29-39.
 21. Whittingham M, Palmer S, Macmillan F. Effects of taping on pain and function in patellofemoral pain syndrome: a randomized controlled trial. J Orthop Sports Phys Ther 2004;34:504-10. doi:10.2519/jospt.2004.34.9.504
 22. Mokhtarinia H, Ebrahimi-Takamjani I, Salavati M, Goharparay S, Khosravi A. The effect of patellar taping on knee joint proprioception in patients with patellofemoral pain syndrome. Acta Med Iran 2008;46:183-90.
 23. Akbaş E, Atay AÖ, Yüksel I. The effects of additional Kinesio taping over exercise in the treatment of patellofemoral pain syndrome. Acta Orthop Traumatol Turc 2011;45:335-41. doi:10.3944/AOTT.2011.2403
 24. Pal S, Besier TF, Draper CE, Fredericson M, Gold GE, Beaupre GS, et al. Patellar tilt correlates with vastus lateralis: vastus medialis activation ratio in maltracking patellofemoral pain patients. J Orthop Res 2012;30:927-33. doi:10.1002/jor.22008
 25. Souza RB, Draper CE, Fredericson M, Powers CM. Femur rotation and patellofemoral joint kinematics: a weight-bearing magnetic resonance imaging analysis. J Orthop Sports Phys Ther 2010;40:277-85. doi:10.2519/jospt.2010.3215
 26. Powers CM, Ward SR, Fredericson M, Guillet M, Shellock FG. Patellofemoral kinematics during weight-bearing and non-weight-bearing knee extension in persons with lateral subluxation of the patella: a preliminary study. J Orthop Sports Phys Ther 2003;33:677-85. doi:10.2519/jospt.2003.33.11.677
 27. Fukuda TY, Rossetto FM, Magalhães E, Bryk FF, Lucareli PR, de Almeida Aparecida Carvalho N. Short-term effects of hip abductors and lateral rotators strengthening in females with patellofemoral pain syndrome: a randomized controlled clinical trial. J Orthop Sports Phys Ther 2010;40:736-42. doi:10.2519/jospt.2010.3246
 28. Earl JE, Hoch AZ. A proximal strengthening program improves pain, function, and biomechanics in women with patellofemoral pain syndrome. Am J Sports Med 2011;39:154-63. doi:10.1177/0363546510379967
 29. Peters JS, Tyson NL. Proximal exercises are effective in treating patellofemoral pain syndrome: a systematic review. Int J Sports Phys Ther 2013;8:689-700.
 30. Magalhães E, Fukuda TY, Sacramento SN, Forgas A, Cohen M, Abdalla RJ. A comparison of hip strength between sedentary females with and without patellofemoral pain syndrome. J Orthop Sports Phys Ther 2010;40:641-7. doi:10.2519/jospt.2010.3120
 31. Prins MR, van der Wurff P. Females with patellofemoral pain syndrome have weak hip muscles: a systematic review. Aust J Physiother 2009;55:9-15. doi:10.1016/s0004-9514(09)70055-8
 32. Ireland ML, Willson JD, Ballantyne BT, Davis IM. Hip strength in females with and without patellofemoral pain. J Orthop Sports Phys Ther 2003;33:671-6. doi:10.2519/jospt.2003.33.11.671
 33. Nakagawa TH, Moriya ÉT, Maciel CD, Serrão FV. Trunk, pelvis, hip, and knee kinematics, hip strength, and gluteal muscle activation during a single-Leg squat in males and females with and without patellofemoral pain syndrome. J Orthop Sport Phys Ther 2012;42:491-501. doi:10.2519/jospt.2012.3987
 34. Cowan SM, Crossley KM, Bennell KL. Altered hip and trunk muscle function in individuals with patellofemoral pain. Br J Sports Med 2009;43:584-8. doi:10.1136/bjism.2008.053553
 35. Barton CJ, Lack S, Malliaras P, Morrissey D. Gluteal muscle activity and patellofemoral pain syndrome: a systematic review. Br J Sports Med 2013;47:207-14. doi:10.1136/bjsports-2012-090953
 36. Chaudhari AM, Andriacchi TP. The mechanical consequences of dynamic frontal plane limb alignment for non-contact ACL injury. J Biomech 2006;39:330-8. doi:10.1016/j.jbiomech.2004.11.013
 37. Powers CM. The influence of altered lower-extremity kinematics on patellofemoral joint dysfunction: a theoretical perspective. J Orthop Sports Phys Ther 2003;33:639-46. doi:10.2519/jospt.2003.33.11.639
 38. Lephart SM, Ferris CM, Riemann BL, Myers JB, Fu FH. Gender differences in strength and lower extremity kinematics during landing. Clin Orthop Relat Res 2002;401:162-9. doi:10.1097/00003086-200208000-00019
 39. Osorio JA, Vairo GL, Rozea GD, Boshia PJ, Millard RL, Aukerman DF, et al. The effects of two therapeutic patellofemoral taping techniques for strength, endurance, and pain responses. Phys Ther Sport 2013;14:199-206. doi:10.1016/j.ptsp.2012.09.006
 40. Miller J, Westrick R, Diebal A, Marks C, Gerber JP. Immediate effects of lumbopelvic manipulation and lateral gluteal kinesio taping on unilateral patellofemoral pain syndrome: a pilot study. Sports Health 2013;5:214-9. doi:10.1177/1941738112473561
 41. Iverson CA, Sutlive TG, Crowell MS, Morrell RL, Perkins MW, Garber MB, et al. Lumbopelvic manipulation for the treatment of patients with patellofemoral pain syndrome: development of a clinical prediction rule. J Orthop Sports Phys Ther 2008;38:297-309. doi:10.2519/jospt.2008.2669
 42. Campolo M, Babu J, Dmochowska K, Scariah S, Varughese J. A comparison of two taping techniques (Kinesio and McConnell) and their effect on anterior knee pain during functional activities. Int J Sports Phys Ther 2013;8:105-10.
 43. Chen PL, Hong WH, Lin CH, Chen WC. Biomechanics effects of Kinesio taping for persons with patellofemoral pain syndrome during stair climbing. IFMBE Proc 2008;21:395-7. doi:10.1007/978-3-540-69139-6_100
 44. Fu TC, Wong AM, Pei YC, Wu KP, Chou SW, Lin YC. Effect of Kinesio taping on muscle strength in athletes-a pilot study. J Sci Med Sport 2008;11:198-201. doi:10.1016/j.jsams.2007.02.011
 45. Aytar A, Ozunlu N, Surenkok O, Baltaci G, Oztop P, Karatas M. Initial effects of Kinesio® taping in patients with patellofemoral pain syndrome: a randomized, double-blind study. Isokinetic Exerc Sci 2011;19:135-42. doi:10.3233/IES-2011-0413
 46. Freedman SR, Brody LT, Rosenthal M, Wise JC. Short-term effects of patellar kinesio taping on pain and hop function in patients with patellofemoral pain syndrome. Sports health 2014;6:294-300. doi:10.1177/1941738114537793
 47. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. J Orthop Sports Phys Ther 2010;40:42-51. doi:10.2519/jospt.2010.3337
 48. Neumann DA. Kinesiology of the musculoskeletal system. 3rd ed. St Louis, MO: Mosby, 2002. p. 1-624.