

The Factors Influencing Sleep Quality and Its Related Disorders in Process, Manufacturing, and Service Industries: A Systematic Review

Soleiman Ramezanifar¹, Ehsan Ramezanifar², Ali Salehi Sahlabadi^{3*}

¹ Occupational Health and Safety Engineering, Department of Occupational Health and Safety Engineering, School of Public Health and Safety, Shahid Beheshti University of Medical Sciences, Tehran, Iran

² Center of Excellence for Occupational Health and Research, Center of Health Sciences, Hamadan University of Medical Sciences, Hamadan, Iran

³ Occupational Health and Safety Engineering, Department of Occupational Health and Safety Engineering, School of Public Health and Safety, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Received: 01 Sep. 2021 Accepted: 01 Oct. 2021

Abstract

Background and Objective: Numerous factors in industrial environments affect employees' sleep quality and cause poor sleep quality and its related disorders. This study was conducted to investigate factors influencing sleep quality and its related disorders in the process, manufacturing, and service industries.

Materials and Methods: In this study, the related articles published in the Embase, PubMed, Google Scholar, Scopus, Web of Science, Magiran, SID, IranMedex, and IranDoc databases from 2010 to 2021 were reviewed. Some of the used search terms were "sleep", "sleep quality", "sleep disorder", "process industry", "manufacturing industry", and "service industry". searched articles were reviewed to extract the required data.

Results: Out of the 431 studies identified, 59 articles were eligible for analysis, of which 19 studies were conducted in process industries, 35 studies in service industries, and 5 studies in manufacturing industries. In process industries, factors such as anxiety, depression, shift work, job stress, and exposure to high noise and vibration reduce workers' sleep quality. In the manufacturing industry, work stress, and shift work impair sleep quality. Inadequate sleep quality was common in the service industry, and several factors, including depression, job stress, shift work, and musculoskeletal disorders play a role in this regard.

Conclusion: The results indicate that working conditions are such that people's sleep can be affected by various factors, so more attention should be paid to sleep health. Shift plans should be implemented following the best method. Moreover, it is critical to focus on sleep quality in the workplace.

Keywords: Sleep quality; Sleep disorders; Sleepiness; Circadian rhythm disorders; Manufacturing industry

Citation: Ramezanifar S, Ramezanifar E, Salehi Sahlabadi A. **The Factors Influencing Sleep Quality and Its Related Disorders in Process, Manufacturing, and Service Industries: A Systematic Review.** *J Sleep Sci* 2021; 6(3-4): 53-66.

Introduction

Maintaining a healthy workforce is the cornerstone of any prosperous society. All individuals' essential characteristics must be considered (1). Sleep is one of these fundamental human characteristics that significantly affect people's health.

Sleep is a dynamic and well-structured biological activity and one of the main aspects of the entire 24-hour life cycle, which is considered an essential part of life and is associated with restoring physical and mental strength (2). If people have a good night's sleep, their performance and alertness will increase throughout the day (3). This increase in awareness and performance impacts individuals' daily life, especially in their work environments, and increases their productivity in the workplace (4).

* **Corresponding author:** A. Salehi Sahlabadi, Occupational Health and Safety Engineering, Department of Occupational Health and Safety Engineering, School of Public Health and Safety, Shahid Beheshti University of Medical Sciences, Tehran, Iran
Tel: +98 21 22432040, Fax: +98 21 22432037
Email: asalehi529@sbumu.ac.ir

Copyright © 2021 Iranian Sleep Medicine Society, and Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Noncommercial uses of the work are permitted, provided the original work is properly cited.

Inadequate sleep quality can decrease mental function, degrade cognition and decision-making and increase error (5). Poor sleep quality increases daily absenteeism from work and increases the frequency of accidents (6). Findings of a cross-sectional study conducted on 200 drivers of heavy vehicles in Iran showed that the prevalence of poor sleep quality is higher in drivers who have a history of accidents (7). Many individual and environmental factors such as age, circadian rhythm, work schedule, and sleep pattern can affect sleep quality (8).

If poor sleep quality persists, people's sleep and its regular rhythm will be affected, this will bring about the conditions for sleep disorders (9). Sleep disorder is a perturbation in the natural cycle of sleep and arousal with adverse effects such as depression, anxiety, lethargy and distraction, and loss of consciousness (10, 11). Sleep disorders affect people in various ways and reduce their quality of life and work performance (12). According to the results of a study conducted in Iran, sleep disorders have adverse effects on the performance of people in the workplace (13). In addition, the findings of a prospective cohort study in the United States showed that sleep disorders and mood problems of the hospital personnel were independently associated with adverse outcomes and further increased risk (10).

Today, industries are divided into the 3 main categories of process, manufacturing, and service industries (14). In each of these industries, people's sleep status is of different importance depending on the nature of their job. In many jobs, long working hours, short rest times, and shift work can affect employees' sleep (15, 16). Many of these factors cause sleep deprivation in employees in various occupations. Lack of sleep causes cognitive function problems such as lower concentration, longer reaction time, lower capacity to acquire and recall new knowledge, and impaired motor skills (17). With the emergence of these problems, people's health, safety, and productivity in these industries are affected (12).

Therefore, it is necessary to study the factors affecting sleep quality and the related disorders in different industries according to their nature. In addition, according to the searches conducted by the authors of this study, no review study has been conducted in this field. For this purpose, the present study was conducted to investigate the factors influencing sleep quality and its related

disorders in the process, manufacturing, and service industries.

Materials and Methods

Search strategy: The two authors used the Embase, PubMed, Google Scholar, Scopus, Web of Science, Magiran, SID, IranMedex, and IranDoc databases to search for related articles. Research articles published in English and Persian from 2010 to 2021 in these valid databases were retrieved. Keywords used for the research included "sleep", "sleep quality", "sleep disorder", "sleep problem", "sleep disturbance", "sleep health", "sleep quality", "sleepiness", "drowsiness", "circadian rhythm disorder", "process industry", "manufacturing industry", and "service industry".

Study selection and data extraction: The two authors independently reviewed the search results and screened the qualified studies for full-text review. All studies that explicitly examined the factors affecting sleep quality and its related disorders in the process, manufacturing, and service industries between 2010 and 2021 were included in this study. However, non-research articles such as editorials, authors' notes, letters to the editor, standard texts, and studies not written in Persian and English were excluded. To extract data from the reviewed articles, the authors used a form that contained items on study design, year of publication, country, the industry studied, sleep measurement tool, sample size (number, gender, and age), and results. Figure 1 shows the general process of choosing articles.

Evaluation criteria for the quality of articles: The Joanna Briggs Institute (JBI) checklist was used to evaluate the quality of the articles (18). This checklist has 9 items with the responses "yes", "no", "unspecified", and "not applicable". This checklist determines the methodological quality of studies and ways to acquire and identify errors in studies, design, and data analysis. The Preferred Reporting Items for Systematic reviews and Meta-Analyses for Protocols 2015 (PRISMA-P 2015) was also applied to write the article itself. This tool includes a 17-item checklist for the preparing and describing of a robust protocol for the systematic review (19).

Results

The process of article selection is presented in figure 1.




Figure 1. The general process of selecting articles

Out of 79 selected articles, 59 studies were finally selected for use in this article. Out of the 59 studies reviewed, 19 studies (32.2%) were conducted in process industries (4, 20-37), 35 studies (59.3%) in service industries (38-72), and 5 studies (8.5%) in manufacturing industries (13, 73-76).

According to tables 1, 2, and 3, of the 59 selected studies, 45 studies (76.3%) were cross-sectional, and 14 studies (23.7%) were conducted with other study designs. In addition, among the studies, 22 studies (37.3%) were performed in Iran, 11 studies (18.7%) in the United States, and 26 studies (44%) in other countries (Figure 2). Moreover, the share of Asia was 35 studies

(59.3%), North America 12 studies (20.3%), and other continents 11 studies (18.6%).

These studies were conducted on 43271 people, 8312 women (19.3%), 34656 men (80%), and 303 unknown (0.7%) participants. Of these, 21048 (48.6%) participated in research related to sleep quality and 22223 (51.4%) in research related to sleep disorders.

Among the tools used to assess participants' sleep status, the Pittsburgh Sleep Quality Index (PSQI), Epworth Sleepiness Scale (ESS), and Berlin Questionnaire (BQ) were the most widely used tools, respectively (Figure 3). The PSQI was used in 69.5% of articles, ESS in 23.7% of articles, and BQ in 11.9% of articles.




Figure 2. Distribution of articles by country (2010-2021)

Table 1. A summary of studies between 2010 and 2020 that examine factors influencing sleep quality and its related disorders in the process industry

Study design	Country (Year)	Industry	Sleep measurement tool	Sample size	Result(s)
Sleep disorders					
Cross-sectional (23)	Iran (2021)	Petroleum industry	ESS, ISI	188 men, mean age: 37.06 ± 9.2 years	- SWD is regarded as a substantial health issue among oil rig shift workers. - SWD is related to daytime sleepiness, insomnia, and depression.
Cross-sectional (33)	Iran (2018)	Petrochemical industry	KSS and employees' periodic medical examination records	501 employees: 477 men and 24 women, 22-70 years	- Participants with sleep disturbances had significantly higher scores on various fatigue dimensions when compared to those without sleep disturbances.
Case-Control (20)	Iran (2013)	Textile Factory	PSQI, BQ, ESS, ISI, and RLSQ	400 male workers: 196 shift workers (mean age: 33.9 ± 5.9 years) and 204 non-shift workers (mean age: 35.1 ± 10.9 years)	Shift workers had substantially higher insomnia and poor sleep quality rates than non-shift workers.
Cross-sectional (21)	Iran (2017)	Steel Factory	PSQI	282 men, mean age: 32.21 ± 6.57 years	- No link was found between sleep disturbances and workplace accidents.
Case Study (37)	Iran (2015)	Wind Farm	ESS	53 men, mean age: 30.8 years	- There was a significant association between age, employees' experience, equivalent sound level, and sleep disorders.
Sleep quality					
Cross-sectional (24)	China (2020)	Petroleum industry	PSQI	2116 workers: 1020 men and 1096 women, 18-60 years	- Workplace stress has a detrimental impact on workers' sleep quality. - The CLOCK and Per3 genes may increase the risk of sleep problems.
Cross-sectional (27)	China (2019)	Petroleum industry	PSQI	2116 workers: 1020 men & 1096 women, 18-60 years	- High job stress increases the likelihood of sleep problems.
Cross-sectional (34)	Iran (2019)	Power plant	PSQI	275 male workers [exposed group (n = 132) and unexposed group (n = 143)], < 30 years (n = 88), 30-40 years (n = 127), ≥ 40 years (n = 48)	- Prolonged industrial ELF-EMF exposure may result in anxiety, depression, stress, and poor sleep quality.
Ex Post Facto (31)	Iran (2017)	Gas refinery	PSQI	255 male workers, 20-60 years	- In terms of sleep disturbances, there was no significant difference between shift workers and day workers. However, the differences in sleep latency and sleep efficiency between the two groups were significant.
Descriptive (35)	Iran (2019)	Gas refinery	PSQI	180 men, 23-59 years	- There was a significant relationship between occupational stress and quality of sleep.
Cross-sectional (22)	Turkey (2018)	Textile Factory	PSQI	100 male workers	- The length of working in the textile industry affects sleep quality metrics.
Case-Control (25)	Iran (2014)	Petrochemical industry	PSQI	62 men: case group (n = 40), control group (n = 22)	- There was no significant association between exposure level, quality of sleep, and overall health.
Quasi-experimental research with a pretest-posttest design (26)	Iran (2014)	Petrochemical industry	PSQI	90 men: (45 case and 45 control), 30-58 years	- Stress management training enhanced staff members' mental health, career satisfaction, and sleep quality.

Table 1. A summary of studies between 2010 and 2020 that examine factors influencing sleep quality and its related disorders in the process industry (continue)

Study design	Country (Year)	Industry	Sleep measurement tool	Sample size	Result(s)
Cross-sectional (30)	Iran (2016)	Petrochemical industry	ESS	180 men, 18-53 years	- Drowsiness, work overload, and work situation awareness had a significant internal relation.
Cross-sectional (36)	Iran (2016)	Steel Factory	PSQI	256 men, mean age: 33.6 ± 6.9 years	- There was a positive and clear link between sleep quality and the overall health of shift workers.
Cross-sectional (32)	Brazil (2018)	Mining	PSQI	89 men, mean age: 32.9 ± 6.3 years	- Positive correlations were found among self-perception of health, and duration and quality of sleep. - Anxiety, and sleep length and quality were found to be related.
Cross-sectional (28)	Canada (2017)	Mining	ESS, KSS, and Actigraph	19 male underground miners, 41.5 ± 5.1 years	-The miners' efficiency decreased overnight.
Cross-sectional (29)	Canada (2020)	Mining	Actigraph	89 male miners: 38 night shift (mean age: 36.8 ± 13.9 years) and 51 day shift (mean age: 37.1 ± 12.8 years)	Despite restricted circadian acclimation in terms of nocturnal vigilance, underground miners have fine sleep quality.
Cross-sectional (4)	Germany (2018)	Wind Farm	PSQI	268 men, 20-34 years (n = 116), 35-49 years (n = 122), ≥ 50 years (n = 30)	- Higher levels of noise exposure, vibrations, and artificial ventilation are related to poor sleep quality.

ESS: Epworth Sleepiness Scale; ISI: Insomnia Severity Index; SWD: Shift Work Disorder; KSS: Karolinska Sleepiness Scale; PSQI: Pittsburgh Sleep Quality Index; BQ: Berlin Questionnaire; RLSQ: Restless Legs Syndrome Questionnaire; ELF-EMF: Extremely Low-Frequency Electric and Magnetic Fields

Table 2. A summary of studies between 2010 and 2020 that examine factors influencing sleep quality in the manufacturing industry

Study design	Country (Year)	Industry	Sleep measurement tool	Sample size	Result(s)
Cross-sectional (74)	Malaysia (2018)	Cable manufacturing	PSQI	140 men, 21-57 years	- A greater level of all aspects of work stress and a higher level of avoidance were associated with poor sleep quality.
Cross-sectional (73)	Korea (2017)	Electronics Manufacturing industry	PSQI	1472 workers: 1178 men and 294 women, 19-39 years	Serum vitamin D insufficiency was found to strongly correlate with poor sleep quality.
Descriptive-Analytical (75)	Iran (2012)	Automobile industry	ESS	120 male workers: 60 fixed morning shifts (31.16 ± 5.55 years) and 60 night rotation shifts (31.83 ± 7.29 years)	- Sleepiness and shifts are harmful to the general health of workers.
Case Study (76)	Taiwan (2010)	Semi-conductor Manufacturing Company	PSQI	239 worker: 198 men and 41 women, 21-50 years	- The effects of varying work shifts on sleep quality and career efficiency were substantial.
Cross-sectional (13)	Iran (2014)	Car accessories manufacturing plant	MUSIC-NQ	727 male workers, 21-63 years	- There was a strong link between sleep disorders and workability.

PSQI: Pittsburgh Sleep Quality Index; ESS: Epworth Sleepiness Scale; MUSIC-NQ: Musculoskeletal Intervention Center-Norrtalje Questionnaire

Table 3. A summary of studies between 2010 and 2020 examining factors influencing sleep quality and its related disorders in the service industry

Study design	Country (Year)	Industry	Sleep measurement tool	Sample size	Result(s)
Sleep disorders					
Pilot (58)	USA (2013)	Nursing	BQ and PSG	21 women, mean age: 40 ± 13 years	- According to the PSG, the prevalence of sleep-disordered breathing was 43%, but only 24% of these cases were regarded as high risk according to the BQ.
Cross-sectional (38)	Iran (2016)	Nursing	PSQI	1456 nurses: 1296 women and 160 men, 19-60 years	- Sleep troubles have a remarkable impact on health-related quality of life.
Cross-sectional (70)	Iran (2019)	Nursing	PSQI, ESS, and ISI	100 nurse: 49 women and 51 men, mean age: 30.4 ± 7.5 years	- Sleep disturbances were more common in nurses working in the surgical ward than those in other parts of the hospital.
Descriptive cross-sectional (45)	Egypt (2014)	Nursing	The two-week sleep diary	52 nurse: 51 women and 1 man, 19-34 years	- The rate of errors peaked with unusual shift work patterns and night sleep patterns in nurses who had worked more than 4 nights in the previous 2 weeks.
Cross-sectional (71)	USA (2015)	Firefighting	BQ, AIS, RLSQ, and SWD screening tool	6933 firefighters: 6360 men & 410 women & 163 unknown, mean age: 40.4 ± 8.9 years	- Sleep disturbances are common among firefighters and are linked to an increased risk of adverse health and safety consequences.
Cross-sectional (39)	USA (2019)	Firefighting	BQ, AIS, RLSQ, ESS	6037 firefighters: 5812 men, 372 women, 123 unknown, mean age: 40.4 ± 8.9 years	- Sleep and mental health troubles were linked to an elevated risk of burnout in firefighters.
Cross-sectional (42)	USA (2019)	Firefighting	PSQI	652 firefighters: 611 men and 36 women, and 5 transgender people, mean age: 38.4 ± 8.6 years	- High alcohol abuse and low distress tolerance were significantly associated with high sleep disturbance.
Cross-sectional (41)	USA (2020)	Firefighting	PSQI, Micro Sleep Watch actigraphs	45 firefighters: 44 men and 1 women, mean age: 37.4 ± 7.22 years	- Even minor sleep disruptions impacted cognitive performance, thus increasing the probability of poor job performance or injury.
Cross-sectional (44)	Iran (2020)	Driving	ESS	281 male locomotive drivers, mean age: 43 ± 10 years	- 166 drivers had 2 or more OSA risk factors, and there was a high rate of OSA among locomotive drivers.
Cross-sectional and prospective cohort (48)	USA (2011)	Police officers	BQ, AIS, RLSQ, CQ, ESS, and PSG	4957 police officers: 861 women and 4079 men, and 17 unknown. 20-77 years	- Sleep disturbances were significantly linked to a higher risk of self-reported detrimental health, performance, and safety consequences.
Sleep quality					
Cross-sectional (63)	Taiwan (2013)	Nursing	PSQI, A take-home Holter ECG SD-100 portable recorder	156 women, mean age: 34.6 ± 8.1 years	- Subjective and objective findings showed that hospital nurses suffered from poor sleep quality.
Descriptive-comparative (57)	Iran (2011)	Nursing	PSQI	180 nurses: 14 men and 166 women, mean age: 31.6 ± 8.5 years	- There was a statistically significant difference in the quality of sleep and the overall health of nursing staff.
Cross-sectional (50)	Iran (2014)	Nursing	PSQI	200 nurses: 175 women and 25 men, mean age: 32.9 ± 5.7 years	- With increased job stress, nurses' sleep quality deteriorates, and their fatigue worsens.
Cross-sectional (62)	China (2020)	Nursing	PSQI	155 women, 21-54 years	- Work-related stress harms sleep quality.

Table 3. A summary of studies between 2010 and 2020 examining factors influencing sleep quality and its related disorders in the service industry (continue)

Study design	Country (Year)	Industry	Sleep measurement tool	Sample size	Result(s)
Quantitative, cross-sectional, descriptive, and comparative (47)	Brazil (2010)	Nursing	PSQI	203 nurses: 179 women and 24 men, < 30 years (n = 31), 30-40 years (n = 65), 40-49 years (n = 85), ≥ 50 years (n = 22)	- There was a significant relationship between sleep and stress.
Cross-sectional (65)	Singapore (2010)	Nursing	KSS	163 nurses: 147 women and 16 men, mean age: 32.8 ± 6.9 years	- Over 70% of nurses said they did not get enough sleep. Older age, poor sleep situation, digestive symptoms, and higher strain and symptom levels were all risk factors for inadequate sleep.
Cross-sectional (43)	Taiwan (2010)	Nursing	PSQI	435 women, 20-52 years	- There was an inverse relationship between sleep quality scores and overall quality of life scales.
Cross-sectional (49)	Korea (2018)	Nursing	PSQI	188 nurses: 182 women and 6 men, mean age: 30.84 ± 7.08 years	Sleep problems and subjective sleep quality were significant predictors of nurse productivity, shift work, and age.
Cross-sectional (72)	Iran (2018)	Firefighting	PSQI	118 men, mean age: 33.2 ± 4.9 years	- Poor sleep quality was associated with musculoskeletal problems, shift work, and a high BMI in firefighters.
Descriptive (67)	USA (2011)	Firefighting	PSQI, ESS	112 firefighters: 107 men and 5 women, mean ages: 43.6 ± 7.7 years	- There was an association between sleep deprivation, depressed mood, physical/mental health, and drinking habits.
Cross-sectional (40)	Australia (2016)	Firefighting	Wrist actigraph	40 firefighters: 31 men and 9 women, 19-65 years	- The amount of sleep achieved during shift work times was limited, and pre-and post-sleep fatigue scores were more significant than non-shift work times.
Cross-sectional (69)	USA (2016)	Firefighting	PSQI	109 male career firefighters, 20-59 years	- Firefighters who work second jobs have significantly poorer sleep quality.
Cross-sectional (68)	Belgium (2011)	Driving	ESS, PSQI, and BQ	476 men, mean age: 42.7 ± 10.2 years	- Lack of sleep and daytime sleepiness were common among truck drivers.
Cross-sectional (46)	Iran (2016)	Driving	PSQI	1500 men, 22-72 years	- Experienced drivers had a high rate of poor sleep quality.
Cross-sectional (66)	Thailand (2019)	Driving	PSQI, BQ	338 men, mean age: 45.8 ± 8.1 years	- Poor sleep quality was linked to night shift work, night or day shift patterns, consumption of alcohol, marriage, and lack of physical activity.
Cross-sectional (60)	Iran (2012)	Driving	PSQI	100 men, 26-63 years	- There was a tangible link between sleep quality, age, and work satisfaction.
Cross-sectional (55)	Korea (2014)	Driving	PSQI	4634 train drivers: 4597 men and 37 women, mean age: 45.3 ± 9.4 years	- Human errors were significantly associated with inadequate sleep quality and posttraumatic stress disorder.
Cross-sectional (53)	Australia (2018)	Driving	ESS, PSQI, KSS, and EEG	63 train drivers: 54 men and 9 women, 24-69 years	- Train drivers' subjective fatigue/sleepiness levels were predicted with different levels of success.
Cross-sectional (59)	USA (2016)	Policing	PSQI	363 police officers: 260 men and 103 women, 27-66 years	- Evening and night work schedules are linked to an increased prevalence of inadequate sleep quality in police officers.
Cross-sectional (64)	Taiwan (2015)	Policing	PSQI	796 male police officers, mean age: 37.36 ± 7.73 years	- Police officers who had more sleep disturbances were more likely to have metabolic syndrome and abdomen obesity.

Table 3. A summary of studies between 2010 and 2020 examining factors influencing sleep quality and its related disorders in the service industry (continue)

Study design	Country (Year)	Industry	Sleep measurement tool	Sample size	Result(s)
Cross-sectional (61)	Australia (2016)	Policing	ESS, PSQI	206 police officers: 140 men and 66 women, mean age: 31.63 ± 8.51 years	- In the survey, bad sleeping quality and fatigue intensity were found to be prevalent.
Cross-sectional (51)	USA (2019)	Policing	PSQI	356 police officers: 256 men and 100 women, mean age: 41.3 ± 6.7 years	- Police officers' perceptions of the severity of stress and the rate of stress factors are linked to poor sleep quality.
Cross-sectional (52)	USA (2011)	Policing	PSQI	391 police officers: 284 men and 107 women, mean age: 40.70 ± 7.14 years	- Quality of sleep was found to be noticeably and separately linked to symptoms of depression.
Cross-sectional (54)	Netherlands (2015)	Policing	A seven-dimensional questionnaire	740 police force: 252 women and 488 men, mean age: 35.7 ± 10.6 years	- Sleep quality was the most crucial factor in shift work tolerance, followed by the need for fatigue, recovery, and work-life balance.
Cross-sectional (56)	Netherlands (2021)	MRI Technicians	The 12-item Medical Outcomes Study sleep scale	233 technicians: 186 women and 47 men, mean age: 42.56 ± 10.6 years	- Contact with MRI-related magnetic fields tends to be related to elevated sleep disruption and non-optimal sleep length.

BQ: Berlin Questionnaire; PSQI: Pittsburgh Sleep Quality Index; ISI: Insomnia Severity Index; AIS: Athens Insomnia Scale; RLSQ: Restless Legs Syndrome Questionnaire; SWD: Shift Work Disorder; ESS: Epworth Sleepiness Scale; CQ: Cataplexy Questionnaire; PSG: Polysomnography; KSQ: Karolinska Sleep Questionnaire; KSS: Karolinska Sleepiness Scale; EEG: Electroencephalography; MRI: Magnetic resonance imaging




Figure 3. Distribution of sleep measurement tools in articles (2010-2021)

In the process industries, factors such as anxiety, depression, shift work, job stress, and exposure to high noise and vibration reduce workers' sleep quality. People with sleep disorders also suffer from more fatigue (4, 20, 23, 27, 33, 35, 37). In addition, exposure to Extremely Low-Frequency Electric and Magnetic Fields (ELF-EMF) may reduce sleep quality (34). Most of the studies conducted in these industries were in the petrochemical, petroleum, and mining industries (Figure 4).




Figure 4. Distribution of articles in process industries (2010-2021)

Work stress and shift work impair sleep quality in the manufacturing industries and reduce job productivity and workers' general health (74-76). In addition, vitamin D deficiency has been suggested to influence sleep quality (73).

In service occupations, such as nursing, sleep quality is related to general health, and job stress affects sleep quality and fatigue (47, 50, 62, 63). In addition, in jobs such as firefighting, factors such as shift work, musculoskeletal disorders, sleep deprivation, depression and working in a second job are also associated with poor sleep quality (40, 67, 69, 72). Inadequate sleep quality is common among drivers, and factors such as night work, poor sleep patterns, alcohol consumption, and lack of physical activity are the cause (46, 66, 68). Poor sleep quality and fatigue are also common among police officers, are more evident in the evening and night work, and are affected by factors such as stress and depression (51, 52, 59, 61). Most of the studies conducted in these industries have been in the nursing and firefighting professions (Figure 5).




Figure 5. Distribution of articles in service occupations (2010-2021)

The distribution of published articles assessing sleep quality and its related disorders by year of publication is shown in figure 6.




Figure 6. Distribution of articles by year of publication

The trend of publishing articles in these 10 years shows the importance of the issue of sleep in the industries.

Discussion

There are several variables in the process, manufacturing, and service industries that affect sleep quality. These variables reduce sleep quality and increase sleep disorders (35). Shift work is one of the main factors affecting sleep quality and its related disorders (77). Shift work is inevitable in many industries due to the requirements and nature of work. The concept of shift work mainly refers to scheduling daily work schedules. In shift jobs, individuals work outside the usual 8 hours a day (78). Problems such as gastrointestinal diseases, diabetes, infectious diseases, cardiovascular disorders, musculoskeletal disorders, and sleep disorders are common in shift workers (79). According to a cross-sectional study conducted on 2,453 employees of the China Petroleum and Xinjiang Petrochemical Company, shift workers had more sleep disorders than non-shift workers (80). In addition, the results of a case study on 239 workers at a semiconductor manufacturing company in Taiwan suggested that different shifts had a significant effect on sleep quality and work quality (76).

Job stress is another component that can affect sleep quality and its related disorders. Occupational stress is a type of psychological stress that occurs due to workers' psychological, physical, and social pressures (81). Problems such as cardiovascular diseases, stomach diseases, digestive issues, headaches, and sleep problems appear as a result of occupational stress (35). Occupational stress is a component that is strongly related to and affects sleep quality. A descriptive study on 180 male employees of Behbahan Bid Boland gas refinery in Iran showed a strong association between job stress and sleep quality (35). Prolonged stress can increase the number of self-regulatory reactions during the night and reduce sleep quality (27).

The impact of factors influencing sleep quality and its related sleep disorders in different industries varies according to the conditions of each industry. Industries and occupations are generally divided into process, production, and services (14). In each of these industries, there are jobs in which the sleep status of employees is of grave importance. If sleep problems occur for employ-

ees of these occupations, their productivity will be significantly reduced. Process industries are the first category to be discussed.

Process industries: Process industries are considered to be complex systems due to the presence of hazardous materials, the large number of workers, and the complexity of operational processes (82). These industries use chemical, biochemical, or physical materials to convert raw materials and/or energy. These industries include various industries such as chemical and petrochemical, gas processing, electricity generation, and water and wastewater industries (83). The topic of sleep is essential in many industries in this category. One of these industries is the petroleum industry. In this industry, due to the nature of the work (workers work 24 hours a day, 7 days a week), the sleep status of employees is severely affected (80). The decline in sleep quality is evident in workers in the petroleum industry. A cross-sectional study on 43 male sailors from the National Iranian Drilling Company found poor sleep quality among oil rig shift workers (84).

The gas industry is also one of the most vital industries in this category. The working conditions in the gas industry are also challenging and dangerous (85). Due to the shift work and the nature of employees' work in the gas industry, the quality of sleep of workers dramatically decreases (86). Due to poor working conditions, most gas refinery workers complain of insufficient sleep because insufficient sleep negatively affects cognitive behaviours such as logical reasoning and working memory (31).

Manufacturing industries: These industries are defined as operational processes that have specific products. In these industries, identifiable objects are produced (14). The steel industry, automotive industry, cable industry, and many other manufacturing industries fall into this category. Numerous studies have evaluated sleep quality and its related disorders among manufacturing workers. A cross-sectional study conducted on 256 steel plant employees in Iran showed a significant relationship between general health, shift work, and sleep disorders (36). In addition, another cross-sectional study was conducted with the participation of 2820 manufacturing workers in China. According to the findings of this study, both insomnia and sleep duration in workers independently and jointly are associated with an increased risk of poor working ability (87).

Service industries: In these industries, fewer industrial resources are usually used. These industries are a large group of occupations working to serve other industries, especially manufacturing industries (14). Driving is one of these occupations. Driving is a high-precision job that requires psychomotor responses and is highly susceptible to fatigue (88). Fatigue and drowsiness are particularly common among truck drivers due to the nature of their job (89).

Inadequate sleep and drowsiness are the leading causes of accidents in these occupations (90). A cross-sectional study conducted on 150 truck drivers in Iran found that about a quarter of road accidents are caused by drowsiness, and driving accidents are less common among drivers who get enough sleep (91).

Firefighting is another occupation in this category. Firefighters are primarily responsible for public safety, and their services are full-time (92). Firefighters often experience irregular and inadequate sleep due to their shift work time (93). Because of their frequent night shift schedules and workplaces, they are at greater risk of sleep deprivation than the general population (94). Sleep disorders in firefighters are significantly more common than in the public adult community (95).

Nursing is another job that falls into this category. Nursing is a distinct occupational group with a different job nature than other occupations (11). Nursing requires focus and often involves activities that require a high level of complexity and responsibility (96). Decreased sleep quality has been shown to impair nurses' decision-making and performance, resulting in errors and accidents (97). Low-quality sleep in this profession increases absenteeism, decreases job satisfaction, and reduces the quality of tasks (98). A cross-sectional study of 315 shift nurses in Italy found that specific characteristics and employment status of shift nurses can indirectly affect sleep quality and burnout (99).

In general, inadequate sleep quality and its related disorders are interrelated with many work environment parameters. Many components of the work environment, such as shift work, job stress, working time, and rest, affect employee sleep. These harmful components of the work environment have affected the quality of sleep of peoples in various industries and created unfavorable conditions for employees' sleep (15, 16). This unhealthy sleep situation has long-term adverse effects on employees in multiple industries.

Conclusion

Employees' sleep status in various industries is unsatisfactory and can be affected by many factors such as job stress, anxiety, stress, and shift work. Poor sleep status can negatively affect employee performance, productivity, accident rates, and many other factors. By taking the necessary measures and through appropriate planning according to employees' quality of sleep in the workplace, poor sleep quality and the resulting sleep disorders can to some extent be prevented.

Among the limitations of most studies was the use of subjective tools to assess the quality of sleep and its mental disorders such as questionnaires. In this regard, researchers are advised to conduct more clinical studies using modern equipment if possible. Moreover, due to the small number of studies in this regard, it is recommended that more researches be conducted on sleep quality and its related disorders in the manufacturing industries.

Conflict of Interests

Authors have no conflict of interests.

Acknowledgments

The authors would like to thank the individuals who helped interpret the findings of some of the studies.

References

1. Zare R, Choobineh A, Keshavarzi S. Association of amplitude and stability of circadian rhythm, sleep quality, and occupational stress with sickness absence among a Gas Company Employees-a cross sectional study from Iran. *Saf Health Work* 2017; 8: 276-81.
2. Khorasani M, Mohammadpoorasl A, Javadi M. The association between sleep quality and metabolic factors and anthropometric measurements. *Biotech Health Sci* 2016; 3: 25-31.
3. Rothstein NH. High-performing workforces demand high-quality sleep. *Target Logistics White Paper*. Target Logistics Management LLC 2013; 10: 1-14.
4. Velasco GM, Mette J, Mache S, et al. Sleep quality of offshore wind farm workers in the German exclusive economic zone: A cross-sectional study. *BMJ Open* 2018; 8: e024006.
5. Lee S, Jackson CL, Robbins R, Buxton OM. Sleep health and the workplace. *Sleep and health*. San Diego, CA: Elsevier Academic Press; 2019. p. 457-71.
6. Nasr Esfahani M, Haghjoo A, Danesh H, et al. Assessment of the relationship between sleep disorder and

- the ability to do work in workers in one of the oil-rich southern regions. *Occupational Medicine* 2017; 9: 14-21. [In Persian].
7. Nodoushan R, Halvani G, Nadjarzadeh A. Relation between road accidents and sleep quality of heavy vehicle drivers in Yazd. *Int J Env Health Eng* 2012; 1: 40.
 8. Yun JA, Ahn YS, Jeong KS, et al. The relationship between chronotype and sleep quality in korean firefighters. *Clin Psychopharmacol Neurosci* 2015; 13: 201-8.
 9. Lim DK, Baek KO, Chung IS, et al. Factors related to sleep disorders among male firefighters. *Ann Occup Environ Med* 2014; 26: 11.
 10. Weaver MD, Vetter C, Rajaratnam SMW, et al. Sleep disorders, depression and anxiety are associated with adverse safety outcomes in healthcare workers: A prospective cohort study. *J Sleep Res* 2018; 27: e12722.
 11. Han Y, Yuan Y, Zhang L, et al. Sleep disorder status of nurses in general hospitals and its influencing factors. *Psychiatr Danub* 2016; 28: 176-83.
 12. Gaultney JF, Collins-McNeil J. Lack of sleep in the workplace: What the psychologist-manager should know about sleep. *Psychol Manag J* 2009; 12: 132-48.
 13. Labbafinejad Y, Ghaffari M, Bahadori B, et al. The effect of sleep disorder on the work ability of workers in a car accessories manufacturing plant. *Med J Islam Repub Iran* 2014; 28: 727-34.
 14. Pont JPD, Debacq M. *Process industries 1: Sustainability, managerial and scientific fundamentals*. Hoboken, NJ: Wiley; 2020.
 15. Palo S, Das M. Prevalence of sleep problems in diverse professions in india and its HR implications. *Sleep Vigilance* 2019; 3: 65-75.
 16. Ozvurmaz S, Mandiracioglu A. Work-related fatigue and related factors among nurses working at the Adnan Menderes University Hospital. *Medical Science and Discovery* 2018; 5: 229-34.
 17. Kazemi R, Haidarimoghadam R, Motamedzadeh M, et al. Effects of shift work on cognitive performance, sleep quality, and sleepiness among petrochemical control room operators. *J Circadian Rhythms* 2016; 14: 1.
 18. Munn Z, Barker TH, Moola S, et al. Methodological quality of case series studies: An introduction to the JBI critical appraisal tool. *JBI Evid Synth* 2020; 18: 2127-33.
 19. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015; 4: 1.
 20. Yazdi Z, Abbasi M, Nabatian M, et al. Comparison of sleep disturbances in shift workers and people working with a fixed shift. *Iran South Med J* 2013; 16: 320-30. [In Persian].
 21. Yazdaniraad S, Khoshakhlagh A, Fallah H, et al. Association between sleep disorders and occupational accidents. *Iran Occup Health* 2017; 1: 35-41. [In Persian].
 22. Sortoglu B, Kurt r, Ozyilmaz S, et al. Influence of the duration of working in a textile factory on respiratory functions and sleep quality. *Eur Respir J* 2018; 52: A4517.
 23. Sadeghniaat-Haghighi K, Mehrabinejad MM, Hajighaderi A, et al. Shift work disorder, insomnia, and depression among offshore oil rig workers. *Iran J Psychiatry* 2021; 16: 162-7.
 24. Ning L, Shi L, Tao N, et al. Effects of occupational stress and circadian clock gene polymorphism on sleep quality of oil workers in Xinjiang, China. *Med Sci Monit* 2020; 26: e924202.
 25. Monazzam MR, Hosseini M, Matin LF, et al. Sleep quality and general health status of employees exposed to extremely low frequency magnetic fields in a petrochemical complex. *J Environ Health Sci Eng* 2014; 12: 78.
 26. Moghadam Dizaj Herik BN, Asle Fattahy B, Taghizadeh H. Effectiveness of stress management training on mental health, job satisfaction and quality of sleep of day work and shift work employees of Tabriz Petrochemical Company. *The Journal of Productivity Management* 2013; 7: 145-64. [In Persian].
 27. Li X, Gao X, Liu J. Cross-sectional survey on the relationship between occupational stress, hormone levels, and the sleep quality of oilfield workers in Xinjiang, China. *Int J Environ Res Public Health* 2019; 16: 28.
 28. Legault G, Clement A, Kenny GP, et al. Cognitive consequences of sleep deprivation, shiftwork, and heat exposure for underground miners. *Appl Ergon* 2017; 58: 144-50.
 29. Lavigne AA, Hebert M, Auclair J, et al. Good sleep quality and progressive increments in vigilance during extended night shifts: A 14-day actigraphic study in underground miners. *J Occup Environ Med* 2020; 62: e754-e759.
 30. Kiani F, Safavi Mirmahale SH, Khodabakhsh MR. Role of work overload and sleep deprivation in predicting work situation awareness among workers of a petrochemical industry. *J Research Health* 2016; 6: 487-94.
 31. Khosrovanmehr N, Mokhtari L, Rajezi Esfahani S, et al. A comparative study of mental health and sleep problems in shift workers and day workers of one gas refinery in Iran. *J Sleep Sci* 2018; 2: 40-5.
 32. Joaquim AC, Lopes M, Stangherlin L, et al. Mental health in underground coal miners. *Arch Environ Occup Health* 2018; 73: 334-43.
 33. Choobineh A, Javadpour F, Azmoon H, et al. The prevalence of fatigue, sleepiness, and sleep disorders among petrochemical employees in Iran. *Fatigue* 2018; 6: 153-62.
 34. Bagheri HM, Khanjani N, Ebrahimi MH, et al. The effect of chronic exposure to extremely low-frequency electromagnetic fields on sleep quality, stress, depression and anxiety. *Electromagn Biol Med* 2019; 38: 96-101.
 35. Asadi N, Hosseini SS. Study of job stress and its association with sleep quality among employees of Behbahan Bid Boland Gas Refinery in 2017: A descriptive study. *J Rafsanjan Univ Med Sci* 2019; 18: 797-810. [In Persian].
 36. Hashemi z, Alidosti M. Association relationship between shift work and sleep quality with public health in personnel Kaavian steel companies Khozestan. *Pa-joohande* 2016; 21: 161-6. [In Persian].

37. Abbasi M, Monnazzam MR, Zakerian S, et al. Effect of wind turbine noise on workers' sleep disorder: A case study of Manjil Wind Farm in Northern Iran. *Fluct Noise Lett* 2015; 14: 155020.
38. Zamanian Z, Nikeghbal K, Khajehnasiri F. Influence of sleep on quality of life among hospital nurses. *Electron Physician* 2016; 8: 1811-6.
39. Wolkow AP, Barger LK, O'Brien CS, et al. Associations between sleep disturbances, mental health outcomes and burnout in firefighters, and the mediating role of sleep during overnight work: A cross-sectional study. *J Sleep Res* 2019; 28: e12869.
40. Vincent GE, Aisbett B, Hall SJ, et al. Fighting fire and fatigue: sleep quantity and quality during multi-day wildfire suppression. *Ergonomics* 2016; 59: 932-40.
41. Stout JW, Beidel DC, Brush D, et al. Sleep disturbance and cognitive functioning among firefighters. *J Health Psychol* 2021; 26: 2248-59.
42. Smith LJ, Bartlett BA, Tran JK, et al. Sleep disturbance among firefighters: understanding associations with alcohol use and distress tolerance. *Cogn Ther Res* 2019; 43: 66-77.
43. Shao MF, Chou YC, Yeh MY, et al. Sleep quality and quality of life in female shift-working nurses. *J Adv Nurs* 2010; 66: 1565-72.
44. Saraei M, Najafi A, Heidarbagi E. Risk factors for obstructive sleep apnea among train drivers. *Work* 2020; 65: 121-5.
45. Saleh AM, Awadalla NJ, El-masri YM, et al. Impacts of nurses' circadian rhythm sleep disorders, fatigue, and depression on medication administration errors. *Egypt J Chest Dis Tuberc* 2014; 63: 145-53.
46. Sadeghniaat-Haghighi K, Yazdi Z, Kazemifar AM. Sleep quality in long haul truck drivers: A study on Iranian national data. *Chin J Traumatol* 2016; 19: 225-8.
47. da Rocha MC, De Martino MM. Stress and sleep quality of nurses working different hospital shifts. *Rev Esc Enferm USP* 2010; 44: 280-6. [In Portuguese].
48. Rajaratnam SM, Barger LK, Lockley SW, et al. Sleep disorders, health, and safety in police officers. *JAMA* 2011; 306: 2567-78.
49. Park E, Lee HY, Park CS. Association between sleep quality and nurse productivity among Korean clinical nurses. *J Nurs Manag* 2018; 26: 1051-8.
50. Najafi Ghezeljeh T, Moradi F, Rafii F, et al. Relationship between job stress, sleep quality and fatigue in nurses. *Iran J Nurs* 2014; 27: 40-9. [In Persian].
51. Ma CC, Hartley TA, Sarkisian K, et al. Influence of work characteristics on the association between police stress and sleep quality. *Saf Health Work* 2019; 10: 30-8.
52. Slaven JE, Mnatsakanova A, Burchfiel CM, et al. Association of sleep quality with depression in police officers. *Int J Emerg Ment Health* 2011; 13: 267-77.
53. Lees T, Chalmers T, Burton D, et al. Electroencephalography as a predictor of self-report fatigue/sleepiness during monotonous driving in train drivers. *Physiol Meas* 2018; 39: 105012.
54. Lammers-van der Holst H, Kerkhof GA. Shift work tolerance and the importance of sleep quality: a study of police officers. *Biological Rhythm Research* 2015; 46: 257-64.
55. Jeon HJ, Kim JH, Kim BN, et al. Sleep quality, posttraumatic stress, depression, and human errors in train drivers: A population-based nationwide study in South Korea. *Sleep* 2014; 37: 1969-75.
56. Huss A, Ozdemir E, Schaap K, et al. Occupational exposure to MRI-related magnetic stray fields and sleep quality among. *Int J Hyg Environ Health* 2021; 231: 113636.
57. Ghaljaei F, Naderifar M, Ghaljeh M. comparison of general health status and sleep quality between nurses with fixed working shifts and nurses with rotating working shifts. *Zahedan J Res Med Sci* 2011; 13: 48-50. [In Persian].
58. Geiger-Brown J, Rogers VE, Han K, et al. Occupational screening for sleep disorders in 12-h shift nurses using the Berlin Questionnaire. *Sleep Breath* 2013; 17: 381-8.
59. Fedekulegn D, Burchfiel CM, Charles LE, et al. Shift work and sleep quality among urban police officers: The BCOPS Study. *J Occup Environ Med* 2016; 58: e66-e71.
60. Emkani M, Khanjani N. Sleep quality and its related factors in intercity bus drivers. *J Mil Med* 2012; 14: 137-41. [In Persian].
61. Elliott JL, Lal S. Blood pressure, sleep quality and fatigue in shift working police officers: Effects of a twelve hour roster system on cardiovascular and sleep health. *Int J Environ Res Public Health* 2016; 13: 172.
62. Deng X, Liu X, Fang R. Evaluation of the correlation between job stress and sleep quality in community nurses. *Medicine (Baltimore)* 2020; 99: e18822.
63. Chien PL, Su HF, Hsieh PC, et al. Sleep quality among female hospital staff nurses. *Sleep Disord* 2013; 2013: 283490.
64. Chang JH, Huang PT, Lin YK, et al. Association between sleep duration and sleep quality, and metabolic syndrome in Taiwanese police officers. *Int J Occup Med Environ Health* 2015; 28: 1011-23.
65. Chan MF. Factors associated with perceived sleep quality of nurses working on rotating shifts. *J Clin Nurs* 2009; 18: 285-93.
66. Chaiard J, Deeluea J, Suksatit B, et al. Factors associated with sleep quality of Thai intercity bus drivers. *Ind Health* 2019; 57: 596-603.
67. Carey MG, Al-Zaiti SS, Dean GE, et al. Sleep problems, depression, substance use, social bonding, and quality of life in professional firefighters. *J Occup Environ Med* 2011; 53: 928-33.
68. Braeckman L, Verpraet R, Van Risseghem M, et al. Prevalence and correlates of poor sleep quality and daytime sleepiness in Belgian truck drivers. *Chronobiol Int* 2011; 28: 126-34.
69. Billings J, Focht W. Firefighter shift schedules affect sleep quality. *J Occup Environ Med* 2016; 58: 294-8.
70. Bazrafshan MR, Moravveji F, Soleymaninejad N, et al. Prevalence and effects of sleep disorders among shift-working nurse. *Ann Trop Med Public Health*

- 2018; 11: 13-8.
71. Barger LK, Rajaratnam SM, Wang W, et al. Common sleep disorders increase risk of motor vehicle crashes and adverse health outcomes in firefighters. *J Clin Sleep Med* 2015; 11: 233-40.
72. Abbasi M, Rajabi M, Yazdi Z, et al. Factors affecting sleep quality in firefighters. *Sleep Hypn* 2018; 20: 283-9.
73. Jung YS, Chae CH, Kim YO, et al. The relationship between serum vitamin D levels and sleep quality in fixed day indoor field workers in the electronics manufacturing industry in Korea. *Ann Occup Environ Med* 2017; 29: 25.
74. Rasdi I. The association between noise, work stress and coping with sleep quality among cable manufacturing workers. *Asian J Agri Biol* 2018; 108-11.
75. Farvaresh E, Monazam MR, Abbassinia M, et al. Investigation the relationship between sleepiness and general health of shift workers in the automobile industry. *J North Khorasan Univ Med Sci* 2012; 4: 221-7. [In Persian].
76. Chou WM, Hsieh L. The impact of Shift Work Implementation on sleeping quality and job performance: A case study of semi-conductor manufacturing company. Proceedings of the 40th International Conference on Computers and Industrial Engineering; 2010 Jul 25-28; Awaji City, Japan.
77. Mousavi SM, Sharifiniya S, Yazdani Rad S, et al. The relationship between shift work and its effects on the health of the operational staff in Abadan Oil Refining Company. *J Prevent Med* 2017; 4: 19-25. [In Persian].
78. McDowall K, Murphy E, Anderson K. The impact of shift work on sleep quality among nurses. *Occup Med (Lond)* 2017; 67: 621-5.
79. Shabani Z, Gholipour M, Mirzaei M, et al. Investigating shift work disorder among workers of Hamadan Glass Factory in 2017. *Occupational Medicine* 2021; 12: 24-34. [In Persian].
80. Zhou Z, Zhang Y, Sang L, et al. Association between sleep disorders and shift work: a prospective cohort study in the Chinese petroleum industry. *Int J Occup Saf Ergon* 2022; 28: 751-7.
81. Khalatbari J, Ghorbanshiroudi S, Firouzbakhsh M. Correlation of job stress, job satisfaction, job motivation and burnout and feeling stress. *Procedia Soc Behav Sci* 2013; 84: 860-3.
82. eskandari T, Mirzaei M, mohammadfam I. Fire and explosion risk analysis using Bow-Tie Method and Fuzzy-Bayesian network in process industries. *Iran Occup Health* 2021; 18: 139-54. [In Persian].
83. Muller R, Oehm L. Process industries versus discrete processing: How system characteristics affect operator tasks. *Cogn Tech Work* 2019; 21: 337-56.
84. Sadeghniaat-Haghighi K, Zahabi A, Najafi A, et al. Evaluating the quality and duration of sleep using actigraphy in petroleum industry shift workers. *Sleep Health* 2020; 6: 407-10.
85. Pandey S, Pestonjee D. A qualitative study of work stress and employees expectations of stress management interventions in context of Indian oil and gas industry professionals. Proceedings of the 2017 International Research Conference in Asia- AHRD; 2017 Nov 8-10; Ahmedabad, India.
86. Sadeghniaat-Haghighi K, Aminian O, Najafi A, et al. Sleep quality in shift workers of offshore petroleum industries. *J Sleep Sci.* 2019; 3: 36-40.
87. Lian Y, Xiao J, Liu Y, et al. Associations between insomnia, sleep duration and poor work ability. *J Psychosom Res* 2015; 78: 45-51.
88. Okrepilov V, Kuzmina S, Kuznetsov S. Tools of quality economics: Sustainable development of a 'smart city' conditions of digital transformation of the economy. *IOP Conf Ser: Mater Sci Eng* 2019; 497: 012134.
89. Phatrabuddha N, Yingratanasuk T, Rotwannasin P, et al. Assessment of sleep deprivation and fatigue among chemical transportation drivers in Chonburi, Thailand. *Safety and Health at Work* 2018; 9: 159-63.
90. Kwon S, Kim H, Kim GS, et al. Fatigue and poor sleep are associated with driving risk among Korean occupational drivers. *J Transp Health* 2019; 14: 100572.
91. Malek M, Halvani GH, Fallah H. A study of the relationship between the Pittsburgh sleep quality index and road accidents among truck drivers. *Occupational Medicine* 2011; 10: 14-20.
92. Jang TW, Jeong KS, Ahn YS, et al. The relationship between the pattern of shift work and sleep disturbances in Korean firefighters. *Int Arch Occup Environ Health* 2020; 93: 391-8.
93. Kim HW, Jung SM, Choi YS, et al. Sleep patterns of firefighters with shift working schedules in Seoul Metropolitan Area. *Sleep Med Res* 2017; 8: 68-75.
94. Shi Y, Bender B, McGovern P, et al. Daytime sleepiness among Midwestern firefighters. *Arch Environ Occup Health* 2021; 76: 433-40.
95. Lin SH, Liao WC, Chen MY, et al. The impact of shift work on nurses' job stress, sleep quality and self-perceived health status. *J Nurs Manag* 2014; 22: 604-12.
96. Palhares VC, Corrente JE, Matsubara BB. Association between sleep quality and quality of life in nursing professionals working rotating shifts. *Rev Saude Publica* 2014; 48: 594-601.
97. Zhang L, Sun DM, Li CB, et al. Influencing factors for sleep quality among shift-working nurses: a cross-sectional study in china using 3-factor Pittsburgh Sleep Quality Index. *Asian Nurs Res (Korean Soc Nurs Sci)* 2016; 10: 277-82.
98. Babamiri M, Moeini B, Tahmasian H, et al. The study of sleep health education effect on sleep quality among Lorestan nursing personnel. *Iran J Ergon* 2017; 4: 8-13. [In Persian].
99. Giorgi F, Mattei A, Notarnicola I, et al. Can sleep quality and burnout affect the job performance of shift-work nurses? A hospital cross-sectional study. *J Adv Nurs* 2018; 74: 698-708.