

## The Use of Nested PCR Method for Rapid Detection of *Rosellinia necatrix* Causal Agent of White Root Disease From Soil and Root

Mohammadi Meymand<sup>1</sup>, S. M., Sharifnabi<sup>1\*</sup>, B. and Bahar<sup>1</sup>, M.

### Abstract

White root rot disease, caused by *Rosellinia necatrix* is one of the most important disease of fruit trees in Iran and worldwide. To detect the pathogen from soil and root, six infected soil samples were collected from cherry crown trees and two samples were collected as positive and negative control. DNA Extraction from eight soil samples including infected cherry roots by white cottony mycelium, roots infected by white mycelial fans and symptom less roots and two faba roots, with white cottony mycelium and symptom less root was performed. Detection of the Pathogen was done by two specific primer pairs R2 .R8 and R7 .R10 in Nested PCR reaction. The results showed the detection from six infected soil, positive control and in roots colonized by white cottony mycelium and roots by white mycelial fans and faba roots colonized by white cottony mycelium. No detection was achieved from symptom less roots. To verify the accuracy of the used detection method, ITS region was amplified and cloned in pTZ57R/T vector and sequenced. Comparison of the sequence showed %98.99 similarities to the recognized isolate of *R. necatrix*. According to our findings, it seems that Nested PCR method could be useful to rapid detection of *R. necatrix* from soil and root of plants in orchards.

**Keywords:** *Rosellinia necatrix*, Nested PCR, White root rot, Detection

### References

- BEHDAD, E. 1975a. Isolation and cultivation of *Rosellinia necatrix* (Hart.) Berl., *in vitro* and its identification. Iran journal of Plant Pathology. 11: 122-137.
- BEHDAD, E. 1975b. Morphology, distribution, importance and hosts of *Rosellinia necatrix* (Hart.) Berl., the cause of white root rot in Iran. Iran journal of Plant Pathology. 11: 30-71
- BEHDAD, E. 1976. Influence of soil amendments, water management and tree species on the growth of *Rosellinia necatrix* (Hartig) Berl., the cause of white root rot. Iran journal of Plant Pathology. 12: 27-41
- CHENARD, V. 2003. Evaluation of methods of soil DNA extraction for PCR. MSc. thesis submitted to Alberta University. 102 pp.
- CULLEN, D. W. AND HIRSCH, P.R. 1998. Simple and rapid method for direct extraction of microbial DNA from soil for PCR. Soil Biology and Biochemistry 30: 983–993.
- FREEMAN, S. AND SZTEJNBERG, A. 1992. *Rosellinia*. PP. 71-73. In: L. L Singleton, J.D. Mihail and C. M. Rush (Eds), Method for research on soil borne phytopathogenic fungi. APS press, USA.
- RUANO-ROSA, D., SCHENA, L., IPPOLITO, A. AND LOPEZ-HERRERA, C.J 2007. Comparison of conventional and molecular method for the detection of *Rosellinia necatrix* in avocado orchards in southern Spain. journal of Plant Patholog. 56: 251-256.
- SCHENA, L. AND IPPOLITO, A. 2003. Rapid and sensitive detection of *Rosellinia necatrix* roots and soils by real time scorpion-PCR. Journal of Plant Pathology 85: 15-25.
- SCHENA, L., NIGRO, F. AND IPPOLITO, A. 2002. Identification and detection of *Rosellinia necatrix* by conventional and real time Scorpion – PCR. European Journal of Plant Pathology 108: 355-366.
- SCHENA, L., NIGRO, F. AND IPPOLITO, A. 2008. Integrated management of *Rosellinia necatrix* root rot on fruit tree crops PP. 137-158 In: Ciancio., A. and Mukerji., K.G. (Eds.), Integrated management of disease caused by Fungi, Phytoplasma and Bacteria. Springer Science.
- WILSON, I. G. 1997. Inhibition and facilitation of nucleic acid amplification. Applied and Environmental Microbiology 63: 3741-3751.

To look at the figures and tables, please refer to the Persian text (pages: 9-15= ۹-۱۵).

1. Former MSc. Student and Associate Professors of Plant Pathology, Department. of Plant Protection, College of Agriculture, Isfahan University of Technology

\*: Corresponding author