Investigation on Interaction Effect of Benzyladenine and Chitosan on *in vitro* Proliferation of Strawberry (*Fragaria* ´ *ananassa* cv. Selva)

Jalili Marandi¹*, R., Naseri¹, L., Mohseniazer¹, M., Hajitagiloo¹, R. and Marhamati¹, M. R.

Abstract

Micropropagation of strawberry provides possibility of fast multiplication. Growth regulators can be effective on micropropagation of strawberry and fast multiplication. To attention to the importance of proliferation stage of micropropagation, in this research an investigation carried out on the interaction effect of benzyladenine (BA) and chitosan. The Murashige and Skoog (MS) culture medium was supplemented with BA in two concentration (1 and 2 mg/l) and Chitosan with high molecular weight in five concentration (0,20,30,40 and 60 mg/l). A factorial experiment based on completely de +sign with eight replication was used. Evaluated characteristics included number of shoots, diameter of shoots, length of shoots, number of leaves, size of leaves, chlorophyll content and dry weight of plant mass. Based on the obtained results there was not significant differences between 1 or 2 mg/l concentration of BA from the point of effect on examined characteristics. The highest number of shoots, diameter of shoots, number of leaves and dry weight of plant mass were obtained in 1 or 2 mg/l BA and without chitosan. But the highest length of shoot and chlorophyll content were observed in combination of 1 or 2 mg/l BA with30 mg/l chitosan. The highest leaf size were obtained in combination of 1 or 2 mg/l BA with30 mg/l chitosan. According to the results, 1mg/l BA was significant effective on proliferation of Selva strawberry.

Keywords: Proliferation, Strawberry, Benzyladenine, Chitosan, in vitro

References

- Abramenko, N. M. 1983. A rapid method of strawberry propagation by meristem culture in vitro. Horticulturae, 53:209-212.
- Ait Barka, E., Eullaffroy, P., Clement, C. and Vernet, G. 2004. Chitosan improves development and protect- *Vitis vinifera* L. Against *Botrytis cinerea*. Plant Cell Report, 22:608-614.
- Arora, D. K., Suri,S. S. and Ramawat, K. G. 2006. Assessment of variability in the regenerants from long- term cultures of sfed musli (*Chlorophytum borivilianum*). Indian Journal of Biotechnology, 5:527-534.
- Asghari- Zakaria, R., Maleki-Zanjani. B. and Sedghi, E. 2009. Effect of in vitro chitosan application on growth and minituber yield of *Solanum tuberosum* L. Plant Soil Environ, 55: 252-254.
- Bhatt, I. D., and Dhar, U. 2000. Micropropagation of Indian wild strawberry. Plant Cell. Tissue and Organ Culture, 60:83-88.
- Biswas, M. K., Hossain, M. and Islam, R. 2007. Virus free plantlets production of strawberry through meristem culture. World Journal of Agricultureal Science, 3:757-763.
- Carelli, B. P., and Echeverrigary, S. 2002. An improved system for the *in vitro* propagation of rose cultivars. Scientia Horticulturae, 92:64-74.
- Chien, P. J., Sheu, F., Huang, W. T. and Su, M. S. 2007. Effect of molecular weight of chitosans on their antioxidative activities in apple juice . Food Chemistry 102 :1192–1198.
- Freepons, D. 1991. Chitosan, does it have a place in agriculture? Proceeding of Plant Growthe Regulation Society of America, p. 11-19.
- Jalili Marandi, R. 2007. Small Fruits. Jihad-e-Daneshgahi. Urmia.297 pp. (In Farsi).
- Khosh-Khui, M. and Sink, K.C. 1982. Rooting enhancement of *Rosa hybrida* for tissue culture propagation. Science Horticulture, 17:371-376.
- Khosh-Khui, M. 1994. (Translator). By Kenneth ,C. Torres. Tissue culture techniques .Shiraz University press.467 pp.
- Kume, T., Nagasawa, N. and Yoshii, F. 2002. Utilization of carbohydrates by radiation processing. Radiation Physics and Chemistry, 63.:625-627.
- Nge, K. L., New, N. Chandrkrachang, S. and Stevens, W.F. 2006. Chitosan as a growth stimultor in orchid tissue culture. Plant Science, 170:1185-1190.
- Owen, H. R., and Miller, A. R. 1996. Haploid plant regeneration from anther cultures of three north American cultivars of strawberry (*Fragaria* × *ananassa* Duch). Plant Cell Reports, 15:905-909.

^{1.} Associate professor, Assistant Professor, MS in Horticulture Respectivly, Department of Horticulture, Faculty of Agriculture, Urmia University, Urmia

^{*:} Corresponding author

AGRICULTURAL BIOTECHNOLOGY, Vol. 10, No. 1, 2011

- Park , S. Y. , Marsh , K. S. and Rhim , J. W. 2002. Characteristics of different molecularweight chitosan films affected by the type of organic solvents. Journal of Food Science, 67(1): 194-197.
- Passey, A. J., Barrett, K. J. and James, D. J. 2002. Adventitious shoot regeneration from seven commercial strawberry cultivars (*Fragaria* × *ananassa* Duch). Using range of explant types. Plant Cell Report, 21:397-401.
- Qiuping, Z., and Wenshui, X. 2007. Effect of 1-methylcyclopropene and/or chitosan coating treatments on storage life and quality maintenance of Indian jujube fruit. Science Direct, 40:404-411.
- Sakila, S., Ahmed, M. B., Roy, U. K. Biswas, M. K., karim, R., Razvy, M. A., Hossain, M., Islam, R. and Hoque, A. 2007. Micropropagation of strawberry (*Fragaria × ananassa* Duch) a newly introduced crop in Bangladesh. American-Eurasian Journal of Scientific Research, 2:151-154.
- Wikins, C. P., and Dodds, J. H. 1982. Effect of various growth regulators on growth *in Vitro* of cherry shoot tips. Plant Growth Regulation, 1:209-216.
- Yokimova, E., Kapchina-Toteva, V., Groshkoff, I. and Ivanova, G. 2000. Effect of BA CPPU on protease and amylase activity of *Rosa hybrida*. Bulgaria Journal of Plant Physiology, 26:39-47.
- Zhang, J., Ningtong, XU., Qihuan, QU. and Xiuhong, XU. 2009. Effect of composite of traditional Chinese medicine and chitosan seed coating agent on growth of soyban. Journal of Northeast Agricultural University, 1:202-207.

To look at the figures and tables, please refer to the Persian text (pages: 27-34=27-34).