The Simulation of Climate Change in Semnan Province with Scenarios of Atmospheric General Circulation Model (Hadcm3)

Zahra Hejazizadeh^a, Sayyed Mohammad Hosseini^b, Alireza Karbalaee Dorei^{c1}

^a Department of Physical Geography, Kharazmi University, Tehran, IRAN

^b Department of Geography, Sayyed Jamaleddin Asadabadi University, Hamadan, IRAN

^c M.A. in Climatology, Kharazmi University, Tehran, IRAN

Received 20 October 2014 Accepted 9 May 2015

1. Introduction

Climatic change is one of the most important challenges human beings have confronted with in recent centuries due to its severe effects on water resources, agriculture, energy, tourism and even human bioclimatic. Developing strategies, making decisions with awareness about water consumption in different parts in the future, and available water resources management require climatic change information (i.e. precipitation and temperature) which can be directly used with hydrology and climatology models.

In recent years, climate prediction models have been used for numerous purposes such as the generation of climate data using different scenarios related to greenhouse gases in general circulation models. By using these models, the output of general circulation models (GCM) can be examined at micro-scales. One of these models is called LARS-WG, which has received considerable attention from the researchers in Iran or elsewhere.

2. Case Study

Using the LAR model, we have collected climate data from among four synoptic stations from their establishment until 2010. Data used in this research including minimum and maximum temperature as well as precipitation were daily recorded. To generate the precipitation values, LAR (a well-known generator model of random data for climatic status) was used. Likewise, the minimum and maximum temperature in every station of Semnan Province was used for both basic and future periods..

In addition,, to investigate the LAR model in simulating, the Ground Truth average bias and absolute error have been calculated for the simulated data and truth in the investigated period:

$$Bias = \frac{1}{n} \sum_{i=1}^{n} (S_i - O_i)$$
$$MAE = \frac{1}{n} \sum_{i=1}^{n} |S_i - O_i|$$

1 Corresponding author: Alireza Karbalaee Dorei. Tel: 09132770511

E-mail: Karbalaee20@gmail.com

1

3. Conclusion

In these scenarios, there was almost full coverage between the Ground Truth and simulated data. In A1B scenario, more stations bring more assurance in simulating data in winter and spring. The maximum observed and simulated precipitation was calculated by Hadcm3 model scenario in all Semnan stations in March. The minimum precipitation in Shahrood and Biarjmand stations was in July and August, respectively. As for Garmsar and Semnan stations, the minimum precipitation was in September. All stations in January had the least minimum and maximum temperature in January, whereas they had the most minimum and maximum temperature in July.

Keywords: Downscaling, Atmospheric general circulation model, Climate change, LARS-WG, Semnan Province.

References (in Persian)

- 1. Ababaei, B., Sohrabi, T., Mirzaei, F., Rezavrdynejad, V., & Karimi, B. (2011). The effects of climate change on wheat yield and risk analysis associated with it (Case study: Isfahan Ruddasht area). *Water and Soil Science*, *3*, 135-150.
- 2. Abbasi, F., Malbousi, S., Babaeian, I., Asmari, M., & Borhani, R. (2010). Climate change prediction of South Khorasan province during 2010–2039 by using statistical downscaling of ECHO-G data. *Journal of Soil and Water*, *24*(2), 218-233.
- 3. Alizadeh, A., Salehnia, N., Sayyaree, N., & Nasiri, M. (2014). Forecasting possible changes in the climatic zones of effective rainfall for wheat (during the next 90 years). *Irrigation and Drainage*, *4*, 575-584.
- 4. Ashraf, B., Baygi Mousavi, M., Kamali, M. Davary, G. A., & Davary, K. (2013). Change of wheat and sugar beet due to the effects of climate change in the next two decades in the plains of Khorasan. *Journal of Irrigation and Drainage*, *2*, 105-117.
- 5. Babaeian, I., & Kohi, M. (2013). Analysis of agricultural climate scenarios of climate change in selected stations in Khorasan Razavi. *Journal of Soil and Water*, 26(4) 953-967.
- 6. Babaeian, I., & Najafi, Z. (2011). The analysis of climate change in Khorasan Razavi during 2010-2039 by using downscaling of GCM model output. *Journal of Geography and Regional Development*, 15, 1-19.
- Babaeian, I., Najafi, Z., Zabul Abbasi, F., Nokhandan Habibi, M., & Malbousi, S. (2010). Assessment of climate change during 2010 and 2039 AD, using downscaling data of atmospheric circulation model ECHO-G. *Geography and Development*, 16, 135-152.
- Haghtalab, N., Goudarzi, M., Habibi Nokhandan, M., Yavari, A. R., & Jafari, H. R. (2014). Tehran and Mazandaran climate modeling using the climate model of LARS-WG and comparing the changes in the front North and South of Central Alborz. *Journal of Environmental Science & Technology*, 1, 37-49.
- Khazanehdaree, L., Zabul Abbasi, F., Ghandhari, S., Kohi, M., & Malbousi, Sh. (2010). The prospect of drought situation in Iran during the next thirty years. *The Journal of Geography and Regional Development*, 12, 83-98.

2

- 10. Khorani, A., & Marvdashti, S. H. (2015). The effect of climate change on visiting in Hengam Island. *Journal of Physical Geography*, *1*, 109-122.
- Kohi, M., Malbousi, S. H., & Sanaei Nejad, H. (2014). A study of climate change scenarios based on the results of the two methods, downscaling for factors of reference evapotranspiration in Orumiyeh. *Journal of Irrigation and Drainage*, 7(4), 559-574.
- 12. Lashkari, A., Alizadeh, A., & Bannayan, M. (2012). Studying the possibility of reducing the effect of climatic factors on corn production in the North East of Iran. *Journal of Soil and Water*, 25(4), 926-939.
- 13. Roshan, G. R., & Ghanghormeh, A. B. (2015). Effects of climate change on intensity and frequency precipitation of North West stations in Iran. *Journal of Geography and Environmental Planning*, *2*, 61-84.
- 14. Roustaee, M., Sohrabi, T., Massahbavani, A., & Ahadi, M. S. (2013). Analysis of Risk Corn Biomass affected by climate change. *Journal of Agricultural Water*, *4*, 425-438.
- 15. Salehnia, N., Mousavi Baygi, M., & Ansari, H. (2014). Rough forecasting using PDE index using LARS-WG models and HadCM3 (Case study: Nishapur basin). *Journal of Irrigation and Drainage*, 7, 93-103.
- Shahkouhi, E., & Roshan, G. R. (2014). Degree day soybean on climate change for decades (Case study: Gorgan). *Quarterly Journal of Geographical Space Planning*, 7, 51-66.
- 17. Sobhani, B., & Fatemi Nia, F. S. (2015). Modeling of climatic parameters in Southern Khorasan. *Natural Geography Journal*, *3*, 311-332.
- 18. Zarghami, M., Babaeian, I., Can'ani, Y., & Can'ani, R. (2013). A study of climate change and its effects on land (Case study: East Azarbaijan). *Journal of Management Science and Engineering*, *18*, 61-80.

References (in English)

- 1. Alison, L. K., Richard, G. L., & Nicholas, S. R. (2004). RCM rainfall for UK flood frequency estimation: Climate change results. *Journal of Hydrology*, *318*(1), 163-172.
- 2. Chang, H., Knight, C. G., Staneva, M. P., & Kostov, D. (2002). Water resource impacts of climate change in Southwestern Bulgaria. *Geo Journal*, *57*(3), 159-168.
- 3. Elshamy, M. E., Wheater, H. S., Gedney, N., & Huntingford, C., (2005). Evaluation of the rainfall component of weather generator for climate change studies. *Journal of Hydrology*, *326*(1), 1-24.
- Gordon, C., Cooper, C., Senior, C. A., Banks, H., Gregory, J. M., Johns, ... & Wood, R. A. (2000). The simulation of SST, seas ice extents and ocean heat transports in a version of the Hadley center coupled model without flux adjustment. *Climate Dynamic*, 16(1-2), 147-168.
- 5. Harmel, R. D. (2002). Evaluating the adequacy of simulating maximum and minimum daily air temperature with the normal distribution. *Journal of Applied Meteorology*, 41(7), 744-753.

Geography and Environmental Hazards 15(2015)

- 6. Harmsen, E., Miller, N. L., Schlegel, N. J., & Gonzalez, J. E. (2009). Seasonal climate change impacts on evapotranspiration, precipitation deficit and cop yield in Puerto Rico. *Agricultural Water Management*, *96*(7), 1085-1095.
- Hashemi, M. Z., Shamseldin, A. Y., & Melville, B. W. (2011). Comparison of SDSM and LARS-WG for simulation and downscaling of extreme precipitation events in a watershed. *Stochastic Environmental Research and Risk Assessment*, 25(4), 475-484.
- 8. Mavromatis, T., & Hansen, J. W. (2001). Interannual variability characteristics and simulated crop response of four stochastic weather generators. *Agricultural and Forest Meteorology*, *109*(4), 283-296.
- 9. Rajabi, A. (2010, January). *Modeling of Kermanshah climate by means of downscaling model of Lars-wg*. Paper presented at the 2nd National Conference on Utilization Integrated Management of Water Resources, Kerman, Iran.
- Rosenberg, N. J., Brown, R. A., Izaurralde, R. C., & Thomson, A. M. (2003). Integrated assessment of Hadley Centre (hadcm2) climate change projections on agricultural productivity and irrigation water supply in the conterminous United States: I. Climate change scenarios and impacts on irrigation water supply simulated with the HUMUS model. *Agricultural and Forest Meteorology*, *117*(1), 73-96.
- Semenov, M. A., & Barrow, E. M. (2002). A stochastic weather generator for use in climate impact studies. User Manual: Hertfordshire, UK. Taulis, M. E., & Milke, M. W. (2005). Estimation of WSIM weather simulation parameters in arid climates. *Ecological Modeling*, 184(2), 177-191.