

# Zonation and Spatial Planning Based on Physical-Spatial Characteristics

## Case Study: City of Urmia

Ali Shamaei<sup>a1</sup>, Nazila Yaghoob Nejad ASL<sup>b</sup>, Khadijeh Saeidi<sup>c</sup>, Roghayeh Amiri<sup>d</sup>

<sup>A</sup> Department of Geography and Urban Planning, University of Kharazmi, Tehran, Tehran, IRAN.

<sup>b</sup> PhD Candidate in Physical Geography, University of Mohaghegh Ardabili, Ardabil, IRAN.

<sup>c</sup> M.S.in Geography and Urban Planning, University of Kharazmi, Tehran, Tehran, IRAN.

<sup>d</sup> M.S.in Geography and Urban Planning, University of Kharazmi, Tehran, Tehran, IRAN.

*Received 27 December 2015*

*Accepted 16 November 2016*

### 1. Introduction

Urmia is one of the important historical cities of Iran which in recent years has been associated with physical-spatial rapid expansion. One of the necessities of spatial planning of Urmia is controlling the rapid expansion of the city in marginal areas that are unbalanced fertile land and industrial and mineral soils are under construction and this problem has presented numerous hazards in the fields of economy and environment. Excessive physical development in the surrounding areas is now one of the major challenges in managing urban spatial development. Recognizing these challenges in identifying the pattern formation and urban structure is important in the context of place and passing of time. The application of this identification helps proper diagnosis and urban land use and appropriate physical development of the city.

Urmia, in West Azarbaijan Province, is one of the ancient cities of Iran. In terms of the geographical coordinate system, it is located in between 44 degree and 3 minute to 47 degree and 23 minute east longitude and between 35 degree and 58 minute to 39 degree and 46 minute north latitude, at an altitude of 1332 meters above sea level. The area is 10 thousand and 548 hectares. The city has been located in the vast and verdure plain in length of 70 km and a width of 30 km. Average of monthly minimum and maximum temperature is 6.4 and 17.5, respectively. The average of relative humidity is 60% and the average of monthly precipitation is 341 mm. Most of the existing deposits in the region are sedimentary deposits, yet highly dynamic structural changes such as orogenic and faulting may be created by dynamic metamorphosis in some parts of the region. More outcrop of sedimentary units of Urmia region is related to Mesozoic Era (Jurassic) and Quaternary (Quaternary).

## 2. Material and Methods

In this study, documents in the library were used for literature review. Next, using climate data and digital elevation model and field surveys, base maps were prepared for zonation spatial planning. Then, the indicators: elevation, slope, soil, land use, climate, and geomorphology, distance from the rural, distance from the city, distance from the roads and distance from the river were used. The selection of criteria was based on field surveys which were performed on the study area. After that, criteria were turned fuzzy using fuzzy membership function and fuzzy gamma 0.9 operator was provided for spatial planning zonation.

Fuzzy model for the first time was provided by Lotfi Zadeh (1965). Fuzzy theory concluded that by creating the membership function in a range of real numbers, new methods develop. Using fuzzy logic in analysis, geographical processes are described more realistically than by Boolean logic. Fuzzy analysis is performed using GIS software.

## 3. Results and Discussion

Urmia has had rapid and inharmonious growth in recent years and due to natural population growth, immigration, expansion of services, the granting of political – official identity to the city as the provincial capital, assignment by various government agencies in the city, country development programs, socio-economic and political changes and finally favorable physical conditions, It has never seen such a large body of demographic and physical changes. Urmia has had high population and area growth in 20 years (1986-2006) t. The present study makes clear that following the land reform in 1964, in the process of physical development from 1966 to 2011 in Urmia, the influx of migrant population in the city of Urmia intensified. Development of marginalization in north and northeast parts of the city, in the communication path to the Salmas town in the north-western edge of Urmia Lake has had its consequences. Since 1987 up to now the highest growth has been in urban areas mostly in the west, North West, east and southern parts of Urmia, which caused the annexation of the surrounding lands to city, settlement cooperation and development and the establishment of inappropriate settlements and marginalization. The issue of ownership had of rapid population and area growth. The growth was fragmented. Given that Urmia is at a distance of approximately 15 kilometers from the east of Lake Urmia, this issue has been an important obstacle for physical development. In addition, in the east, northeast and southeast of the city there are agricultural lands and horticultural products. City development in these areas has been in contrast with these lands and is causing hazards and limitations for these lands. Thus, the development of the city stopped in this direction and development trends in the west, north, south west and south of the city are spreading. These directions which have followed the development process are the acceptable level of suitability. Overall, based on the assessments done in this study, the west, north, south west and south areas of the city are appropriate for development and consist of high and very high zones. Actually, the city has a single central structure – radial. With regard to the recognition and implementation of aerial photographs and

maps available of physical development, the results are largely acceptable. Obviously, the increased standards and assessment in lesser extent will bring more precision. So, with proper selection of base maps and merge and their detailed analysis, fuzzy logic allows more accurate assessments and provides more accurate locations and can be used for planning urban development.

#### 4. Conclusion

The results showed that zones with high suitability for the growth and development of the city are located in the west and southwest of Urmia and also in the south and northeast, in a scattered manner. In coastal areas of Urmia Lake and its immediate zones we see alluvial terraces deposits and very low, low and moderate areas. These areas have low ability in terms of physical development. Moreover, areas with low suitability can be seen in the northeast to southeast in a scattered manner.

**Keywords:** Zonation of urban land, Spatial planning, Spatial indices, Spatial- physical development, City of Urmia.

#### References (in Persian)

- Abbas Zadegan, M., & Rostam Yazdi, B. (1999). Utilization of smart growth and organizing urban sprawl. *Journal of Technology of Education*, 3(1), 33–48.
- Amiri, M.J., Jalali, S.G.H., Salman Mahini, A.A., Hosseini, S.M., & Azari Dehkordi, F. (2009). Ecological land capability assessment for forestry in 2000 and 3000 watersheds in the north of Iran using GIS. *Journal of Environmental Studies*, 35(50), 33–44.
- Amiri, M.J., Salman Mahini, A.A., Jalali, S.G.H., Hosseini, S.M., & Azari Dehkordi, F. (2009). A comparison of maps overlay systemic method and Boolean-Fuzzy logic in the ecological capability evaluation of No. 33 and 34 watershed forests in northern Iran. *Journal of Environmental Sciences*, 7(2), 109–124.
- Babaie Kafaki, S. (2001). *Environmental evaluation for forest land classification by using of GIS (Case study in Kazem-rood Watershed, north of Iran)*(Unpublished doctoral dissertation). Islamic Azad University, Science and Research Branch, Tehran, Tehran, Iran.
- Bagher Zadeh, H.R., Bagher Zadeh, A., & Moein Rad, H. (2012). Qualitative evaluation of Neishabur plain lands for wheat (*Triticum aestivum*. L), corn (*Zea mays*. L) and cotton (*Gossypium herbaceum*. L) Using GIS. *Journal of Agroecology*, 4(1), 41–51.
- Bayat, B., Motakan, A.A., Rahmani, B., & Arabi, B. (2011). Comprehensive planning of land use and land use planning in urban watershed using GIS- Case study: Catchment of Mahidasht. *Journal of Management System*, 4(13), 119–135.
- Fallah Shmasi, S.R. (2004). *Economic evaluation of different applications in middle Keliber Chaei (North Arasbaran) using linear planning*(Unpublished doctoral dissertation). University of Tehran, Tehran, Iran.

- Faraj Zadeh, M., & Karami, T.A. (2004). Land use planning using remote sensing and Geography Information System (Case study: Khoram Abad). *Journal of Physical Geography Research Quarterly*, 36(47), 81–94.
- Farid, Y. (2007). *Geography and urban logy*. Tabriz: University of Tabriz Press.
- Ghadimi, M., Hosseini, M.S., Pour Ghasemi, H.R., & Moradi, H.R. (2010). Protective modeling of Manesht protected area and Qalarang using fuzzy logic. *Journal of Environmental Science*, 8(1), 85–105.
- Ghorbani, R., Pour Mohamadi, M.R., & Mahmoud Zadeh, H. (2013). Ecological Approach in land use chang modeling of Tabriz metropolitan using multi temporal satellite images, multi criteria analysis and Cellular Automata Markov Chain (1984-2038). *Journal of Urban Studies*, 2(8), 13–30.
- Iran Meteorological Organization.*
- Makhdoom, M. (1987). New method to analyze and data summary in land use planning process. *Iranian Journal of Natural Recourses*, 1(41), 68–78.
- Makhdoom, M. (1993). Development of the proposed method in 1987 to analyze and data summary in land use planning process. *Iranian Journal of Natural Recourses*, 3(46), 109–112.
- Makhdoom, M. (1997). Land use planning of woodsy six sub-basins Fars Province: Guideline for Zagros strategic planning. *Journal of Environmental Studies*, 19(19), 41–50.
- Makhdoom, M., Darvish Sefat, A.A., Jafar Zadeh, H., & Makhdoom, A. (2001). *Environmental assessment and planning using Geographical Information System*. Tehran: University of Tehran Press.
- Mir Katouli, J., & Kanani, M.R. (2011). Assessment of the potential ecological urban development using multi-criteria decision-making models of MCDM and GIS (Case study: City of Sari, Mazandaran). *Journal of Human Geography Research Quarterly*, 43(77), 75–88.
- Mohamadi, J., & Give, J. (2001). Land evaluation for irrigated wheat in Felaverjan Area (Isfahan) using Fuzzy sets theory. *Journal of Sciences and Technology of Agriculture and Natural Resources*, 5(1), 103–115.
- Mokhtari Kashki, D. (2010). Assessment of ecotourism potential of geomorphic sites at Asyab-Kharabeh Catchment area in North West of Iran by Prolong Method. *Geography and Development Iranian Journal*, 8(18), 27–52.
- Najafifar, A. (2011). The role of land use planning in optimal management of Zagros forests (Case study: Sarab, Darreh Shahr Basin, Ilam Province). *Iranian Journal of Rangelands and Forests Plant Breeding and Genetic Research*, 19(4), 510–522.
- Nazarian, A., Zeiaean, P., & Abdollahi, A.A. (2005). *Detection of optimal sites for future physical development of Kangan City using Boolean logic model and GIS and RS techniques*. 9<sup>th</sup> Symposium of Geological Society of Iran, 5-7 October 2005 (pp. 5-18). Tehran, University of Kharazmi, Geological Society of Iran.
- Nohegar, A., Hossein Zadeh, M.M., & Pirasteh, A. (2009). Assessment capabilities of ecotourism Qeshm island using SWOT strategy management model. *Iranian Journal of Geography and Development*, 7(15), 151–172.

- Salman Mahini, A.R., Reaizi, B., Naeimi, B., Babaie Kafaki, S., & Javadi Larijani, A. (2009). Evaluation of potential ecotourism in Behshahr City based on multi-criteria analysis method using GIS. *Journal of Environmental Science and Technology*, 11(1), 187–198.
- Salman Mahini, A.R., Reaizi, B., Naeimi, B., Babaie Kafaki, S., & Javadi Larijani, A. (2009). Evaluation of potential ecotourism in Behshahr City based on multi-criteria analysis method using GIS. *Journal of Environmental Science and Technology*, 11(1), 187–198.
- Sheeyeh, E. (1998). *An introduction to the principles of urban planning*. Tehran: Iran University of Science & Technology Press.
- Zareei, R., & Alsheikh, A.A. (2012). Modeling of urban development using cellular automation and genetic algorithm (Case study: City of Shiraz). *Journal of Research and Urban Planning*, 3(11), 1–16.

### References (in English)

- Bullard, R. D., Johnson, G.S., & Torres, A. O. (2003). Atlanta mega sprawl. Clark Atlanta University. U.S. State of Georgia: The Environmental Justice Resource Center.
- Chang, N. B., Parvathinathan, G., & Breeden, J. B. (2008). Combining GIS with fuzzy multicriteria decision making for land fill siting in a fast-growing urban region. *Journal of Environmental Management*, 1(87), 139 - 153.
- Davidson, D. A., Tehocharopoulos, S. P., & Jbloksma, R. (1994). A land evaluation project in Greece using GIS and based on Boolean and fuzzy set methodologies. *Journal of Geographical Information Systems*, 8(4), 369-889.
- Devana, A., M., & Yamaguchib, Y. (2009). Land use and land cover change in Dhaka, Bangladesh: Using remote sensing to promote sustainable urbanization. *Journal of Applied Geography*, 29(3), 390-401.
- Dixon, B. (2005). Ground water vulnerability mapping: A GIS and fuzzy rule based integrated tool. *Journal of Applied Geography*, 4(25), 327-347.
- Eastman, J. R. (2006). *IDRISI Andes tutorial*. Worcester, MA: Clark University.
- Eastman, J., Weigen, J., Peter, A., & Kwaku, K. (1995). Raster procedure for multicriteria multi objective decisions. *Journal of Photogram Metric Engineering & Remote Sensing*, 61(5), 539-547.
- Foroutan, E., & Delavar, M.R. (2012). *Urban growth modeling using fuzzy logic*. ASPRS2012 Annual Conference, 19-23 March 2012 (pp. 1-10). , Sacramento, Cardinal Systems Esri. .
- Kumar, K., & Lesslie, A. (2008). Geomorphological mapping planning and development- A case for Korba City, Chhattisgarh. *Journal of Research Communication*, 97(15), 12-25.
- Lotfi Zadeh, A. (1965). Fuzzy sets. *Journal of Information and Control*, 8(1), 338-353.
- May, M. E., Dlalala, M., & Chenini, I. (2010). Urban geological mapping: Geotechnical data analysis for rational development planning. *Journal of Engineering Geology*, 1(116), 129-138.

- McBratney, A. B., & Odeh I. O. A. (1997). Application of fuzzy sets in soil sciences: Fuzzy logic, measurements and fuzzy classifications. *Journal of Geoderma*, 77(1), 85–113.
- Mez'osi, G., Blanka, V., Bata, T., Kovács, F., & Meyer, B. (2015). Estimation of regional differences in wind erosion sensitivity in Hungary. *Journal of Natural Hazards and Earth System Science*, 1(15), 97- 107.
- Ronld, W. T. (1973). *Focus on environmental geology*. Oxford: Oxford University Press.
- Sui, Z. D. (1992). A fuzzy GIS modeling approach for urban land evaluation computers. *Journal of Environment and Urban System*, 16(2), 101-115.
- Sukoop, H., Numata, M., & Huber, A. (1995). *Urban ecology as basis of urban planning*. The University of Chicago Press Books. Chicago. State of Illinois: Chicago Distribution Center