

Received 2018-04-01
Revised 2018-04-07
Accepted 2018-04-08

The Role of Herbal Medicine as Anti-Cancer Medicine: From the Claim to Truth

Hoda Aryan^{1,2}✉

¹Young Researchers and Elite Club, Tehran Medical Sciences Branch, Islamic Azad University, Tehran, Iran

²Medical Students' Scientific Association (MSSA), Tehran Medical Sciences Branch, Islamic Azad University, Tehran, Iran

Dear Editor,

One of the main causes of death in the world is cancer, so that has involved more than one-third of the population, and is considered as the cause of more than 20 percent of all deaths in the world [1]. More than 100 different types of cancer have been identified. Regarding the International Agency for Research on Cancer (IARC) reports, the most commonly diagnosed cancers worldwide were those of the lung (1.8 million, 13.0% of the total), breast (1.7 million, 11.9%), and colorectum (1.4 million, 9.7%) [2]. If the cancer is detected in its early stages, surgery or radiation therapy can be used to treat, but advanced cancer treatment needs chemotherapy. Although chemotherapy drugs are effective, their use is associated with abundant adverse effects and drug resistance [3]. Herbal remedies have been used over the centuries to treat a variety of diseases. Most of these treatments as alternatives or supplements to consider a way that helps patients to have better physical and mental status. Currently, many in-vitro and in vivo studies indicated the beneficial effect of medicinal plants and their bioactive compounds could induce the apoptosis in neoplastic cells [4-8]. Previous studies showed the anti-cancer activity of the ten widely used herbs that are commonly used in the context of cancer by patients

in the Middle East: *Olea Europea* (Olive), *Nigella Sativa* (Black Seeds), *Crocus Sativus* (Saffron), *Punica Granatum* (Pomegranate), *Urtica Dioica* (Nettle), *Allium Sativum L.* (Garlic), *Allium Ceba* (Onion), *Curcuma longa* (Curcumin), *Arum Palaestinum* (Palestinian Arum), and *Vitis Vinifera* (Grapes) [9,10]. However, researchers have believed to confirm the usefulness of herbal medicines in the treatment or prevention of the cancer is essential to conduct extensive clinical trials to determine what plants can be used alongside conventional therapeutic methods for cancer treatment. Nowadays, more than 25% of drugs used during the last 20 years are directly derived from plants, while the other 25% are chemically altered natural products. Still, only 5–15% of the approximately 260,000 higher plants have ever been investigated for bioactive compounds [3,4]. The advantage of using such compounds for cancer treatment is their relatively low/non-toxic nature. According to the WHO, 80 percent of the world benefit from traditional treatment. Sixteen percent of drugs approved by the FDA in the years 1984 to 1994 have been obtained from natural resources, especially plants [4]. Vinca Alkaloids (Vinblastine and Vincristine), Taxanes, podophyllotoxin are the best known FDA approved plant-derived anticancer agents. Vinblastine and Vincristine are the first herbal anti-cancer medicine that intro-

GMJ

Copyright© 2018, Galen Medical Journal. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>)
Tel/Fax: +98 71 36474502
Email: info@gmj.ir



✉ Correspondence to:

Hoda Aryan, Young Researchers and Elite Club, Tehran Medical Sciences Branch, Islamic Azad University, Tehran, Iran
Telephone Number: +982122006660
Email Address : hodaaryan@gmj.ir

duced in 1960 which are widely used to treat breast cancer, Hodgkin's Lymphoma, William's tumor, lymphoblastic leukemia and Kaposi's sarcoma [4-6]. In 1992, Taxanes, the most efficient anti-tumor agent, was approved by FDA to treat breast, head, and neck, prostate, and gastric cancers [4,6,7]. Etoposide and teniposide are two active and semi-synthetic compounds of Epipodophylotoxin that is approved by FDA for the treatment of choriocarcinoma, lung cancer, ovari-

an and testicular cancers, lymphoma and acute myeloid leukemia. [8] Considering the fact that little is known about efficacy and safety of herbal products, and not paying attention to commonly used products setting, further research can improve appropriate use of plant products drastically. [GMJ.2018;7:e1179]
DOI: 10.22086/gmj.v0i0.1179

Keywords: Cancer; Herbal Medicine; Complementary; Anti-cancer

References

1. Mirmalek S, Jangholi E, Jafari M, Yadollah-Damavandi S, Javidi M, Parsa Y, et al. Comparison of in Vitro Cytotoxicity and Apoptogenic Activity of Magnesium Chloride and Cisplatin as Conventional Chemotherapeutic Agents in the MCF-7 Cell Line. *Asian Pac J Cancer Prev*. 2016;17:131-134.
2. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2015;136: E359-E386.
3. Mirmalek SA, Azizi MA, Jangholi E, Yadollah-Damavandi S, Javidi MA, Parsa Y, et al. Cytotoxic and apoptogenic effect of hypericin, the bioactive component of *Hypericum perforatum* on the MCF-7 human breast cancer cell line. *Cancer Cell Int*. 2015; 16:3.
4. Hosseini MM, Karimi A, Behroozaghdam M, Javidi MA, Ghiasvand S, Berimipour A, et al. Cytotoxic and Apoptogenic Effects of Cyanidin-3-Glucoside on the Glioblastoma Cell Line. *World Neurosurg*. 2017;108:94-100.
5. Safarzadeh E, Sandoghchian Shotorbani S, Baradaran B. Herbal medicine as inducers of apoptosis in cancer treatment. *Adv Pharm Bull*. 2014 ;4:421-427.
6. Balunas MJ, Kinghorn AD. Drug discovery from medicinal plants. *Life Sci*. 2005;78:431-441.
7. Cragg GM, Newman DJ. Plants as a source of anti-cancer agents. *J Ethnopharmacol*. 2005;100:72-79.
8. Nobili S, Lippi D, Witort E, Donnini M, Bausi L, Mini E, et al. Natural compounds for cancer treatment and prevention. *Pharmacol Res*. 2009;59:365-578.
9. Ben-Arye E, Ali-Shtayeh MS, Nejmi M, Schiff E, Hassan E, Mutafoglu K, et al. Integrative oncology research in the Middle East: weaving traditional and complementary medicine in supportive care. *Support Care Cancer*. 2012;20:557-564.
10. Zaid H, Silbermann M, Ben-Arye E, Saad B. Greco-Arab and Islamic Herbal-Derived Anticancer Modalities: From Tradition to Molecular Mechanisms. *Evid Based Complement Alternat Med*. 2012; 2012: 349040.