



# Relationship Between Posttraumatic Growth and Meaning in Life Among Patients with Myocardial Infarction

Seyede Zahra Hosseinigolafshani <sup>1,\*</sup>, Farshad Taheri<sup>2</sup>, Hamid Sharif Nia<sup>3</sup> and Sakineh Moghaddam Zeabadi<sup>4</sup>

<sup>1</sup>Department of Critical Care Nursing, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>2</sup>Student Research Committee, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>3</sup>School of Nursing and Midwifery Amol, Mazandaran University of Medical Sciences, Sari, Iran

<sup>4</sup>Emergency Department, School of Paramedicine, Qazvin University of Medical Sciences, Qazvin, Iran

\*Corresponding author: Assistant Professor, Department of Critical Care Nursing, Qazvin University of Medical Sciences, Qazvin, Iran. Email: z.hoseinigolafshani@qums.ac.ir

Received 2019 May 09; Revised 2019 June 19; Accepted 2019 June 24.

## Abstract

**Background:** A great number of people who have survived after stressful events experience positive psychological changes. These positive changes are recognized as posttraumatic growth.

**Objectives:** The purpose of the current study was to examine the relationship between posttraumatic growth and meaning in life in patients with myocardial infarction.

**Methods:** This is a descriptive correlational study conducted on 191 patients referring to the largest cardiology center (Bu-Ali Sina) in Qazvin, Iran. The sample was selected through a convenience sampling method based on the inclusion criteria. The data were collected using a demographic questionnaire, the posttraumatic growth inventory, and the meaning in life questionnaire. Data were analyzed with descriptive analysis. Inferential statistics (Pearson's test) were used to evaluate the relationship between posttraumatic growth and meaning in life.

**Results:** The sample included 49 (25.7%) women and 142 (74.3%) men with the mean age of  $59.6 \pm 1.2$  years. The mean total score of posttraumatic growth was  $64.6 \pm 11.5$  and the mean total score of meaning in life was  $49 \pm 6.04$ . Based on the results, there was a significant positive correlation between posttraumatic growth and meaning in life ( $r = 0.71, P < 0.001$ ).

**Conclusions:** The occurrence of myocardial infarction can lead to some positive psychological changes called posttraumatic growth. Nurses, informal caregivers, and policymakers can use these findings to help patients cope with challenging conditions after myocardial infarction.

**Keywords:** Posttraumatic Growth, Meaning in Life, Myocardial Infarction, Iran

## 1. Background

In today's life, we are witnessing the occurrence of stress and its complications more than ever. Psychologically, people who suffer from repeated, prolonged stress in life are more likely to experience loneliness, social isolation, sense of difference with peers, poor education, and limited community participation (1). However, recent studies have addressed new concepts in this matter using the Seligman positive psychology approach and argued that the opposite side of people who give disappointing answers to severe crises are those who have more effective responses and choose unique and creative ways to manage their position; they not only adapt to crises and changes caused by them and accept them as a stage of life, but also often go beyond their previous position and experience

great profoundness and evolution in various aspects of life (2).

In fact, the results of studies conducted using this approach since 1996 have shown that a great number of people who have survived after stressful events experience positive psychological changes. These positive changes can be caused by stressful events or they might be a type of education that a person learns when tries to adapt to stressful events (3-5). An interesting point is that the review of the literature over the past decade in this matter suggests that, contrary to expectations, the evidence of growth experiences is much greater than the evidence of posttraumatic stress disorder (PTSD) at different levels (6). This positive change, which is recognized as posttraumatic growth (PTG), is the experience or mental perception of positive psychological changes that result from the struggle with a

stressful event and includes the growth and excellence in five categories: (1) new possibilities, (2) relating to others, (3) personal strength, (4) spiritual change, and (5) appreciation of life. Special attention to the term “struggle” in the literature is related to the reason that growth is not necessarily the result of an event, but the individual’s efforts to cope with new realities are crucial in determining the extent of growth experience (4, 7, 8).

In several studies, the concept of PTG has been studied with other terms such as “benefit finding”, “stress-related growth”, and “flourishing”. However, most researchers in this field tend to use the term “posttraumatic growth” to describe such positive changes (5, 9). A review of the literature shows that most studies examined posttraumatic growth and its association with various factors in non-clinical samples. This is despite the fact that posttraumatic growth can be the subject of research in patients experiencing chronic diseases or medical conditions (4, 8).

The high rates of chronic diseases are one of the new challenges that the healthcare system is facing; the coronary artery disease, and at the top of it, myocardial infarction, places in this category (10, 11). In the United States, 1.5 million people are affected by myocardial infarction each year, leading to 400,000 deaths. In Iran, of all the 700 to 800 deaths per day, 317 are due to cardiovascular diseases and 166 of them are because of myocardial infarction. The direct and indirect costs of cardiovascular diseases in the United States are estimated to be 4753 billion dollars a year. In Iran, 15 billion dollars a year are spent on treatment besides 50 million dollars on the purchase of related medical equipment (10).

Myocardial infarction is one of the most stressful events that can cause serious physical, psychological, and social problems for patients. Patients with myocardial infarction face multiple sequelae such as congestive heart failure, cardiac arrhythmias, unemployment, physical disabilities, and mental disorders such as anxiety and depression. These problems can affect patients’ well-being and quality of life. On the other hand, the Seligman’s positive psychological approach can stimulate the development of positive attitudes and modify the individuals’ perception and the philosophy of life, creating the ground for making positive changes in people who experience life-threatening illnesses such as myocardial infarction that suddenly and unexpectedly claim the patients’ lives (12, 13). The results of several studies investigating the concept of posttraumatic growth in post-myocardial infarction patients suggest that people first affected by infarction and then experiencing posttraumatic growth during the course of disease are less likely to have recurrent heart attack than other patients are; moreover, the occurrence of related unpleasant side effects is also lower due to the pres-

ence of several and different support systems along with the improvement of the patients’ ability to adapt better to new stressors (4, 5, 7).

Meaning in life refers to a sense of connection with the creator of the universe, having a goal in life, pursuing and achieving valuable goals, and ultimately achieving evolution. One of the most famous definitions of meaning in life was presented by Emil Frankl (1982). He believes that when a person feels that his being is linked to a perfect source and sees himself reliant on extensive and reliable frameworks such as the religion and philosophy that he chose to live in, he finds the meaning and feels it (14, 15). These people show resilience and tolerance against stressful events of life (9, 16).

In general, the perception of the meaning of life (MOL) is one of the strong predictors of well-being and life satisfaction. The results of studies also show that having a meaningful life has a significant positive correlation with hope, happiness, life satisfaction, and quality of life and a significant negative association with depression, anxiety, and mental disorders. All of these verify that a meaningful life is a valuable indicator for assessing mental health in individuals (17).

The results of some studies indicate that meaning in life is one of the variables associated with posttraumatic growth (18). However, some other studies have not shown a clear relationship between the two mentioned concepts (3, 19, 20). In a study by Rahimi and Heidarzadeh on patients with myocardial infarction experiencing a traumatic event, the mean total score of posttraumatic growth was moderate to high (13). In another study, Mousavi and Vatankhah stated that meaningful life was a positive predictor of posttraumatic growth in women with breast cancer (21).

Since myocardial infarction is the most common and a major cause of death in the world and its consequences can affect the lives of individuals and the community, the understanding of different dimensions of the concept of posttraumatic growth and the meaning of life, and more importantly, examining the type and the relationship between these concepts based on different components can provide valuable information for patients and healthcare providers. Applying these concepts to plan daily care and interactions leads to a more effective compromise of patients with stressful situations and better care provided by their caregivers.

## 2. Objectives

Considering the relationship between posttraumatic growth and social, cultural, and supporting systems of the target population (22) and the lack of sufficient studies on

the relationship between these two variables in patients with myocardial infarction in Iran, the aim of this study was to investigate the relationship between posttraumatic growth and meaning in life in myocardial infarction patients.

### 3. Methods

#### 3.1. Study Design and Population

The present descriptive-correlational study was performed on a sample of 191 patients referring to the largest cardiology center (Bu-Ali Sina) in Qazvin, Iran. Patients were selected through a convenience sampling method based on the inclusion criteria. The inclusion criteria included the age of 18 years or older, the definitive diagnosis of myocardial infarction (for the first time) by a cardiologist, the ability to read and write, the experience of at least three months from the onset of myocardial infarction (23), and performing PCI process in all samples (24, 25). The exclusion criteria were unwillingness to participate in the study, experience of severe stress during the past month based on the patient's self-report, having mental disorders during data collection based on the information recorded in the patient's file, and inappropriate physical condition at the time of data collection if prevented the patient from responding appropriately to questions (Figure 1). After explaining the research objectives and how to participate in the study, written consent was obtained from the patients.

#### 3.2. Sample Size

The sample size was calculated using the relevant statistical formula considering the significance level of 0.05, the power of 80%, and the effect size of  $d = 0.2$  obtained according to the coefficient of determination (0.04) using G\*power 3.1.7 software.

- Analysis: A priori: Computing required sample size
- Input: Tail(s) = Two
- Effect size  $|\rho| = 0.2000000$
- $\alpha$  err prob = 0.05
- Power ( $1 - \beta$  err prob) = 0.8
- Output: Noncentrality parameter  $\delta = 2.8210518$
- Critical t = 1.9725951
- Df = 189
- Total sample size = 191

#### 3.3. Instruments

The instruments used in the present study included a demographic characteristics questionnaire, the posttraumatic growth inventory (PTGI), and the meaning in life questionnaire (MLQ).

The demographic characteristics consisted of data about patients' age, gender, marital status, level of education, and occupation.

#### 3.4. Posttraumatic Growth Inventory (PTGI)

The posttraumatic growth inventory (PTGI; Tedeschi and Calhoun, 1996) was used to assess growth-related changes experienced by traumatic individuals (26). This 21-item scale evaluates five domains of PTG consisting of (1) new possibilities, (2) relating to others, (3) personal strength, (4) spiritual change, and (5) appreciation of life. The items are rated from 0 (I did not experience this situation as a consequence of my crisis) to 5 (I fully experienced this change to a very great degree). The total score ranges from 0 to 105 with a higher score indicating a greater experience of posttraumatic growth (4). The reliability and validity of the scale were evaluated in a sample of patients with cardiac disorders in the United States by Sheikh and Marotta. The obtained Cronbach  $\alpha$  for the total scale was 0.96 (27).

Triplett et al. reported that the internal reliability of the scale was 0.90 and the test-retest consistency with a two-month interval was 0.71 (18, 28). The PTGI was translated into Persian by Mahmoudi (2008) and then revised and adapted by Mahmoudi et al. According to the results of PTGI construct analysis, three factors were confirmed including changes in the relationship with others ( $\alpha = 0.86$ ), philosophy of life ( $\alpha = 0.87$ ) and self-perception ( $\alpha = 0.88$ ) in an Iranian population (29, 30).

#### 3.5. The Meaning in Life Questionnaire (MLQ)

This instrument was designed by Steger et al. to assess the presence of meaning in life and search for it. The validity, reliability, and factor structure were studied in various studies with different samples. The researchers found two factors, the presence of meaning in life and search for meaning in life, with a total number of 17 items using exploratory factor analysis. Then, in confirmatory factor analysis, seven items were removed and a suitable structure of two factors with 10 items was achieved. According to Steger et al., the validity of the subscale "the presence of meaning" was 0.86 and that of the subscale "search for meaning" was 0.87. Moreover, the reliability coefficients of the subscales "the presence of meaning" and "search for meaning" were 0.70 and 73.1, respectively (31). Jamali obtained its reliability coefficient as 0.9 using Cronbach's alpha (32). In addition, the reliability coefficient of the MLQ was determined to be 0.89 in Peymanfar and Akbari study using Cronbach's alpha (33). The scale for scoring the items of the questionnaire included absolutely untrue (1), mostly untrue (2), somewhat untrue (3), cannot say true or false




Figure 1. The study flow chart

(4), somewhat true (5), mostly true (6), and absolutely true (7). The scores of items 2, 3, 7, 8, and 10 are summed up to obtain the subscale score of “search for meaning” and the scores of items 1, 4, 5, 6, and 9 are summed up to calculate the subscale score of “the presence of meaning”. The total score of the scale ranges from 10 to 70. A higher score indicates a greater meaning in one’s life and a lower score indicates a lower meaning.

### 3.6. Ethical Consideration

This study was approved by the Ethics Committee of Qazvin University of Medical Sciences (IR.QUMS.REC.1397.022). Before signing informed consent forms, all the participants were informed about the study objectives and procedures. They were also assured that participation in the study would be voluntary and would not affect their medical care. Permission was obtained from the PTGI and MLQ developers (Tedeschi and Calhoun (5) and Steger et al. (31) respectively).

### 3.7. Statistical Analysis

Data were analyzed using SPSS (version 24). Descriptive analysis, frequency, mean, and standard deviation (SD) were used to describe the participants’ demographic variables. Inferential statistics (Pearson’s test) were used to evaluate the relationship between PTG and MOL. The significance level was set at  $P < 0.05$ .

## 4. Results

### 4.1. Socio-Demographic Characteristics of Patients with Myocardial Infarction

Based on the results of the study (Table 1), the mean age of the patients was  $59.6 \pm 1.2$  years. Of the 191 participants, 49 (25.7%) were female and 142 (74.3%) were male. Moreover, 5 (2.6%) patients were single and 186 (97.4%) were married. In terms of education level, 36 (18.8%) were illiterate, 117 (61.3%) were under diploma, 30 (15.7%) had diploma degrees, and 8 (4.2%) had university degrees. Regarding the occupation status, 133 (69.6%) were employed, 51 (26.7%) were retired, and 7 (3.6%) were unemployed.

### 4.2. Posttraumatic Growth and Its Dimensions

Based on the results (Table 2), the mean total score of posttraumatic growth in patients was  $64.6 \pm 11.5$ . In terms of the domains of posttraumatic growth, the mean scores of appreciation of life, new possibilities, relating to others, personal strength, and spiritual change were 9.4, 15.8, 21.2, 12.4, and 5.7, respectively. Given the unequal number of items in each domain, the weighted mean scores of domains (based on the number of items of each domain)

**Table 1.** Sociodemographic Characteristics of Patients with Myocardial Infarction<sup>a</sup>

Variable	Values
Age, mean $\pm$ SD	12.1 $\pm$ 59.6
<b>Gender</b>	
Female	49 (25.7)
Male	142 (73.3)
Total	191 (100)
<b>Marital status</b>	
Single	5 (2.6)
Married	186 (97.4)
Total	191 (100)
<b>Education level</b>	
Illiterate	36 (18.8)
Under diploma	11 (61.3)
Diploma	30 (15.7)
Academic	8 (4.2)
Total	191 (100)
<b>Occupation status</b>	
Employed	133 (69.6)
Retired	51 (26.8)
Unemployed	7 (3.6)
Total	191 (100)

<sup>a</sup>Values are expressed as No. (%) unless otherwise indicated.

were estimated to be 3.13, 3.16, 3.02, 3.1, and 2.8, respectively. Accordingly, the subscale “new possibilities” scored the highest and “spiritual change” scored the lowest.

### 4.3. Meaning in Life and its Dimensions

Based on the results (Table 3), the mean total score of meaning in life in patients was  $49 \pm 6.04$ . The mean scores of subscales “presence of meaning” and “search for meaning” were 26.03 and 23, respectively.

### 4.4. Relationship Between MOL and PTG

Based on the results (Tables 4 and 5), there was a significant positive correlation between posttraumatic growth and meaning in life ( $r = 0.71$ ,  $P < 0.001$ ). Thus, the mean score of posttraumatic growth increases by an increase in the score of the meaning in patients’ lives. Pearson correlation coefficient showed significant positive correlations between meaning in life and all the domains of posttraumatic growth ( $P < 0.05$ ).

**Table 2.** Posttraumatic Growth and Its Dimensions Scores

Variable	Mean $\pm$ SD	Min	Max	Number of Items	Weighted Average
Appreciation for life	9.4 $\pm$ 1.9	3	15	3	3.13
New possibilities	15.8 $\pm$ 3.4	4	23	5	3.16
Relating to others	21.2 $\pm$ 4.1	7	30	7	3.02
Personal strength	12.4 $\pm$ 2.5	4	18	4	3.1
Spiritual change	5.7 $\pm$ 1.07	1	8	2	2.8
Posttraumatic growth	64.6 $\pm$ 11.5	21	92	21	3.07

**Table 3.** Meaning of Life and its Dimension Scores

Variable	Min	Max	Mean $\pm$ SD
The existence of meaning in life	15	34	26.03 $\pm$ 3.9
Search for meaning in life	16	29	23 $\pm$ 2.4
Meaning of life	32	63	49 $\pm$ 6.04

**Table 4.** Correlation Matrix of Posttraumatic Growth and Meaning of Life in Patients

Variable	Meaning of Life	Posttraumatic Growth
<b>Posttraumatic growth</b>		
r	0.71	1
P value	< 0.001	-
<b>Meaning of life</b>		
r	1	-
P value	-	-

## 5. Discussion

This study aimed to investigate the relationship between posttraumatic growth and meaning in life among patients with myocardial infarction. The results indicated that the mean total score of posttraumatic growth was moderate to high among the participants (64.6  $\pm$  11.5), which demonstrated a degree of growth experience in patients. Consistent with the results of this study, the mean total score of posttraumatic growth was reported high by Zarin et al. (27) in patients with spinal cord injury and Rahimi and Heidarzadeh (13) in patients with myocardial infarction. Heidarzadeh et al. (34) and Jansen et al. (35) stated that the mean total score of posttraumatic growth was moderate to high in cancer patients. Meanwhile, the results of Morris et al. (22) and Bellizzi et al. (36) showed that the mean total score of posttraumatic growth was low in patients with cancer. Regardless of the cultural differences between participants in the above studies, the results of the present study confirmed the theoretical framework of the concept of posttraumatic growth, which believes that exposure to stressful events, such as myocar-

dial infarction, could be associated with the experience of growth in various areas.

Based on the results of the present study, the domain of "new possibilities" scored the highest and the domain of "spiritual change" scored the lowest among the five domains of posttraumatic growth. It means that participants in the study experienced the most growth in the domain of "new possibilities" and the least growth in "spiritual change". Consistent with the results of the present study, the results of studies by Brix et al. (37), Morris et al. (22), and Bellizzi et al. (36) showed that the domain of spiritual change had the least growth in patients with cancer. Hooper et al. found that the highest growth was in terms of new possibilities and the least growth was in the domain of spiritual change (38). Contrary to the present study, Rahimi and Heidarzadeh (13) study in patients with myocardial infarction and Heidarzadeh et al. (34) study in cancer patients reported the highest growth in the domain of spiritual change and the least growth in the domain of new possibilities. On the other hand, Aflakseir et al. (39) in women with breast cancer and Teodorescu et al. (40) in Norwegian immigrants concluded that the lowest posttraumatic growth was in the domain of new possibilities. The contradictory results of studies concerning the scores of different domains of posttraumatic growth in different events and societies can be reasonably expected due to differences in the type and quality of the experienced stress, cultural differences, religious teaching, and values of each society that requires further in deep examination using a qualitative approach.

In terms of meaning in life, the results of the study showed that the mean total score of meaning in life was 49  $\pm$  6.04 among the participants. The mean scores of "presence of meaning" and "search for meaning" subscales were 26.03  $\pm$  3.9 and 23  $\pm$  2.4, respectively. Consistent with the current study, Abedi et al. stated that elderly people achieved a higher score on meaning in life (41). Thompson also reported that breast cancer patients had a high meaning in life (42). Reza Zadeh and Rahmani Asl (9) and Park and Baumeister (16) believe that individuals are more re-

**Table 5.** Relationship Between the Components of Posttraumatic Growth and Meaning of Life in Patients

Posttraumatic Growth	Appreciation for Life	New Possibilities	Relating to Others	Personal Strength	Spiritual Change	Total
<b>Meaning of life</b>						
r	0.61	0.67	0.66	0.62	0.62	0.71
P value	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001

sistant and patient in facing stressful events if they have a meaning in life. Several studies have shown meaning in life had a direct and meaningful relationship with quality of life, satisfaction, and optimism while it was inversely related to anxiety, depression, and psychological distress (43). Nasiri showed a significant positive correlation between meaning in life and life expectancy, happiness, and life satisfaction and a significant negative correlation between meaning in life and depression (17). All of these results indicate that patients participating in the present study had a valuable indicator for assessing mental health by having a high mean score of meaning in life.

Contrary to the results of the present study, Kord and Rahbari reported that the mean score of “search for meaning” was high in patients with cardiovascular diseases (44). Hassankhani et al. (45) and Eric (46) reported low meaning in life in cancer patients. After experiencing severe stress, the person who searches for the meaning needs some times to achieve a fixed and definite meaning. Therefore, one of the reasons for the contradictory results of the above studies can be the different times of the studies.

In terms of the relationship between the domains of posttraumatic growth and meaning in life, the results of the study showed a significant positive relationship between the scores of posttraumatic growth and meaning in life ( $P < 0.05$ ). Patients achieved higher scores on different domains of posttraumatic growth by an increase in the scores of meaning in life. Consistent with the results of this study, Mousavi and Vatankhah stated that meaning in life was a predictor of posttraumatic growth in women with breast cancer (21). Dursun et al. showed a significant positive relationship between posttraumatic growth and meaning in life in university students (47). Garcini et al. concluded that students whose lives were more meaningful could cope better with the anxiety and challenges of life (48). Shafiee et al. stated that the mean score of meaning in life was lower in the injured women than in other women (49). According to the researchers, experiencing posttraumatic growth can facilitate the process of recognizing the stressful events in patients, create a positive view in patients and their relatives, and change their lifestyles; this concept creates meaningful positive changes in the emotional and cognitive lives of individuals, which positively affect their behaviors and functions (4, 5, 50). Healthcare

providers can apply the concepts to plan the daily care and interactions, which may lead to an effective compromise of patients with stressful events and better care provided by caregivers.

### 5.1. Conclusions

The understanding of variables related to posttraumatic growth can help clinicians change this process in a useful way. We believe that clinicians are the facilitators of this process because posttraumatic growth is likely inhibited by heavy-handed attempts to move trauma survivors toward the understanding they have not yet directly experienced.

### 5.2. Limitations and Suggestions

The use of self-report tools was one of the limitations of this study. Patients might not have a clear understanding of the concepts of this study to answer the related questions. Thus, it is suggested that extensive studies concentrate on qualitative factors and open interviews.

It is also suggested that the predictors of posttraumatic growth be evaluated in Iranian society in future studies to effectively plan and provide a context for improving traumatic patients' adaptability, development, and quality of life.

### Acknowledgments

The authors would like to appreciate the authority of the Department of Nursing of Qazvin University of Medical Sciences, the staff of Bu-Ali Sina Hospital for their cooperation in data collection, and all myocardial infarction patients who participated in this study.

### Footnotes

**Authors' Contribution:** Seyedeh Zahra Hosseinigolafshani was the main investigator. Farshad Taheri collected the data and wrote the first draft. Hamid Sharif Nia analyzed the data. Seyedeh Zahra Hosseinigolafshani led the team designed the study and read and correct the final draft. Sakineh Moghaddam Zeabadi collected the data and helped in writing the final draft.

**Conflict of Interests:** The authors declare no conflict of interests.

**Ethical Considerations:** IR.QUMS.REC.1397.022.

**Funding/Support:** This research project was approved and supported by the Research Deputy of Qazvin University of Medical Sciences (IR.QUMS.REC.1397.022).

## References

- Khodayari Fard M, Parand A. [Stress and ways to deal with it]. Tehran: University of Tehran; 2013. Persian.
- Abedi H, Hosseini Golafshani S, Ahmadi F. Tailor the experiences of people with disabilities: Circumstances that cause growth or impediment to growth. *J Qual Res Health Sci*. 2013;**2**(3):248-60.
- Dursun P, Steger MF, Bentele C, Schulenberg SE. Meaning and post-traumatic growth among survivors of the september 2013 colorado floods. *J Clin Psychol*. 2016;**72**(12):1247-63. doi: [10.1002/jclp.22344](https://doi.org/10.1002/jclp.22344). [PubMed: [27459242](https://pubmed.ncbi.nlm.nih.gov/27459242/)].
- Rahimi R, Heidarzadeh M, Shoaee R. The Relationship between Post-traumatic Growth and Social Support in Patients with Myocardial Infarction. *Can J Cardiovasc Nurs*. 2016;**26**(2):19-24. [PubMed: [27382668](https://pubmed.ncbi.nlm.nih.gov/27382668/)].
- Tedeschi RG, Calhoun LG. The posttraumatic growth inventory: Measuring the positive legacy of trauma. *J Trauma Stress*. 1996;**9**(3):455-71. doi: [10.1007/BF02103658](https://doi.org/10.1007/BF02103658). [PubMed: [8827649](https://pubmed.ncbi.nlm.nih.gov/8827649/)].
- Tedeschi RG, Calhoun LG. 2004 , editor. *Target article: "Posttraumatic growth: Conceptual foundations and empirical evidence"*. 15. Philadelphia, PA: Lawrence Erlbaum Associates; 2004. doi: [10.1207/s15327965pli1501\\_01](https://doi.org/10.1207/s15327965pli1501_01).
- Garnefski N, Kraaij V, Schroevers MJ, Somsen GA. Post-traumatic growth after a myocardial infarction: A matter of personality, psychological health, or cognitive coping? *J Clin Psychol Med Settings*. 2008;**15**(4):270-7. doi: [10.1007/s10880-008-9136-5](https://doi.org/10.1007/s10880-008-9136-5). [PubMed: [19104983](https://pubmed.ncbi.nlm.nih.gov/19104983/)].
- Gul E, Karanci AN. What determines posttraumatic stress and growth following various traumatic events? A study in a Turkish community sample. *J Trauma Stress*. 2017;**30**(1):54-62. doi: [10.1002/jts.22161](https://doi.org/10.1002/jts.22161). [PubMed: [28103412](https://pubmed.ncbi.nlm.nih.gov/28103412/)].
- Reza Zadeh H, Rahmani Asl M. [Comparing the meaning of life from the perspective of Mohammad Taghi Jafari Tabrizi and Friedish Nijah]. *Religious Sci J*. 2014;**10**(30):115-34. Persian.
- Baghaee R, Alinejad V, Sharifi A. [Impact of cardiac rehabilitation on myocardial infarction patient's anxiety]. *J Urmia Nurs Midwifery Fac*. Persian.
- Thrift AG, Thayabaranathan T, Howard G, Howard VJ, Rothwell PM, Feigin VL, et al. Global stroke statistics. *Int J Stroke*. 2017;**12**(1):13-32. doi: [10.1177/1747493016676285](https://doi.org/10.1177/1747493016676285). [PubMed: [27794138](https://pubmed.ncbi.nlm.nih.gov/27794138/)].
- Gardner MH, Mrug S, Schwebel DC, Phipps S, Whelan K, Madan-Swain A. Demographic, medical, and psychosocial predictors of benefit finding among caregivers of childhood cancer survivors. *Psychosomatics*. 2017;**26**(1):125-32. doi: [10.1002/pon.4014](https://doi.org/10.1002/pon.4014). [PubMed: [26489522](https://pubmed.ncbi.nlm.nih.gov/26489522/)]. [PubMed Central: [PMC4959962](https://pubmed.ncbi.nlm.nih.gov/PMC4959962/)].
- Rahimi R, Heidarzadeh M. [Post-traumatic growth among patients with myocardial infarction]. *Cardiovascular Nurs J*. 2015;**4**(2):44-52. Persian.
- Ho MY, Cheung FM, Cheung SF. The role of meaning in life and optimism in promoting well-being. *Pers Individ Dif*. 2010;**48**(5):658-63. doi: [10.1016/j.paid.2010.01.008](https://doi.org/10.1016/j.paid.2010.01.008).
- Lavasani M, Ejei JM. The Relationship between meaning of life and optimism with subjective well-being. *J Psychol*. 2013;**17**(3):17.
- Park J, Baumeister RF. Meaning in life and adjustment to daily stressors. *J Posit Psychol*. 2016;**12**(4):333-41. doi: [10.1080/17439760.2016.1209542](https://doi.org/10.1080/17439760.2016.1209542).
- Nasiri HJ. [Meaning in life, hope, life satisfaction and mental health in women]. *Women Dev Polit*. 2008;**6**(2):157-76. Persian.
- Triplett KN, Tedeschi RG, Cann A, Calhoun LG, Reeve CL. Posttraumatic growth, meaning in life, and life satisfaction in response to trauma. *Psychol Trauma Theory, Res Pract Policy*. 2012;**4**(4):400-10. doi: [10.1037/a0024204](https://doi.org/10.1037/a0024204).
- Linley PA, Joseph S. Meaning in life and posttraumatic growth. *J Loss Trauma*. 2011;**16**(2):150-9. doi: [10.1080/15325024.2010.519287](https://doi.org/10.1080/15325024.2010.519287).
- Merrill N, Waters TE, Fivush R. Connecting the self to traumatic and positive events: links to identity and well-being. *Memory*. 2016;**24**(10):1321-8. doi: [10.1080/09658211.2015.1104358](https://doi.org/10.1080/09658211.2015.1104358). [PubMed: [26512915](https://pubmed.ncbi.nlm.nih.gov/26512915/)].
- Mousavi SV, Vatankhah M. [Prediction of posttraumatic growth through religion, meaning of life and social support in female breast cancer]. *J Achiev Clin Psychol Shahid Chamran Univ Ahvaz*. 2014;**1**(3):33-48. Persian. doi: [10.22055/jacp.2015.12405](https://doi.org/10.22055/jacp.2015.12405).
- Morris BA, Shakespeare-Finch J, Scott JL. Posttraumatic growth after cancer: The importance of health-related benefits and newfound compassion for others. *Support Care Cancer*. 2012;**20**(4):749-56. doi: [10.1007/s00520-011-1143-7](https://doi.org/10.1007/s00520-011-1143-7). [PubMed: [21494782](https://pubmed.ncbi.nlm.nih.gov/21494782/)].
- Grewal K, Stewart DE, Abbey SE, Leung YW, Irvine J, Grace SL. Timing of depressive symptom onset and in-hospital complications among acute coronary syndrome inpatients. *Psychosomatics*. 2010;**51**(4):283-8. doi: [10.1176/appi.psy.51.4.283](https://doi.org/10.1176/appi.psy.51.4.283). [PubMed: [20587755](https://pubmed.ncbi.nlm.nih.gov/20587755/)]. [PubMed Central: [PMC4474645](https://pubmed.ncbi.nlm.nih.gov/PMC4474645/)].
- Husson O, Zebrack B, Block R, Embry L, Aguilar C, Hayes-Lattin B, et al. Posttraumatic growth and well-being among adolescents and young adults (AYAs) with cancer: A longitudinal study. *Support Care Cancer*. 2017;**25**(9):2881-90. doi: [10.1007/s00520-017-3707-7](https://doi.org/10.1007/s00520-017-3707-7). [PubMed: [28424888](https://pubmed.ncbi.nlm.nih.gov/28424888/)]. [PubMed Central: [PMC5527055](https://pubmed.ncbi.nlm.nih.gov/PMC5527055/)].
- Staniute M, Brozaitiene J, Bunevicius R. Effects of social support and stressful life events on health-related quality of life in coronary artery disease patients. *J Cardiovasc Nurs*. 2013;**28**(1):83-9. doi: [10.1097/JCN.0b013e318233e69d](https://doi.org/10.1097/JCN.0b013e318233e69d). [PubMed: [22067721](https://pubmed.ncbi.nlm.nih.gov/22067721/)].
- Ochoa C, Casellas-Grau A, Vives J, Font A, Borrás JM. Positive psychotherapy for distressed cancer survivors: Posttraumatic growth facilitation reduces posttraumatic stress. *Int J Clin Health Psychol*. 2017;**17**(1):28-37. doi: [10.1016/j.ijchp.2016.09.002](https://doi.org/10.1016/j.ijchp.2016.09.002). [PubMed: [30487878](https://pubmed.ncbi.nlm.nih.gov/30487878/)]. [PubMed Central: [PMC6236322](https://pubmed.ncbi.nlm.nih.gov/PMC6236322/)].
- Zarin S, Khanjani M, Foroughan M, Hosseini M, Bakhshi E, Kamali M. Relationship between locus of control with posttraumatic growth among individuals with spinal cord injury. *J Mod Rehabil*. 2017;**11**(2):109-18.
- Seidm Mahmoodi J, Rahimi C, Mohammadi N. Resiliency and religious orientation: Factors contributing to posttraumatic growth in Iranian subjects. *Iran J Psychiatry*. 2011;**6**(4):145-50. [PubMed: [22952540](https://pubmed.ncbi.nlm.nih.gov/22952540/)]. [PubMed Central: [PMC3395957](https://pubmed.ncbi.nlm.nih.gov/PMC3395957/)].
- Rahmani A, Mohammadian R, Ferguson C, Golizadeh L, Zirak M, Chavoshi H. Posttraumatic growth in Iranian cancer patients. *Indian J Cancer*. 2012;**49**(3):287-92. doi: [10.4103/0019-509X.104489](https://doi.org/10.4103/0019-509X.104489). [PubMed: [23238146](https://pubmed.ncbi.nlm.nih.gov/23238146/)].
- Seyed Mahmoudi S, Rahimi C, Mohammadi N, Hadianfard H. [Increasing post traumatic growth during time and its relationship with coping styles and optimism in patients with positive HIV]. *J Psychol Achiev*. 2010;**17**(2):165-86. Persian.
- Steger MF, Frazier P, Oishi S, Kaler M. The meaning in life questionnaire: Assessing the presence of and search for meaning in life. *J Couns Psychol*. 2006;**53**(1):80-93. doi: [10.1037/0022-0167.53.1.80](https://doi.org/10.1037/0022-0167.53.1.80).
- Jamali F. *The study of the relationship between religious attitudes, sense of life and mental health in Tehran University students [dissertation]*. Tehran, Iran: Alzahra University; 2004.
- Peymanfar A, Akbari A. [Comparing the feeling of loneliness and sense of meaning in elderly life with different levels of religious attitude]. *J Psychol Religion*. 2013;**5**(4):42-54. Persian.
- Heidarzadeh M, Rassouli M, Shahbolaghi F, Majd H, Karam AM, Ghanavati A. The relationship of posttraumatic growth with quality



- of life in cancer patients. *Bull Environ Pharmacol Life Sci*. 2014;**3**(9):98-102.
35. Jansen L, Hoffmeister M, Chang-Claude J, Brenner H, Arndt V. Benefit finding and post-traumatic growth in long-term colorectal cancer survivors: Prevalence, determinants, and associations with quality of life. *Br J Cancer*. 2011;**105**(8):1158-65. doi: [10.1038/bjc.2011.335](https://doi.org/10.1038/bjc.2011.335). [PubMed: [21878935](https://pubmed.ncbi.nlm.nih.gov/21878935/)]. [PubMed Central: [PMC3208486](https://pubmed.ncbi.nlm.nih.gov/PMC3208486/)].
  36. Bellizzi KM, Smith AW, Reeve BB, Alfano CM, Bernstein L, Meeske K, et al. Posttraumatic growth and health-related quality of life in a racially diverse cohort of breast cancer survivors. *J Health Psychol*. 2010;**15**(4):615-26. doi: [10.1177/1359105309356364](https://doi.org/10.1177/1359105309356364). [PubMed: [20460418](https://pubmed.ncbi.nlm.nih.gov/20460418/)].
  37. Brix SA, Bidstrup PE, Christensen J, Rottmann N, Olsen A, Tjonneland A, et al. Post-traumatic growth among elderly women with breast cancer compared to breast cancer-free women. *Acta Oncol*. 2013;**52**(2):345-54. doi: [10.3109/0284186X.2012.744878](https://doi.org/10.3109/0284186X.2012.744878). [PubMed: [23240637](https://pubmed.ncbi.nlm.nih.gov/23240637/)].
  38. Hooper LM, Marotta SA, Depuy V. A confirmatory factor analytic study of the posttraumatic growth inventory among a sample of racially diverse college students. *J Ment Health*. 2009;**18**(4):335-43. doi: [10.1080/09638230802522502](https://doi.org/10.1080/09638230802522502).
  39. Aflakseir A, Nowroozi S, Mollazadeh J, Goodarzi MA. The role of psychological hardiness and marital satisfaction in predicting posttraumatic growth in a sample of women with breast cancer in Isfahan. *Iran J Cancer Prev*. 2016;**9**(4). e4080. doi: [10.17795/ijcp-4080](https://doi.org/10.17795/ijcp-4080). [PubMed: [27761204](https://pubmed.ncbi.nlm.nih.gov/27761204/)]. [PubMed Central: [PMC5055761](https://pubmed.ncbi.nlm.nih.gov/PMC5055761/)].
  40. Teodorescu DS, Siqueland J, Heir T, Hauff E, Wentzel-Larsen T, Lien L. Posttraumatic growth, depressive symptoms, posttraumatic stress symptoms, post-migration stressors and quality of life in multi-traumatized psychiatric outpatients with a refugee background in Norway. *Health Qual Life Outcomes*. 2012;**10**:84. doi: [10.1186/1477-7525-10-84](https://doi.org/10.1186/1477-7525-10-84). [PubMed: [22824521](https://pubmed.ncbi.nlm.nih.gov/22824521/)]. [PubMed Central: [PMC3416737](https://pubmed.ncbi.nlm.nih.gov/PMC3416737/)].
  41. Abedi S, Foroughan M, Khanjani MS, Bakhshi EA, Farhadi A. Relationship between meaning of life and spiritual well-being in the older people residing in nursing homes Shemiranat, 2014. *Salmand*. 2016;**11**(3):456-65. doi: [10.21859/sija-1103456](https://doi.org/10.21859/sija-1103456).
  42. Thompson P. The relationship of fatigue and meaning in life in breast cancer survivors. *Oncol Nurs Forum*. 2007;**34**(3):653-60. doi: [10.1188/07.ONF.653-660](https://doi.org/10.1188/07.ONF.653-660). [PubMed: [17573324](https://pubmed.ncbi.nlm.nih.gov/17573324/)].
  43. Yeganeh Khah M, Abedini A, Akbari H, Ziyayi Nezhad M. *Iran J Nurs*. 2012;**24**(74):36-44.
  44. Kord B, Rahbari P. The prediction of subjective well-being based on meaning of life and mindfulness among cardiovascular patients. *Iran J Psychiatric Nurs*. 2018;**5**(6):16-23. doi: [10.21859/ijpn-05063](https://doi.org/10.21859/ijpn-05063).
  45. Hassankhani H, Soheili A, Hosseinpour I, Eivazi Ziaei J, Nahamin M. A comparative study on the meaning in life of patients with cancer and their family members. *J Caring Sci*. 2017;**6**(4):325-33. doi: [10.15171/jcs.2017.031](https://doi.org/10.15171/jcs.2017.031). [PubMed: [29302572](https://pubmed.ncbi.nlm.nih.gov/29302572/)]. [PubMed Central: [PMC5747591](https://pubmed.ncbi.nlm.nih.gov/PMC5747591/)].
  46. Erci B. Meaning in life of patients with cancer. *Palliat Support Care*. 2015;**13**(1):3-10. doi: [10.1017/S1478951513000254](https://doi.org/10.1017/S1478951513000254). [PubMed: [23759250](https://pubmed.ncbi.nlm.nih.gov/23759250/)].
  47. Dursun P, Saracli S, Konuk O. The roles of meaningful life and posttraumatic stress in posttraumatic growth in a sample of Turkish university students. *Procedia Soc Behav Sci*. 2014;**159**:702-6. doi: [10.1016/j.sbspro.2014.12.454](https://doi.org/10.1016/j.sbspro.2014.12.454).
  48. Garcini L, Short M, Norwood W. Affective and motivational predictors of perceived meaning in life among college students. *J Happiness Well Being*. 2013;**1**(2):47-60.
  49. Shafiee M, Basharpour S, Heydaryrad H. [Comparison of the meaning of life, suicidal thoughts, and cognitive distortions between female patient affected by post-traumatic stress disorder and non-patient]. *Thought Behav Clin Psychol*. 2016;**10**(40):67-76. Persian.
  50. Weiss T, Berger R. *Posttraumatic growth and culturally competent practice: Lessons learned from around the globe*. John Wiley Sons; 2010. doi: [10.1002/9781118270028](https://doi.org/10.1002/9781118270028).