



Comparing the Effect of Walking and Yoga on Clinical and Laboratory Parameters in Men with Type II Diabetes: A Randomized Controlled Clinical Trial

Bayan Saberipour ¹, Mahin Gheibizadeh ^{1,*}, Elham Maraghi ² and Leila Moradi ³

¹Nursing Care Research Center in Chronic Diseases, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

²Department of Biostatistics and Epidemiology, School of Public Health, Ahvaz Jundishapur University of Medical sciences, Ahvaz, Iran

³Diabetes Research Center of Persian Gulf, Health Research Institute, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

*Corresponding author: Nursing Care Research Center in Chronic Diseases, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. Email: gheibizadeh-m@ajums.ac.ir

Received 2019 December 15; Revised 2020 April 08; Accepted 2020 April 09.

Abstract

Background: Identifying safe strategies to regulate the laboratory and clinical parameters of patients with diabetes is critical.

Objectives: The study aimed to compare the effect of walking and yoga on clinical and laboratory indicators in men with type II diabetes.

Methods: This randomized clinical trial was conducted on 108 patients with diabetes who were randomly assigned to walking, yoga, and control groups. Data were collected using a demographic information questionnaire, mercury sphygmomanometer, height, and weight scale, and blood serum biochemical tests before the intervention and after the end of the eighth week of the intervention. Interventions were performed in both yoga and walking groups for 8 weeks, three times a week for 60 minutes each time, and the control group did not receive any intervention. Data were analyzed by SPSS V. 20 software. The significance level was less than 0.05.

Results: The results showed after the intervention the mean of FBS ($P = 0.045$), LDL ($P = 0.005$), HDL ($P = 0.001$), systolic blood pressure ($P = 0.002$), and diastolic blood pressure ($P = 0.003$) were significantly different between the three groups. Changes in the mean of fasting blood sugar (FBS), cholesterol, and body mass index (BMI) were significant only in the yoga group, and changes in the mean of high-density lipoprotein (HDL) were significant solely in the walking group ($P < 0.05$).

Conclusions: Yoga and walking had a desirable effect on improving the clinical and laboratory indicators in men with type II diabetes. However, yoga regulated more indicators in the studied patients in comparison to walking.

Keywords: Walking, Yoga, Blood Sugar, Blood Pressure, Diabetes Mellitus, Body Mass Index

1. Background

Diabetes Mellitus (DM) is a chronic metabolic disorder caused by impaired insulin release, insulin function, or both, which includes 90% of all types of diabetes (1, 2). Diabetes Mellitus has a growing trend in recent years and was increasingly prevalent. According to the World Health Organization (WHO), adults with DM throughout the world were 422 million in 2014, which is expected to reach over 366 million by 2030 (3). In Iran, it is estimated that the number of people with diabetes will reach 9.2 million by the year 2030 (4).

Various factors have been identified as risk factors of DM, including smoking, being overweight and obese, low physical activity, hypertension, high cholesterol, and genetic (5, 6). Moreover, DM has many chronic compli-

cations, including decreased physical function, mental health problems, impaired personal and interpersonal relationships, neuropathy, nephropathy, retinopathy, and cardiovascular disorders (7, 8). Also, high levels of low-density lipoprotein (LDL) and very-low-density lipoprotein (VLDL), and low levels of high-density lipoprotein (HDL) with high blood pressure in patients with DM make them more susceptible to atherosclerosis, which is the most common cause of death in these patients type II DM (8).

Exercise is one of the therapeutic options for patients with DM, which lowers blood glucose, glycosylated hemoglobin, and cardiovascular risk factors, and prevents long-term complications of DM. Walking is one of the activities that has a positive effect on diabetes and leads to increased norepinephrine, serotonin synthesis, and endorphins, which are effective in causing vitality and prevent-

ing fatigue in people with diabetes (9, 10). Walking is a simple sport that does not require any special equipment and can be combined in the daily life of people with DM (11, 12). In this regard, the study by Yekkeh Fallah et al. showed that walking is effective in lowering blood sugar in patients with DM (13).

Yoga is another exercise that enhances muscle flexibility and strength, and also improves blood circulation, oxygen uptake, and hormone function. In addition, yoga stabilizes the autonomic nervous system, controls emotions, and improves one's sense of health (13, 14). Various studies have investigated the effects of yoga that have reported different results (13, 15, 16). For example, the study of Yekkeh Fallah et al. showed that decrease in fasting blood sugar (FBS) at the end of the intervention in the yoga group was significant (13), but Sreedevi et al. (15) and Shiju et al. (16) did not report significant changes in FBS and lipid profile.

Although the effects of walking and yoga on improving some laboratory and clinical indicators have been identified separately, there is not enough knowledge about the effectiveness of these two methods compared to each other. Therefore, it is necessary to compare these two activities so that we can recommend the most appropriate method to patients. Therefore, this study aimed to compare the effect of walking and yoga on the clinical and laboratory parameters in men with type II DM.

2. Objectives

The aim of this study was to compare the effect of walking and yoga on clinical and laboratory indicators in men with type II DM.

3. Methods

This randomized clinical trial (IRCT20180511039610N1) was conducted with a pre-test/post-test design on 108 men with type II DM referring to Golestan and Imam Khomeini and hospitals in Ahvaz, Iran, from May to August 2019. The sample size was calculated as 108 individuals using table B.12 (17) and considering 10% attrition and $\alpha = 0.05$, $\beta = 0.95$, and $\Delta/\sigma \sim 1$ according to the previous studies (9).

The inclusion criteria were men aged 30 to 65 years, at least 1 year has been passed since the diagnosis of DM, not participating in another exercise programs at the same time, lack of exercise prohibitions, lack of any physical activity limitations, lack of insulin treatment, and not taking antidepressants. The exclusion criteria were hospitalization during the intervention, change of treatment from oral medications to insulin therapy during the study period, migration to another city, and medical forbiddance

for physical activity during the intervention. One hundred and eight eligible men were randomly assigned to walking ($n = 36$), yoga ($n = 36$), and control ($n = 36$) groups using a block permutation method with block size 6. The random allocation was made by a statistical consultant who was a member of the research team (Figure 1).

After the random allocation of the patients to the groups, the pre-test was conducted in all three groups. The walking group was asked to walk for 8 weeks three times a week for 60 minutes each time. The yoga group also received yoga training for 8 weeks, 3 sessions a week, and 60 minutes each time, under the supervision of the researcher and professional instructor in a designated gym. Yoga included warm-up, physical and breathing exercises, relaxation, and meditation. The control group did not receive any intervention. A post-test was conducted on the three groups at the end of the eighth week of interventions. Data collection tools were a demographic information questionnaire (containing questions about age, occupation, educational level of the patient and his wife, economic status, history of hospitalization), mercury sphygmomanometer (Yamasa Model 600), and height and weight scale (Seka model 786). The same tools were used for all participants, and the tools were calibrated before use. Simultaneous measurements of height, weight, and blood pressure were performed by two people, including the researcher and a research assistant nurse. To perform biochemical tests, a blood sample of 10 ml was taken in two stages (pre-test and post-test) in the fasting state by the researcher. All blood samples were evaluated using quantitative diagnostic kits (fasting blood sugar: FBS, Cholesterol, Triglycerides: TG, HDL-C, and LDL-C) of Pars Azmun Company after storage at -70°C . All participants' blood tests were evaluated in the same laboratory.

Data analysis was performed using SPSS software version 20. To examine the relationship between qualitative variables, the chi-square test was used and to compare quantitative variables between three independent groups (yoga, walking, and control), one-way analysis of variance, or its non-parametric equivalent (Kruskal-Wallis test) was used. The Mann-Whitney test was used to determine the mean difference between the independent groups, and the Wilcoxon test was used to compare the mean within the groups. The significance level was considered to be less than 0.05.

3.1. Ethical Considerations

This study was approved by the Research Ethics Committee of Ahvaz Jundishapur University of Medical Sciences (IR.AJUMS.REC.1397.101). Ethical considerations included explaining essential information to participants, obtaining informed consent, voluntary participation in




Figure 1. CONSORT flowchart

the study, freedom in leaving the study at any time, the confidentiality of the information, and no imposing costs on the participants.

4. Results

In this study 108 men with type II DM participated. Their mean (standard deviation: SD) age was 49.91 (9.42)

years. The mean (SD) of participants' height was 170.99 (7.02) cm, weight was 81.82 (9.73) kg, Body Mass Index (BMI) was 28.06 (3.31) kg/m², and systolic blood pressure (SBP) and diastolic blood pressure (DBP) were 135.22 (13.71) and 83.48 (8.80) mmHg, respectively. There was no statistically significant difference between the three groups in terms of demographic characteristics (Table 1).

The results showed that there was no difference be-

Table 1. Homogeneity of Demographic Variables in Intervention and Control Groups^a

Variables	Groups			P Value
	Yoga	Walking	Control	
Occupation^b				0.126
Employee in governmental sections	2 (5.6)	5 (13.9)	6 (16.7)	
Self-employed	24 (66.7)	16 (44.4)	23 (63.9)	
Unemployed	5 (13.9)	3 (8.3)	3 (8.3)	
Other	5 (13.9)	12 (33.3)	4 (11.1)	
Education^b				0.518
Illiterate	1 (2.8)	3 (8.3)	1 (2.8)	
Primary school	15 (41.7)	11 (30.6)	11 (30.6)	
High school diploma or associate degree	18 (50)	17 (47.2)	22 (61.1)	
Bachelor's degree and above	2 (5.6)	5 (13.9)	2 (5.6)	
Financial status^b				0.249
Good	2 (5.6)	4 (11.1)	4 (11.1)	
Average	29 (80.6)	20 (55.6)	23 (63.9)	
Weak	5 (13.9)	12 (33.3)	9 (25)	
History of hospitalization^b				0.510
Yes	12 (33.3)	15 (41.7)	11 (30.6)	
No	24 (66.7)	21 (58.3)	25 (69.4)	
Education of spouse^b				0.634
Illiterate	3 (8.3)	4 (11.1)	5 (13.9)	
Primary school	14 (38.9)	21 (58.3)	16 (44.4)	
High school diploma or associate degree	18 (50)	10 (27.8)	14 (38.9)	
Bachelor's degree and above	1 (2.8)	1 (2.8)	1 (2.8)	
Age^c	48.25 ± 7.14	49.83 ± 9.58	51.66 ± 11.06	0.308

^aValues are expressed as No. (%) or mean ± SD.

^bChi-square test.

^cOne-way ANOVA.

tween the mean scores of the clinical and laboratory parameters between the three groups before the intervention ($P > 0.05$). However, data analysis of post-test showed significant differences in the mean scores of FBS ($P = 0.045$), LDL ($P = 0.005$), HDL ($P = 0.001$), SBP ($P = 0.002$), DBP ($P = 0.003$), and BMI ($P = 0.024$) between the three groups after the intervention. There were no significant differences between the three groups in terms of TG ($P = 0.163$) and cholesterol levels ($P = 0.061$) (Table 2).

The comparison of groups using Post hoc test showed that there were statistically significant differences after intervention between the two groups of walking and control in terms of FBS level ($P = 0.028$); between the two groups of yoga and control ($P = 0.007$) and walking and control groups ($P = 0.033$) in terms of LDL level; between the two groups of yoga and control ($P = 0.001$), and the two groups

of walking and control ($P = 0.016$) in terms of HDL level. Moreover, significant differences were seen between the two groups of yoga and control ($P = 0.004$) and the two groups of walking and control ($P = 0.001$) in SBP; and between yoga and control ($P = 0.02$) and walking and control ($P = 0.005$) groups in DBP. Regarding BMI, there was a significant difference between the two groups of walking and control ($P = 0.027$).

5. Discussion

In this study that was carried out with the aim of comparing the effects of walking and yoga on the clinical and laboratory parameters in men with type II DM, the findings revealed that walking and yoga training had an effective role in improving clinical and laboratory parameters

Table 2. Comparison of Laboratory and Clinical Parameters in Men with Type II DM in the Three Groups (Yoga, Walking, and Control) Before and After Intervention^a

Variables	Before Intervention	After Intervention	P Value ^b
FBS (mg/dL)			
Yoga	181.11 (56.48)	159.62 (49.40)	< 0.0001
Walking	162.66 (59.89)	165.30 (64.81)	0.668
Control	179.05 (75.03)	195.57 (63.37)	0.022
P value ^c	0.232	0.045	
LDL (mg/dL)			
Yoga	80.97 (30.07)	79.34 (25.83)	0.724
Walking	84.67 (35.79)	83.24 (29.10)	0.872
Control	88.91 (28.66)	101.81 (29.17)	0.001
P value ^c	0.478	0.005	
HDL (mg/dL)			
Yoga	46.25 (9.02)	48.71 (7.53)	0.107
Walking	45.52 (11.16)	48.49 (10.91)	0.022
Control	42.30 (7.72)	41.96 (7.07)	0.274
P value ^c	0.150	0.001	
TG (mg/dL)			
Yoga	151.02 (87.07)	139.68 (71.81)	0.166
Walking	161.11 (93.75)	149.45 (99.06)	0.245
Control	161.11 (91.64)	171.21 (82.63)	0.391
P value ^c	0.809	0.163	
Cholesterol (mg/dL)			
Yoga	154.61 (42.04)	145.65 (38.37)	0.003
Walking	151.40 (42.14)	149.09 (35.28)	0.681
Control	163.52 (31.31)	166.15 (36.12)	0.754
P value ^c	0.426	0.061	
SBP (mmHg)			
Yoga	134.44 (13.61)	127.34 (11.07)	0.001
Walking	132.05 (13.53)	126.36 (11.94)	< 0.0001
Control	139.16 (13.38)	136.66 (12.16)	0.059
P value ^c	0.062	0.002	
DBP (mmHg)			
Yoga	82.08 (8.81)	82.03 (6.93)	0.824
Walking	82.38 (9.54)	81.66 (7.03)	0.514
Control	85.97 (7.63)	86.51 (5.22)	0.533
P value ^c	0.222	0.003	
BMI (kg/m²)			
Yoga	28.45 (3.06)	27.88 (2.77)	0.001
Walking	27.01 (2.92)	26.69 (3.03)	0.07
Control	28.72 (3.74)	28.82 (3.68)	0.761
P value ^c	0.058	0.024	

Abbreviations: BMI, body mass index; DBP, diastolic blood pressure; SBP, systolic blood pressure; TG, triglycerides.

^aStatistically significant (P < 0.05).

^bWilcoxon test.

^cMann-Whitney U test.

in men with type II DM.

Accordingly, the results indicated that there was a significant difference in mean of FBS, LDL, HDL, SBP, and DBP,

and BMI indices between the three groups of yoga, walking, and control after the intervention, but cholesterol and TG levels showed no significant differences in the three groups. The yoga group showed a greater decrease in FBS, cholesterol, and BMI compared to the walking group. However, the decrease in SBP was more in the walking group than that in the yoga group. The inter-group comparisons showed a statistically significant difference in the mean of BMI in the two groups of walking and yoga, which was in favor of the yoga group, but there was no significant difference between the two groups in other indices. Consistent with the present study, the study by Sreedevi et al., which aimed to determine the effect of yoga and peer support on blood sugar (BS) in women with type II DM, showed that the mean of SBP decreased significantly after yoga intervention (15). However, the results of Kim et al. did not show such effectiveness (18). The difference between the results of the researches can be attributed to the difference between the intensity and duration of the training programs as well as the difference between the age and gender of the research samples. In a study by Rafiei et al., with the purpose of comparing the effect of Tai Chi exercise and walking on FBS in patients with type II DM, there was a significant difference between the mean of BS levels before and after the intervention in the Tai Chi and walking groups. Although FBS in the walking group had more decline than in Tai Chi and control groups, the difference was not significant (19). Yekkeh Fallah et al. in their study which aimed to compare the effect of one month of regular yoga and walking exercises on BS and the number of pills used in type II DM patients, declared that there was a significant decrease in FBS at the end of the intervention in the yoga group, but it was not significant in the walking group (13). In a study by Bello-Ovosi et al., who aimed to investigate the effects of aerobic exercise on physiological parameters and quality of life in patients with type II DM, the results showed that FBS, LDL, and HDL were significantly improved significantly in the intervention group (20). Chimkode et al., in a study with the purpose of investigating the effect of yoga on blood glucose levels in healthy and diabetic subjects, reported that yoga practices led to a significant reduction in the mean of fasting and after-breakfast BS levels in the diabetic group compared to the healthy group (2). However, some studies have shown no significant changes in clinical and laboratory parameters of patients after exercise interventions (15, 16). For example, in the study by Sreedevi, cholesterol level, FBS, and BMI did not change significantly after the intervention (15). Also, the results of a study by Shiju et al. showed that yoga intervention did not change or improve the BS level and lipid profile of studied patients (16). In the study by Dabbagh Nikukheslat et al. (21), BMI did not show any significant change after the

intervention. Shantakumari et al. reported that after performing yoga exercises, although BMI decreased and HDL levels increased in type 2 DM patients, it was not statistically significant. They stated that yoga exercises significantly reduced cholesterol, TG, and LDL levels in the intervention group (22). It seems that practicing yoga exercises and regular walking in proper time intervals can improve some of the laboratory and clinical parameters. Based on the results of various studies as well as the present study, what is understood is that it is not possible to say with certainty which of the two methods of walking and yoga is superior to the other. So patients can take advantage of any of these exercises that they find more enjoyable.

One of the limitations of this study, which limited decision-making to more effective intervention, was the failure to measure glycosylated hemoglobin (HgA1C) due to time constraints. Another limitation of the present study was that it was not possible to observe the exact course of walking and its severity in patients in the walking group, so the self-report method was used to evaluate it, and patients were monitored by telephone. Therefore, further studies are recommended with careful control of activity intensity, as well as by measuring HgA1C to determine the effectiveness of the intervention in controlling BS over time.

5.1. Conclusions

The findings of the study show that yoga and walking exercises improve the FBS level and lipid profile, especially cholesterol and HDL levels in men with type II DM. Yoga and walking exercises are also effective in reducing SBP and BMI. However, in comparison to walking, yoga had a more desirable effect on lowering the levels of FBS and cholesterol in the patients. Therefore, considering the effectiveness of yoga and walking exercises, it is recommended to consider these exercises in the training and self-care programs of patients with diabetes. It is likely that the duration of training in these patients should be longer for further influence on other parameters (TG and LDL).

Acknowledgments

This study is part of the M.Sc. thesis of the first author (Bayan Saberipour), which was financially supported by the Nursing Care Research Center in Chronic Diseases of Ahvaz Jundishapur University of Medical Sciences (NCRCCD-9708). The authors would like to express their appreciation to the sponsor of the study as well as all patients who participated in our study.

Footnotes

Authors' Contribution: Bayan Saberipour, Mahin Gheibizadeh, Elham Maraghi, and Leila Moradi were responsible for the study conception and design, data analysis, and drafting of the manuscript. Data collection was conducted by Bayan Saberipour. Elham Maraghi was responsible for statistical analysis.

Clinical Trial Registration Code: IRCT20180511039610N1.

Conflict of Interests: The authors declare no conflict of interest.

Ethical Approval: This article is the result of research approved by the Ethics Committee of Ahvaz Jundishapur University of Medical Sciences (IR.AJUMS.REC.1397.101).

Funding/Support: This study was financially supported by Nursing Care Research Center in Chronic Diseases of Ahvaz Jundishapur University of Medical Sciences (NCRCCD-9708).

Patient Consent: All research ethics considerations, including explaining essential information to participants, obtaining informed consent, voluntary participation right, freedom in leaving the study at any time of the research, the confidentiality of the information and the non-imposition of costs on the participants were considered.

References

1. Al-Kuraishy HM, Al-Gareeb AI. Erectile Dysfunction and Low Sex Drive in Men with Type 2 DM: The Potential Role of Diabetic Pharmacotherapy. *J Clin Diagn Res.* 2016;**10**(12):FC21-6. doi: [10.7860/JCDR/2016/19971.8996](https://doi.org/10.7860/JCDR/2016/19971.8996). [PubMed: [28208875](https://pubmed.ncbi.nlm.nih.gov/28208875/)]. [PubMed Central: [PMC5296448](https://pubmed.ncbi.nlm.nih.gov/PMC5296448/)].
2. Chimkode SM, Kumaran SD, Kanhere VV, Shivanna R. Effect of yoga on blood glucose levels in patients with type 2 diabetes mellitus. *J Clin Diagn Res.* 2015;**9**(4):CC01-3. doi: [10.7860/JCDR/2015/12666.5744](https://doi.org/10.7860/JCDR/2015/12666.5744). [PubMed: [26023550](https://pubmed.ncbi.nlm.nih.gov/26023550/)]. [PubMed Central: [PMC4437062](https://pubmed.ncbi.nlm.nih.gov/PMC4437062/)].
3. Ebrahimi M, Guilan-Nejad TN, Pordanjani AF. Effect of yoga and aerobics exercise on sleep quality in women with Type 2 diabetes: a randomized controlled trial. *Sleep Sci.* 2017;**10**(2):68-72. doi: [10.5935/1984-0063.20170012](https://doi.org/10.5935/1984-0063.20170012). [PubMed: [28966742](https://pubmed.ncbi.nlm.nih.gov/28966742/)]. [PubMed Central: [PMC5612039](https://pubmed.ncbi.nlm.nih.gov/PMC5612039/)].
4. Safari-Faramani R, Rajati F, Tavakol K, Hamzeh B, Pasdar Y, Moradinazar M, et al. Prevalence, Awareness, Treatment, Control, and the Associated Factors of Diabetes in an Iranian Kurdish Population. *J Diabetes Res.* 2019;**2019**:5869206. doi: [10.1155/2019/5869206](https://doi.org/10.1155/2019/5869206). [PubMed: [31565657](https://pubmed.ncbi.nlm.nih.gov/31565657/)]. [PubMed Central: [PMC6745166](https://pubmed.ncbi.nlm.nih.gov/PMC6745166/)].
5. McDermott KA, Rao MR, Nagarathna R, Murphy EJ, Burke A, Nagendra RH, et al. A yoga intervention for type 2 diabetes risk reduction: a pilot randomized controlled trial. *BMC Complement Altern Med.* 2014;**14**:212. doi: [10.1186/1472-6882-14-212](https://doi.org/10.1186/1472-6882-14-212). [PubMed: [24980650](https://pubmed.ncbi.nlm.nih.gov/24980650/)]. [PubMed Central: [PMC4096416](https://pubmed.ncbi.nlm.nih.gov/PMC4096416/)].
6. Valizadeh R, Vali L, Bahaadinbeigy K, Amiresmaili M. The Challenges of Iran's Type 2 Diabetes Prevention and Control Program. *Int J Prev Med.* 2019;**10**:175. doi: [10.4103/ijpvm.IJPVM_371_17](https://doi.org/10.4103/ijpvm.IJPVM_371_17). [PubMed: [32133093](https://pubmed.ncbi.nlm.nih.gov/32133093/)]. [PubMed Central: [PMC6826765](https://pubmed.ncbi.nlm.nih.gov/PMC6826765/)].
7. Defeudis G, Gianfrilli D, Di Emidio C, Pofi R, Tuccinardi D, Palermo A, et al. Erectile dysfunction and its management in patients with diabetes

- mellitus. *Rev Endocr Metab Disord*. 2015. doi: [10.1007/s11154-015-9321-4](https://doi.org/10.1007/s11154-015-9321-4). [PubMed: [26497842](https://pubmed.ncbi.nlm.nih.gov/26497842/)].
8. Stoa EM, Meling S, Nyhus LK, Glenn S, Mangerud KM, Helgerud J, et al. High-intensity aerobic interval training improves aerobic fitness and HbA1c among persons diagnosed with type 2 diabetes. *Eur J Appl Physiol*. 2017;**117**(3):455-67. doi: [10.1007/s00421-017-3540-1](https://doi.org/10.1007/s00421-017-3540-1). [PubMed: [28160083](https://pubmed.ncbi.nlm.nih.gov/28160083/)].
 9. Azimi H, Yekkehfallah L, Ghorbani A. [Comparing Effects Of Walking And Yoga Exercise On Quality Of Life In Patients With Type 2 Diabetes]. *Preventive Care In Nursing & Midwifery Journal*. 2015;**5**(1):46-58. Persian.
 10. Ezema CI, Omeh E, Onyeso OKK, Anyachukwu CC, Nwankwo MJ, Amaeze A, et al. The Effect of an Aerobic Exercise Programme on Blood Glucose Level, Cardiovascular Parameters, Peripheral Oxygen Saturation, and Body Mass Index among Southern Nigerians with Type 2 Diabetes Mellitus, Undergoing Concurrent Sulfonylurea and Metformin Treatment. *Malays J Med Sci*. 2019;**26**(5):88-97. doi: [10.21315/mjms2019.26.5.8](https://doi.org/10.21315/mjms2019.26.5.8). [PubMed: [31728121](https://pubmed.ncbi.nlm.nih.gov/31728121/)]. [PubMed Central: [PMC6839658](https://pubmed.ncbi.nlm.nih.gov/PMC6839658/)].
 11. Algeffari M, Jayasena CN, MacKeith P, Thapar A, Dhillo WS, Oliver N. Testosterone therapy for sexual dysfunction in men with Type 2 diabetes: a systematic review and meta-analysis of randomized controlled trials. *Diabet Med*. 2018;**35**(2):195-202. doi: [10.1111/dme.13553](https://doi.org/10.1111/dme.13553). [PubMed: [29171080](https://pubmed.ncbi.nlm.nih.gov/29171080/)].
 12. Kim JM, Kim MK, Joung KH, Lee JH, Kim HJ, Ku BJ. Association between glycemic state and pulmonary function and effect of walking as a protective factor in subjects with diabetes mellitus. *Ann Transl Med*. 2019;**7**(20):530. doi: [10.21037/atm.2019.09.139](https://doi.org/10.21037/atm.2019.09.139). [PubMed: [31807512](https://pubmed.ncbi.nlm.nih.gov/31807512/)]. [PubMed Central: [PMC6861773](https://pubmed.ncbi.nlm.nih.gov/PMC6861773/)].
 13. Yekkeh Fallah L, Azimi H, Sadeghi T. [Comparison of the Effects of Two Types of Exercise on Blood Glucose Levels and Anti-diabetics Tablets Used by People with Type II Diabetes]. *Iran Journal of Nursing*. 2014;**27**(87):79-87. Persian. doi: [10.29252/ijn.27.87.79](https://doi.org/10.29252/ijn.27.87.79).
 14. Rendell MS, Rajfer J, Wicker PA, Smith MD. Sildenafil for treatment of erectile dysfunction in men with diabetes: a randomized controlled trial. Sildenafil Diabetes Study Group. *JAMA*. 1999;**281**(5):421-6. doi: [10.1001/jama.281.5.421](https://doi.org/10.1001/jama.281.5.421). [PubMed: [9952201](https://pubmed.ncbi.nlm.nih.gov/9952201/)].
 15. Sreedevi A, Gopalakrishnan UA, Karimassery Ramaiyer S, Kamalamma L. A Randomized controlled trial of the effect of yoga and peer support on glycaemic outcomes in women with type 2 diabetes mellitus: a feasibility study. *BMC Complement Altern Med*. 2017;**17**(1):100. doi: [10.1186/s12906-017-1574-x](https://doi.org/10.1186/s12906-017-1574-x). [PubMed: [28173786](https://pubmed.ncbi.nlm.nih.gov/28173786/)]. [PubMed Central: [PMC5297169](https://pubmed.ncbi.nlm.nih.gov/PMC5297169/)].
 16. Shiju R, Thomas D, Al Arouj M, Sharma P, Tuomilehto J, Bennakhi A. Effect of Sudarshan Kriya Yoga on anxiety, depression, and quality of life in people with type 2 diabetes: A pilot study in Kuwait. *Diabetes Metab Syndr*. 2019;**13**(3):1995-9. doi: [10.1016/j.dsx.2019.04.038](https://doi.org/10.1016/j.dsx.2019.04.038). [PubMed: [31235126](https://pubmed.ncbi.nlm.nih.gov/31235126/)].
 17. Neter J Km, Nachtsheim CJ, Wasserman W. *Applied Linear Statistical Models*. Chicago: Irwin; 1996.
 18. Kim HN, Ryu J, Kim KS, Song SW. Effects of yoga on sexual function in women with metabolic syndrome: a randomized controlled trial. *J Sex Med*. 2013;**10**(11):2741-51. doi: [10.1111/jsm.12283](https://doi.org/10.1111/jsm.12283). [PubMed: [23899008](https://pubmed.ncbi.nlm.nih.gov/23899008/)].
 19. Rafiei F, Masroor D, Haghani H, Azimi H. [The comprison of exercise as a copplementary therapy and walking exercise on the fasting blood glucose of people with type 2 diabetes]. *J Urmia Nurs Midwifery Fac*. 2017;**15**(3):218-29. Persian.
 20. Bello-Ovosi BO, Asuke S, Abdulrahman SO, Ibrahim MS, Ovosi JO, Ogunsina MA, et al. Prevalence and correlates of hypertension and diabetes mellitus in an urban community in North-Western Nigeria. *Pan Afr Med J*. 2018;**29**:97. doi: [10.11604/pamj.2018.29.97.14191](https://doi.org/10.11604/pamj.2018.29.97.14191). [PubMed: [29875978](https://pubmed.ncbi.nlm.nih.gov/29875978/)]. [PubMed Central: [PMC5987122](https://pubmed.ncbi.nlm.nih.gov/PMC5987122/)].
 21. Dabbagh Nikukheslat S, Gholami F, Salek-Zamani Y, Jafari A, Faraji E. [Effect of 12-Week Aerobics Training on Metabolic Control in Type-2 Diabetic Men with Peripheral Neuropathy]. *Med J Tabriz Uni Med Sciences Health Services*. 2018;**39**(6):21-8. Persian.
 22. Shantakumari N, Sequeira S, El deeb R. Effects of a yoga intervention on lipid profiles of diabetes patients with dyslipidemia. *Indian Heart J*. 2013;**65**(2):127-31. doi: [10.1016/j.ihj.2013.02.010](https://doi.org/10.1016/j.ihj.2013.02.010). [PubMed: [23647889](https://pubmed.ncbi.nlm.nih.gov/23647889/)]. [PubMed Central: [PMC3861018](https://pubmed.ncbi.nlm.nih.gov/PMC3861018/)].