

# Characterization of the Extended-Spectrum beta-Lactamase Producers among Non-Fermenting Gram-Negative Bacteria Isolated from Burnt Patients

Mojdeh Hakemi Vala<sup>a</sup>, Masoumeh Hallajzadeh<sup>a</sup>, Fatemeh Fallah<sup>b,a\*</sup>,  
Ali Hashemi<sup>a</sup>, Hossein Goudarzi<sup>a</sup>

<sup>a</sup> Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

<sup>b</sup> Pediatric Infections Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

\*Correspondence should be addressed to Dr. Fatemeh Fallah; Email: [fallah@pirc.ir](mailto:fallah@pirc.ir)

## A-R-T-I-C-L-E I-N-F-O

### Article Notes:

Received: Dec 1, 2012

Received in revised form: Jan 15, 2013

Accepted: Jan 16, 2013

Available Online: Jan 20, 2013

### Keywords:

*Acinetobacter baumannii*

beta-Lactamases

Drug Resistance

Iran

Patients, Burnt

*Pseudomonas aeruginosa*

## A-B-S-T-R-A-C-T

**Background & Aims of the Study:** Extended-spectrum beta-Lactamases (ESBLs) represent a major group of beta-lactamases which are responsible for resistance to oxyimino-cephalosporins and aztreonam and currently being identified in large numbers throughout the world. The objective of this study was to characterize ESBL producers among non-fermenter gram-negative bacteria isolated from burnt patients.

**Materials & Methods:** During April to July 2012, 75 non-fermenter gram-negative bacilli were isolated from 240 bacterial cultures collected from wounds of burnt patients admitted to the Burn Unit at Shahid Motahari Hospital (Tehran, Iran). Bacterial isolation and identification was done using standard methods. Antimicrobial susceptibility testing was performed by disk diffusion method for all strains against selected antibiotics and minimum inhibitory concentration was determined by microdilution test. The ability to produce ESBL was detected through double disk synergy test among candidate strains.

**Results:** Of 75 non-fermenter isolates, 47 *Pseudomonas aeruginosa* and 28 *Acinetobacter baumannii* were identified. The resistance of *P. aeruginosa* isolates to tested antibiotics in antibiogram test were 100% to cefpodoxime, 82.98% to ceftriaxone, 78.73% to imipenem, 75% to meropenem, 72.72% to gentamicin, 69.23% to ciprofloxacin and aztreonam, 67.57% to ceftazidime, 65.95% to ceftazidime, and 61.53% to piperacillin. The results for *Acinetobacter baumannii* were 100% to ceftazidime, cefepime, ciprofloxacin, imipenem, meropenem, cefpodoxime, and cefotaxim, 96.85% to gentamicin, 89.65% to ceftriaxone, 65.51% to aztreonam, and 40% to piperacillin. Double disk synergy test showed that 21 (28%) of non-fermenter isolates were ESBL producer.

**Conclusions:** None of the third or fourth generation of cephalosporins is suitable for treatment of *Pseudomonas aeruginosa* and *Acinetobacter baumannii* isolated from burnt patients.

**Please cite this article as:** Hakemi Vala M, Hallajzadeh M, Fallah F, Hashemi A, Goudarzi H. Characterization of the extended-spectrum beta-lactamase producers among non-fermenting gram-negative bacteria isolated from burnt patients. Arch Hyg Sci 2013;2(1):1-6.

## Background

Non-fermenting gram-negative bacteria are a group of bacteria which are unable to produce energy by fermenting sugars. Because of

emerging multidrug resistant (MDR) strains, the members of this group have become a particular challenge in healthcare management. Most of the time, gram-negative non-fermenters are pathogens that cause opportunistic

infections in patients who are critically ill or immunocompromised (1).

Non-fermenting bacteria are distributed in nature, especially in soil and water. In the hospital environment, they may be isolated from humidifiers, ventilator machines, mattresses, and other equipment, as well as from the skin of healthcare workers (1). Therefore, recent clinical attention has focused on the increasing frequency of non-fermenting gram-negative pathogens, which account for hospital-acquired infections (2).

Among non-fermenting gram-negative rods, the most clinically important species are *Pseudomonas aeruginosa*, *Stenotrophomonas maltophilia*, and *Acinetobacter baumannii*, which are frequently MDR (3). Infections with *Pseudomonas aeruginosa* and *Acinetobacter baumannii* are of particular concern for hospitalized patients, especially whom with MDR infection (4).

Multidrug resistant *Pseudomonas aeruginosa* and *Acinetobacter baumannii* isolates are increasingly causing hospital-acquired infections. These clones are spreading into new geographic areas, and causing susceptible strains to acquire resistance genes. New extended-spectrum beta-lactamase and carbapenemase producers are emerging and leading to the evolution of pan-resistant strains.

Extended-spectrum beta-lactamases (ESBLs) represent a major group of beta-lactamases responsible for resistance to oxyimino-cephalosporins and aztreonam, and currently being identified in large numbers throughout the world (5). ESBLs are encoded by both chromosomal or plasmid genes (4-7). Hence, acquired resistance to beta-lactams can result in therapeutic failure, particularly when it is associated with resistance to other classes of drugs, such as aminoglycosides and fluoroquinolones (8).

The risk of infection in burns is well-known, and in the recent decades, the frequency of antimicrobial resistance is increasing among isolated bacteria from burnt patients (9). Burn

wound is an ideal environment for bacterial growth, and are potentially richer sources of infection than surgical wounds, predominantly due to the larger area involved and the longer duration of hospitalization (10).

**Aims of the study:** The objective of this study was to characterize extended-spectrum beta-lactamase (ESBL) producers among non-fermenter gram-negative bacteria isolated from burnt patients.

## Materials & Methods

**Bacterial isolates:** During April to July 2012, 75 non-fermenter gram-negative bacilli were isolated from 240 bacterial cultures collected from wounds of burnt patients admitted to the Burn Unit at Shahid Motahari Hospital (Tehran, Iran). The wound exudates were collected by swabbing, and immediately transported in transport culture media under standard conditions to the laboratory of the Department of Microbiology of Shahid Beheshti University of Medical Sciences (Tehran, Iran).

Bacterial isolation and identification was done using standard methods such as "oxidase", "triple sugar iron", "oxidative-fermentative", and "motility" tests (11). Based on the oxidase test, the isolated bacteria were divided into two groups (oxidase-positive and oxidase-negative). Oxidase-positive strains were considered as members of *Pseudomonas* family, and further confirmation tests were performed, including growth on ceftrimid agar, growth at 42°C, decarboxylation of lysine on lysine decarboxylase and arginine dihydrolase test. Oxidase-negative isolates were presumed as members of *Acinetobacter* family, and were confirmed by Microgen identification kit (Microgen<sup>TM</sup>, UK).

**Antibiotic susceptibility (antibiogram) test:** Based on Clinical Laboratory Standards Institute (CLSI) 2012 protocol (12), antimicrobial susceptibility testing was

Piperacillin (100 µg), meropenem (10 µg), imipenem (10 µg), ceftazidime (30 µg), cefotaxime (30 µg), cefepime (30 µg), cefttriaxone (30 µg), cefpodoxime (10 µg), aztreonam (30 µg), ciprofloxacin (5 µg), and gentamicin (10 µg), all purchased from MAST Co. (Mast Diagnostics, UK).

**ESBL screening:** The ability to produce ESBL was detected by double disk synergy test (DDST) among candidate strains.

The ceftazidime resistant strains were candidates for ESBL detection by DDST method. In this test, colonies from overnight cultures on blood agar plates were suspended in Mueller-Hinton broth and their turbidity was adjusted to 0.5 McFarland standard. Then, the suspension was streaked onto Mueller-Hinton agar plates (Qlab, Montreal, Canada) (10). DDST test was performed by using plain ceftazidime disk (30 µg) and ceftazidime/clavulanic acid (30 µg/10 µg) and putting other third generation cephalosporins like cefotaxime, ceftriaxone, ceftazidime, and

**Data analysis:** The results were reported as percentages.

Totally, 75 non-fermenter gram-negative bacilli were isolated from 240 bacterial cultures collected from wounds of burnt patients admitted to the Burn Unit at Shahid Motahari Hospital (Tehran, Iran).

Out of 75 non-fermenter isolates, 47 (62.67%) *Pseudomonas aeruginosa* and 28 (36.33%) *Acinetobacter baumannii* were identified. Antimicrobial resistance data are presented in table 1.

Using DDST, an increase in inhibition zone diameter of 5 mm for ceftazidime/clavulanic acid versus ceftazidime alone were detected among 21 (28%) non-fermenter isolates (table 2).

The MICs of ceftazidime for *Pseudomonas aeruginosa* and *Acinetobacter baumannii* are presented in table 3.

Antibiotics	imipenem	meropenem	ceftazidim	cefotaxime	ceftriaxone	cefepodoxim	cefepime	piperacilin	ciprofloxacin	aztreonam	gentamycin
<i>Pseudomonas aeruginosa</i>	78.73	75	65.95	82.97	82.98	100	67.57	61.53	69.23	69.23	72.72
<i>Acinetobacter baumannii</i>	100	100	100	100	89.65	100	100	40	100	65.51	96.85

Species	Number	extended-spectrum beta-lactamase -positive (%)
<i>Pseudomonas aeruginosa</i>	15	31.91
<i>Acinetobacter baumannii</i>	6	21.42

Species	MIC50 (µg/mL)	MIC90 (µg/mL)	MIC range (µg/mL)
<i>Pseudomonas aeruginosa</i>	128	256	4>X≥256
<i>Acinetobacter baumannii</i>	256	>256	4>X>256

## Discussion

In this study, existence of 100% resistance to 7 tested antibiotics, including imipenem, meropenem, ceftazidime, cefepime, cefpodoxime, cefotaxim, and ciprofloxacin among *Acinetobacter baumannii* shows that these groups of bacteria are more resistant to common antibiotics than *Pseudomonas aeruginosa* isolates. Therefore, based on the antibiotic resistance pattern, none of antibiotics tested in antibiogram test are effective against *Pseudomonas aeruginosa* or *Acinetobacter baumannii* (table 1).

Detection of ESBL producers among gram-negative bacilli is necessary for successful therapy, improving clinical outcomes, and limiting the spread of these MDR organisms (14). ESBL producing strains are usually found in hospitals where antibiotic usage is high and patients are in critical conditions (5). Importance of prolonged duration of staying in burn ward is for its serious threat of the spread of MDR bacterial pathogens, and consequently causing nosocomial infection (10).

31.91% of *Pseudomonas aeruginosa* isolates were ESBL-positive, which is in accordance with other studies conducted in Iran (28%), Bangladesh (37.8%), India (32.6%), and Turkey (31.5%) (5,15-17).

However, it is in contrast with two other studies in Iran that their results were lower (18%) and higher (42.8%) than our result. This discrepancy may be related to more usage of beta-lactam drugs and time of study (18,19).

In addition, 21.42% of *Acinetobacter baumannii* were identified as ESBL producer by phenotypic tests, which was similar to Owlia *et al.*'s study (21%) in Iran and another study in Poland (20%) (10,20). Despite the above findings, a study in Chile has reported a low rate (10%) of ESBL producer among *A. baumannii* strains, which were collected from pathological products of hospitals by the same DDST method (21).

Resistance to 3<sup>rd</sup> and 4<sup>th</sup> generation cephalosporins and carbapenems among *Pseudomonas aeruginosa* isolated in our study is comparable with a study by Rossolini *et al.* in Latin America, which reported the same rate of resistance to the following antibiotics: ceftazidime (65.95%), cefepime (67%), imipenem (78%) (23). Rate of resistance in our study were 66, 67, and 76 percent, respectively.

Similarity between the resistance rates, which were reported among *Acinetobacter baumannii* in the Owlia's study (98% to ceftazidime, 95% to cefotaxim, and 85% to imipenem) and in our study (100% for all) is another evidence that confirms our results (10).

Whereas, a kind of beta-lactams is inhibited by beta-lactamase inhibitors like clavulanic acid, the use of beta-lactam/beta-lactamase inhibitor combination (21%) therapy may be an alternative treatment, but the effect of this combination varies dependent on the subtype of ESBL present (5).

In conclusion, the antibiogram results in this study reveal that none of the third or fourth generation cephalosporins are suitable for treatment of *Acinetobacter baumannii*.

## Footnotes

### Acknowledgments:

We would like to say our thanks to the personnel of "Pediatric Infections Research Center" for their cooperation.

### Funding/Support:

This study was supported by "Pediatric Infections Research Center" at Mofid Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran (Grant Number: 390). Also this study was part of a Master Degree thesis (Registration Number: 137) in Shahid Beheshti University of Medical Sciences, Tehran, Iran.

### Conflict of Interest:

The authors declare no conflict of interest.



## References

1. McGowan JE Jr. Resistance in nonfermenting gram-negative bacteria: multidrug resistance to the maximum. *Am J Med* 2006 Jun;119(6 Suppl 1):S29-36.
2. Bonomo RA, Szabo D. Mechanisms of multidrug resistance in *Acinetobacter* species and *Pseudomonas aeruginosa*. *Clin Infect Dis* 2006;43(Suppl 2):S49-56.
3. Vila J, Marco F. [Interpretative reading of the non-fermenting gram-negative bacilli antibiogram]. *Enferm Infecc Microbiol Clin* 2002;20(6):304-10. (Full Text in Spanish)
4. Navon-Venezia S, Ben-Ami R, Carmeli Y. Update on *Pseudomonas aeruginosa* and *Acinetobacter baumannii* infections in the healthcare setting. *Curr Opin Infect Dis* 2005;18(4):306-13.
5. Begum S, Salam MA, Alam KhF, Begum N, Hassan P, Haq JA. Detection of extended spectrum  $\beta$ -lactamase in *Pseudomonas spp.* isolated from two tertiary care hospitals in Bangladesh. *BMC Res Notes* 2013;6:7.
6. Bradford PA. Extended-Spectrum B-Lactamases in the 21<sup>st</sup> Century: Characterization, Epidemiology, and Detection of This Important Resistance Threat. *Clin Microbiol Rev* 2001;14(4):933-51.
7. Polotto M, Casella T, de Lucca Oliveira MG, Rúbio FG, Nogueira ML, de Almeida MTG, et al. Detection of *P. aeruginosa* harboring blaCTX-M-2, blaGES-1 and blaGES-5, blaIMP-1 and blaSPM-1 causing infections in Brazilian tertiary-care hospital. *BMC Infect Dis* 2012;12:176.
8. Szabó D, Szentandrassy J, Juhász Z, Katona K, Nagy K, Rókusz L. Imported PER-1 producing *Pseudomonas aeruginosa*, PER-1 producing *Acinetobacter baumannii* and VIM-2-producing *Pseudomonas aeruginosa* strains in Hungary. *Ann Clin Microbiol Antimicrob* 2008;30(7):12.
9. Bayram Y, Parlak M, Aypak C, Bayram I. Three-year review of bacteriological profile and antibiogram of burn wound isolates in Van, Turkey. *Int J Med Sci* 2013;10(1):19-23.
10. Owlia P, Azimi L, Gholami A, Asghari B, Lari AR. ESBL- and MBL-mediated resistance in *Acinetobacter baumannii*: a global threat to burn patients. *Infect Med* 2012;20(3):182-7.
11. Murray PR, Baron EJ, Jorgensen JH, Landry ML, Pfaller MA. Manual of clinical microbiology. 9<sup>th</sup> ed. Washington DC: ASM Press 2007; p: 164, 472-3, 719-28, 749,
12. Performance standards for Antimicrobial Susceptibility Testing (CLSI); Twenty-second Informational Supplement. Wayne, PA, USA: Clinical and Laboratory Standards Institute. CLSI Document M100-S22. 2012, 32(3).
13. Polotto M, Casella T, de Lucca Oliveira MG, Rúbio FG, Nogueira ML, de Almeida MT, et al. Detection of *P. aeruginosa* harboring blaCTX-M-2, blaGES-1 and blaGES-5, blaIMP-1 and blaSPM-1 causing infections in Brazilian tertiary-care hospital. *BMC Infect Dis* 2012;12:176.
14. Lascols C, Hackel M, Hujer AM, Marshall SH, Bouchillon SK, Hoban DJ, et al. Using Nucleic Acid Microarrays To Perform Molecular Epidemiology and Detect Novel B-Lactamases: a Snapshot of Extended-Spectrum B-Lactamases throughout the World. *J Clin Microbiol* 2012;50(5):1632-9.
15. Kalantar E, Taherzadeh S, Ghadimi T, Soheili F, Salimizand H, Hedayatnejad A. *Pseudomonas aeruginosa*, an emerging pathogen among burn patients in Kurdistan Province, Iran. *Southeast Asian J Trop Med Public Health* 2012 ,43(3);712-7.
16. Tsering DC, Das S, Adhiakari L, Pal R, Singh TS. Extended Spectrum Beta-lactamase Detection in Gram-negative Bacilli of Nosocomial Origin. *J Glob Infect Dis* 2009;1(2):87-92.
17. Zarakolu P, Metan G, Hasçelik G, Akova M. [Comparison of different phenotypic methods detecting extended spectrum beta-lactamases in *Pseudomonas aeruginosa* and *Acinetobacter baumannii* strains]. *Mikrobiyol Bul* 2005;39(3):265-72. (Full Text in Turkish)
18. Jabalameli F, Mirsalehian A, Sotoudeh N, Jabalameli L, Aligholi M, Khoramian B, et al. Multiple-locus variable number of tandem repeats (VNTR) fingerprinting (MLVF) and antibacterial resistance profiles of extended spectrum beta lactamase (ESBL) producing *Pseudomonas aeruginosa* among burnt patients in Tehran. *Burns* 2011;37(7):1202-7.
19. Vahdani M, Azimi L, Asghari B, Bazmi F, RastegarLari A. Phenotypic screening of extended-spectrum  $\beta$ -lactamase and metallo- $\beta$ -lactamase in multidrug-resistant *Pseudomonas aeruginosa* from infected burns. *Ann Burns Fire Disasters* 2012;25(2):78-81.
- 20- Sacha P, Wiczorek P, Ojdana D, Czaban S, Kłosowska W, Jurczak A, et al. Susceptibility, phenotypes of resistance, and extended-spectrum  $\beta$ -lactamases in *Acinetobacter baumannii* strains. *Folia Histochem Cytobiol* 2012;50(1):46-51.

21. Pino I C, Domínguez Y M, González R G, Bello T H, Sepúlveda A M, Mella M S, et al. [Extended spectrum beta lactamases (ESBL) production in *Acinetobacter baumannii* strains isolated from Chilean hospitals belonging to VIII Region]. Rev Chilena Infectol 2007;24(2):137-41. (Full Text in Spanish)

22. Jacoby GA, Muñoz-Price LS. The New beta-Lactamases. N Engl J Med 2005;352:380-91.

23. Rossolini GM, Mantengoli E. Treatment and control of severe infections caused by multi resistant *Pseudomonas aeruginosa*. Clin Microbiol Infect 2005;11 Supplement 4:17-32.

Archive of SID