ORGINAL ARTICLE

The Relationship between Blood Buffer Capacity and Aerobic Power with Ability to Perform High Intensity Intermittent Activities in Young Football Players

*Gaeini, A.A;*¹ *Kashef, M;*² *Joudaki Nejad, H.R;*³ *Kazemi, F*⁴ ^{1,4} *Department of Exercise Physiology, Tehran University, Tehran, Iran* ^{2,3} *Department of Exercise Physiology, Shahid Rajaee Teacher Training University, Tehran, Iran*

Abstract

Background: The purpose of this study was to determine the relationship between blood buffer capacity and aerobic power (Vo_{2max}) with ability to perform high intensity intermittent activities in young football players. Methods: 27 male football players (age 17.66 ± 0.49 years) selected as subjects. Blood samples were collected before and after the yoyo intermittent test. Then, after 48 hours, the Bruce treadmill test was performed for measuring Vo_{2max}. The Pearson coefficient correlation was used for analyzing data. Results: A significant relationship was found between H^+ , Hco_3^- , BE and ability to intensity perform high intermittent activities (r=0.609, p=0.001; r= -0.381, p=0.050; r=-0.401, p=0.038), a significant relationship between Vo_{2max} and ability to perform high intensity intermittent activities (r=0.761, p=0.000) and no significant relationship between Pco2 and perform ability to high intensity intermittent activities (r=-0.130, p=0.517). Conclusion: Results indicate that blood buffer capacity and Vo_{2max} are effective factors in the performance of intermittent activities in football players.

Key Words: Blood buffer capacity,

Gaeini, A.A (⊠) aagaeini@yahoo.com

Vo_{2max}, high intensity intermittent activities, young football players

Introduction

During the recent years, sport and physical education has advanced increasingly. Therefore. identification of effective factors in performance may help coaches and sport scientists to develop better training programs. A large number of team games such as football, basketball and hockey require participants to have the ability to perform high intensity intermittent activities with short- term recovery periods and maintain speed until termination of exercise. This performance ability requires desired recovery intervals during high intensity intermittent activities for fighting fatigue (Wadely & Rossignol., 1998; Bangsbo et al., 1996). Following the repeated intermittent activities such as football, the recovery should not last less than a few minutes. Such activities which cause depletion of ATP-PCr and increase in lactate and H⁺ accumulation require longer recovery periods (Tamlin & wenger., 2001). Increased buffer capacity and blood flow elevate Lactate removal from muscles (Holoszy & coyle., 1984). Decrease in intracellular PH reduces ability perform high intensity to intermittent exercises. Decrease in intracellular acidosis declines ability to perform intermittent sprint exercises via inhibition

of

glycolysis

1981). There is no clear evidence how

fatigue occur during the high intensity

(Hermansen.,

1986). In a research Krustrup et al (2003) physiological considered intermittent activities in football players. results during the

intermittent activities. In most studies it is known that metabolic factors (Lactate accumulation and decreased PH) (Tamlin & wenger., 2001) and depletion of muscle glycogen (Krustrup et al., 2003) play an important role in the development of muscle fatigue during high intensity intermittent activities. Base on some researches, the lactate accumulation does not make fatigue during high intensity intermittent activities. There is also no relationship between decrease in PH and decrease in performance during football race (Krustrup et al., 2006). Increase in H+ buffering may improve the ability to intensity perform high intermittent activities by facilitating resynthesis of PCr or decreasing inhibition of glycolysis (Bishop & Spencer., 2004). Lactate removal from muscles resulting from increasing buffer capacity is one of effective factors in increasing performance during the intermittent activities (Holoszy & coyle., 1984). On the other hand, VO_{2max} is a main factor during the intermittent activities and during recovery intervals of the intermittent activities (McMahon & Wenger., 1998)⁻ Thus, the buffer capacity and VO_{2max} may be the main factors in performing intermittent activities (Bishop & Spencer., 2004). **Bicarbonate** (Hco_3) is important extracellular buffer and phosphates, proteins and hemoglobin are intracellular buffers (Holoszy & coyle., 1984).

In general, only a few studies have been published to determine muscle metabolism and fatigue during the high intensity team games with brief recovery periods (Bishop et al., 2004). Zoladez et al (1993) reported a significant increase in blood lactate and decrease in Hco₃, Pco₂, PH and BE after the incremental activities (Zoladz et al., 1993). McCarteney et al (1993) found a decrease in blood PH and no change in Hco_3 , Pco_2 and Po_2 during each stage of intermittent work (McCartney et al.,

showed no relationship The between fatigue and muscle lactate and H⁺ levels high intensity activities(Krustrup et al., 2003). The results of research Bishop et al (2004) indicated significant relationship a between muscle buffer capacity and significant aerobic power and no relationship between total work and buffer capacity during the repeated- sprint activities (Bishop et al., 2004). In a research of Krustrup et al (2006) muscle PH of 40 elite football player during the yo vo intermittent test reaches to 6.8 in a fatigued state. The results suggested that buffer capacity and aerobic power are main factors in performing intermittent activities (Krustrup et al., 2006). Edje et al (2006) reported decrease in pre-exercise H^+ , no change in pose-exercise H^+ and improved endurance performance and lactate threshold lactate after 8- weeks consuming of NaHco3 supplement (EdJe et al., 2006). Böning et al (2007) indicated changes in intracellular buffers and no change in the rate of Hco_3^- during and after exercise (Böning et al., 2007). Bishop et al (2008) found decrease in lactate and H^+ accumulation after 5-week high intensity intermittent training (Bishop et al., 2008). Based on the previous studies the factors such as PH, lactate and CP play an important role in the development of muscle fatigue during high intensity intermittent activities (Krustrup et al., 2006). It seems that the intensity of activity, the number of repetitions, duration of exercise and rest intervals in recovery periods may influence on performance during intensity high intermittent activities (Bangsbo et al., 1996). Therefore the present study designed to determine the relationship between blood buffer capacity and aerobic power with ability to perform high intensity intermittent activities in young football player and evaluate how much the

to

response

performance ability can be influenced by buffer capacity and aerobic power.

Methods

Subjects

Among of the members of two teams "Naft and Mehrene- reza", 36 male football players in the first class league of Tehran city (age 17.66 \pm 0.49 years) volunteered as subjects. 9 of them were deleted during the test and 21 of them performed the test. Subjects participated in this study after given their written informed consent.

Procedures

Exercise test

The Bruce treadmill test

The Bruce treadmill test was used as exercise test for measuring VO_{2max} . In the Bruce test, treadmill is set up with the Stage 1 speed (2.74 km / hr) and grade of slope (10%) and the athlete commences the test. At the appropriate times during the test, the speed and slope of the treadmill are adjusted. Therefore, after 3 minutes into the test the speed is adjusted to 4.02 km/hr and the slope to 12%, after 6 minutes into the test the speed is adjusted to 5.47 km/hr and the slope to 14%, and so on. The test stops when the athlete is unable to continue - this ideally should be between 9 and 15 minutes.

The Yo-Yo intermittent recovery test

The test is measured the variables H^+ , Pco_2 . Hco₃ and BE. The Yo-Yo intermittent recovery test consists of repeated 2.20 -m runs back and forth between the starting, turning, and finishing line at a progressively increased speed controlled by audio bleeps from a tape recorder. Between each running bout, the subjects have a 10-s active rest period, consisting of 2.5 m of jogging. When the subjects twice have failed to reach the finishing line in time, the distance covered is recorded and represents test result. The test may be performed at two different levels with differing speed profiles (level 1

and 2). in the present study, we used the Yo-Yo intermittent recovery test, level 1, which consist of 4 running bouts at 10-13 km.h-1 (0-160m) and another 7 runs at 13.5-14 km .h-1 (160-440m), where after it continues with stepwise 0.5 km.h1 speed increments after every 8 running bouts (i.e., after 760, 1080, 1400, 1720 m. etc.) until exhaustion .the test was performed indoor on running lanes, marked by cones,, having a width of 2 m and a length of 20m. Another cone placed 5 m behind the finishing line marked the running distance during the active recovery period. Before the test, all subjects car tried out a warm up period consisting of the first four running bouts in the test. The total duration of the test was 6-20 min. all subject were familiarized to the test by at least one pretest (Aziz et al., 2003).

The Blood samples were collected before the yoyo test. Then after warm- up, the yoyo intermittent test was performed and immediately after the test, blood samples were collected. After 48 hours, subjects arrived at Olympic national Academy in part "the evaluation center of physical capabilities" before noon and they performed the Bruce test for evaluating Vo_{2max}. The Bruce test was performed using treadmill (k₄b₂, techno Gym, Italy). The blood samples were collected pre and post- vovo test. 1 cm³ of blood sample was collected from the radial artery in wrist using plastic syringe. After heparin of syringe, the blood sample was collected from radial artery. Then the needle was bended to prevent the air enters the saying and the lid was covered with a plastic and transformed to blood laboratory. The samples were examined using device of the measurement of blood gas and the

Statistical analysis

Statistical significance was assessed using Pearson coefficient correlation. The statistical package SPSS Version 16 was used for statiscal analysis. Differences

result was printed on special sheets.

were considered significant at the $p \le 0.05$ and results were presented as means \pm SD.

Results

The physical characteristics, variables resulting from the yoyo intermittent test and Bruce test in male athletes (n=27) (Table 1) were recorded.

Table 1. Physical characteristics of the athletes (n=27)									
Variables	Age (yrs)	Height (cm)	Weight (kg)	Fat (%)					
Distance(cr	m) Vo _{2m}	_{ax} (ml/kg/min)							
(M ±SD)	17.66 ± 0.49	173.47 ± 6.03	64.33 ± 5.31	14.28 ± 3.38	1640.74				
±410.94	49.76 ± 5.33								

Table 2. Descriptive statistics (M±SD) for variables in pre and post-test in male athletes (n=27)

Variables		M±SD	Max	Min			
	Pre-test	38.802±1.4	40.65	36.131			
$\mathbf{H} \vdash (\mathbf{mag}/\mathbf{I})$	0 0						
$\Pi + (\Pi eq/L)$	Post-test	55.84 ± 2.76	63.37	43.241			
			8				
U_{aa} (mm a $1/L$)	Pre-test	22.53 ± 1.90	25.9	18.2			
HCO3-(IIIIIIOI/L)	Post-test	12.75 ± 2.76	18.8	9.2			
Dag 2 (mmIIa)	Pre-test	36.44±3.26	42.1	31.3			
PCO2 (IIIIIFIG)	Post-test	28.44 ± 4.55	33.3	26.10			
DE(mmol/L))	Pre-test	-1.14 ± 1.50	1	-3.8			
DE (IIIII0I/L))	Post-test	-12.65±3.21	-5.1	-18.5			

(M±SD) Descriptive statistics for measured variables in pre and post-test (Table 2) and the results of Pearson coefficient correlation between variables in male athletes (n = 27) (Table 3) were recorded. Using Pearson coefficient correlation, significant relationship were found between H+ (r= 0.609, $p \le 0.001$); Hco_3 (r= -0.381, p ≤ 0.05); BE (r= 0.401, $p \le 0.05$), Vo_{2max} (r= 0.761, $p \le 0.000$) and performance ability of High intensity intermittent activities. Also no significant relationship was found between Pco₂ (r= 0.130, p= 0.517) and performance ability of High intensity intermittent activities.

Discussion

There was a significant relationship between H^+ and ability to perform high intensity intermittent activities. This result is supported by the results of Pilegaard et al (2000), Edje et al (2003) and Bishop et al (2008)^[15], but is not in agreement with the results of Edje et al (2003) and Krustrup et al (2003). This contradiction may be due to the difference in training protocol (in terms of intensity, number of repetition and recovery periods) and also type of training (intermittent vs resistance). During the high intensity exercise, H^+ is produced by volatile acids (e.g carbonic acid), Non-volatile acids (e.g sulfuric acid and phosphoric acid) and organic acids (e.g lactic acid and acetoacetic acid). Our result indicate that during high intensity intermittent activities the concentration of intracellular H^+ increase through production of mentioned acids and their metabolic processes (Scott & Powers., 2006). As bishop et al (2003) reported positive significant relationship between H^+ concentration and the repeated sprint ability in elite female (bishop et al., 2003). In this matter, some researchers found similar results. Krustrup et al (2006) concluded that the H^+ concentration of elite players increased during the yo yo intermittent test and decreased after 3-mim rest (Krustrup et al., 2006).

There was a significant relationship between Hco₃⁻ and ability to perform high intensity intermittent activities (r=-0.381, $p \le 0.05$). The findings of results of McCarteney et al (1986), Zoladez et al (1993) and Krustrup et al (2006) are in agreement with our result. However, the result of Böning et al (2007) is not supported our finding and indicated that intermittent activities had more effect on buffer capacity compared with the continuous activities. This may be due to type of training protocol and physical activity (Böning et al., 2007). Edje et al (2003) found the relationship between the Hco₃ buffer and the repeated sprint ability in 3 groups (team athlethes, trained and untrained endurance athletes). However, this relationship between team athlethes more than others. The results was that intermittent suggested trainings improved the buffering system (Edge & Bishop., 2003).

There was no significant relationship between Pco_2 and ability to perform high intensity intermittent activities (r=-0.130, p=0.517). The previous study of Dona et al (2001) and Hermansen., (1981) are supported this result, but the result of Böning et al (2007) is not in agreement with our finding. This may be attributed to difference in the type of blood sample (venous vs. artery) and type of exercise (swimming vs. football). During high intensity activities, pulmonary ventilation increases because of metabolic acidosis. Increased pulmonary ventilation cause more exertion of Co_2 from pulmonary capillaries and decrease in Pco_2 .

There was a significant relationship between BE (Base excess) and ability to perform high intensity intermittent activities (r=0.401, p≤0.05). This result is supported by the results of Böning et al (2007). However, our result is not in agreement with the result of McCartney et al (1986). This may be related to difference in intensity, duration, number of repetition and recovery periods. BE or the measurable concentration of blood Base decrease after high intensity activities. Decrease in BE resulting from increasing H⁺ disorder Acid-Base balance. Therefore, there is relationship between BE and H⁺ levels after high intensity activities (Scott & Powers., 2006). Zoladez et al (1993) examined changes in acid and base in runner during an incremental intermittent test and found a significant increase in lactic acid and a significant decrease in BE in a fatigued state (Zoladez et al (1993).

There was a significant relationship between Vo_{2max} and ability to perform activities high intensity intermittent (r=0.761, p≤0.001). The strong relationship between Vo_{2max} and ability to high intensity intermittent perform activities show that Vo_{2max} is an important factor during high intensity intermittent activities. Our result is in agreement with results of Bishop et al (2004), Edge et al (2005) and Krustrup et al (2006) and is not in agreement with the results of Bishop et al (2003. It seems that sexuality, type of exercise, protocol of training (in terms of the number of the repetition, rest intervals periods) of recovery may effect on performance high during intensity intermittent activities. However, Tamlin and venger (2001) reported a significant relationship between Vo_{2max} and aerobic responses to fast recovery periods.the Mcmahon and Venger (1998) indicated that Vo_{2max} is a main factors during high intensity intermittent activities. Bishop et

al (2004) showed a significant relationship between Vo_{2max} and the ability to perform repeated sprint activities (Bishop al., 2004). Although many factors effect on the perform ability to high intensity intermittent activities, acid lactate production and increased H⁺ concentration in sarcoplasm of muscle cell disorder Acid-Base balance and decrease partly intracellular PH and finally cause fatigue. Studies show that increased intracellular H^+ may result a disturbance in energy production and in fact, H^+ can be an inhibitory factor in contractile mechanism (McCartney., 1986). Increased blood lactic acid enhances plasma lactate and H^+ concentration. Also Increased H^+ concentration elevate depth and frequency of breathing (Scott & Powers., 2006). The factors improving Vo_{2max} can be defined as: Increased oxidative enzymes; increased number, size and rate of mitochondria and myoglobin, increased blood flow (by increased cardiac increased output), capillary Network of muscle tissue and vasodilation. In fact, increased aerobic during intermittent portion activities decrease lactate production. Also lactate removal from muscle resulting from increasing buffer capacity and blood flow improve the ability to perform high intensity intermittent activities (Holoszy & coyle., 1984).

Conclusion

Blood buffer capacity and Vo_{2max} are effective factors in the performance of intermittent activities in football players.

References

- Aziz, A.R., Tan, F.Y.H; Teh, k. (2005). "Comparing two field tests with the treadmill Run test in soccer players". J sci and med, 4, 105-112.
- Bangsbo J, Madsen K, Kiens B, Richter EA.. (1996). "Effect of muscle acidity on muscle metabolism and fatigue during intense exercise in man". j physiol, 1;495 (Pt 2):587-96..

- Bishop D, Edge J, Goodman g. (2004). "Muscle buffer capacity and aerobic fitness is associated with repeated – sprint ability in women does". sic med sport, 7, 199-209.
- Bishop D, Lawrence s, Spencer M. (2003)." Predictors of repeated sprint ability in elite female hockey players". sci med sport, 6 (2), 199-209.
- Bishop D; Edge J; Tomas C; Mercier J. (2008). "Effects of high-intensity training on muscle lactate turn sport and post exercise recovery of muscle lactate and hydrogen ion in woman" .j physiol ,295,118-132.
- Bishop D; Spencer M. (2004). "Determinants of repeated – sprint ability in well – trained team sport athletes and endurance – trained athletes". j sports med fitness' 44, 1-7.
- Böning D, Klarholz C, Himmelsbach B, Hütler M, Maassen N. (2007). "Extra cellular bicarbonate and non bicarbonate buffering against lactic acid during and after exercise". j Apple physiol, 100(4):457-67
- Edge j, Bishop d, Goodman c, Dowson, B. (2005). "The effects of high and moderate – intensity training of metabolism and repeated – sprints". Med Sci Sports Exerc. 37(11):1975-82.
- Edge J, Hill-Haas S, Goodman C, Bishop D. (2006). "Effects of resistance training on H⁺ regulation capacity, and repeated sprints". Med sci sports Exec 38(11):2004-11.
- Edge J; Bishop D. (2003). "Difference muscle buffer capacity between team sports, endurance trained and untrained". Sci and med sport, 6, 11-21.
- EdJe J, Bishop D, and Goodman c. (2006). "Effects of chronic NaHco3 ingestion during interval training on changes to muscle buffer capacity, metabolism and short – term endurance performance". Appl physio, 225-234.
- Edje J, Bishop D.duffield R. (2002). "Muscle metabolism during repeated

spring in untrained femal". J sci and med in sport, 3, 190-202.

- Hermansen D. (1981). "Muscle fatigue maximal exercise of short duration in: physiological chemistry of exercise and training". Medicine and sport science, karager, Basel, 45-52.
- Holoszy j, coyle EF. (1984). "Adaptation of skeletal muscle to Endurance exercise and their metabolic convey uneces". J Apple physiol, 56, 831-838.
- Krustrup, P; Mohr, M; Amstrup, T; Rysgaard, T., Johnjsen, J; Bangsbo, J. et al (2003). "The yo yo intermittent Recovery test: physiological reponse, Relibility and validity". Med. Sci. sports Exer. 35 (4), 697-705.
- Krustrup; p; mohr, M; Nybo, G., Jensen, J., Nimlsen, g., Bangs Bo b. (2006).
 "The yo yo IR2 Test: physiological Response, Reliability, and Application to elite Soccer". Med. Sci sports Exerc. 38(9): 1666-73.
- Mc Mahon S, Wenger HA. (1998). "The relation between aerobic fitness and both power output and subsequent recovery during maximal intermittent exercise". j sci Med sport,1,4; 219-227.
- McCartney N, Spriet LL, Heigenhauser GJ, Kowalchuk JM, Sutton JR, Jones NL (1986). "Muscle power and

metabolism in maximal intermittent exercise". J appl, physiol. 1164-1169.

- Pilegaard H, Domino K, Noland T, Juel C, Hellsten Y, Halestrap AP, et al (2000).
 "Effect high – intensity exercise training on lactate/ H + transport capacity in human skeletal muscle". j, physiol, 276, 255-261.
- Scott k,.Powers T. (2006). "Exercise physiology: theory and application to fitness and performance".252-324.
- Tamlin DL; wenger H. (2001). "The relationship between aerobic fitness and recovery from high intensity intermittent exercise". Sport med, 31(1), 1-11.
- Tomlin DL, Wenger HA. (2001)."Relationship Between Aerobic Fitness and Recovery from High intensity intermittent Exercise". Sports Med. 31(1):1-11.
- Wadely G, Rossignol P. (1998). "The relation between repeated sprint ability and aerobic and an aerobic energy system". sci med sport, 1, (2), 100-110.
- Zoladz J. A, Sargeant A. J, Emmerich J, Stoklosa J, Zychowski A. (1993). "Changes in acid-bas status of marathon runners during an incremental field test". Frg 67(1): 71-76.