

A systematic review of questionnaires used on oral health knowledge, attitude, and practice in 12-year-olds

Maryam Rad DDS, MSc¹, Arash Shahravan DDS, MSc²,
Ali Akbar Haghdooost MD, PhD³

Review Article

Abstract

BACKGROUND AND AIM: National oral health knowledge, attitude, and practice (KAP) data among 12-year-old children need nation-wide programs to help promote oral health. In most countries, oral epidemiologic data are collected by self-administered structured questionnaires. The aim of this study was to undertake a systematic review of the existing literature about questionnaires used for analyzing the oral health knowledge, attitude, and behavior profile of the 12-year-old children.

METHODS: The search was conducted in PubMed and Google Scholar search engines. The Medical Subject Heading (MeSH) search was performed applying singularly and by combining the following terms retrieved from the MeSH browser provided by MEDLINE: "Knowledge, awareness, attitude, practice, behavior, 12-year-old children, oral health."

RESULTS: Of 176 records found, 29 evaluated oral health KAP in 12-year-old children by structured questionnaires. The most important questions on knowledge (23 of 29 questionnaires) included some items which are focused on the importance of preserving natural teeth, effects of brushing, and sweets and soft drinks on the dentition. The most common questions on practice (28 questionnaires) were brushing activity and dental visits. The attitudes were evaluated by questions about fear of dental treatment, opinion about dentists and dental care (10 questionnaires).

CONCLUSION: Considering differences in the available questionnaires showed that despite the importance of promotion of oral health by increasing knowledge, and improving attitudes and practice in 12-year-old children, more work is needed to form a standard questionnaire.

KEYWORDS: Oral Health; Knowledge; Attitude; Behavior; Review

Citation: Rad M, Shahravan A, Haghdooost AA. **A systematic review of questionnaires used on oral health knowledge, attitude, and practice in 12-year-olds.** J Oral Health Oral Epidemiol 2016; 5(1): 1-12.

Dental caries and gum disease are the most common diseases in human populations that affecting over 80% of school children in some countries.^{1,2} During the past decades, the common consensus from many reports was that prevalence of dental caries in children and adolescents had declined significantly in developed countries in contrast to developing countries.^{3,4} There are, however, recent studies

that clearly indicates a marked increase in the prevalence of dental caries in many countries. It appears that the main reasons for this global increase are unhealthy dietary habits, and inadequate oral hygiene practices.⁵

To prevent and control oral diseases, improvements in knowledge, attitudes, and behaviors related to oral health among school children have been recommended by the World Health Organization (WHO).⁶ The

1- PhD Candidate, Oral and Dental Diseases Research Center AND Kerman Social Determinants on Oral Health Research Center, Kerman University of Medical Sciences, Kerman, Iran

2- Associate Professor, Endodontology Research Center, Kerman University of Medical Sciences, Kerman, Iran

3- Professor, Regional Knowledge Hub, WHO Collaborating Centre for HIV Surveillance, Research Center for Modeling in Health, Institute of Futures Studies in Health, Kerman University of Medical Sciences, Kerman, Iran

Correspondence to: Ali Akbar Haghdooost MD, PhD

Email: ahaghdooost@gmail.com

12-year age group is especially important as it is the age at which all the permanent teeth, except the third molars, have erupted. Therefore, this age has been selected as the global monitoring age for caries for international comparisons and future planning of oral health programs.⁷

To promotion of the oral health, the planning and assessing of school-based oral health programs would be needed. For planning these programs, analysis of the oral health situation including information on oral health knowledge, attitude, and practice (KAP), would be essential.⁸ Despite the high rate of dental caries in 12-year-old,⁹ socio-epidemiologic data about oral health behavior of children are not available at national level. Therefore, KAP studies to collect such information and the assessment of oral health in children and adolescents are considered to be an essential prerequisite.

In most countries, oral epidemiologic data about KAP were collected by self-administered structured questionnaires.^{6,8} In some studies, the construction of questionnaires has been based on experiences gained from surveys carried out by the WHO. These questionnaires vary widely in terms of the number of items and format of questions and responses. Therefore, there is a need for a standard questionnaire. The aim of this study was to carry out a systematic review of the existing literature on these questionnaires. This report is a part of a study with the aim of designing an appropriate questionnaire for the analysis of oral health behavior profile of 12-year-old children.

Methods

A computerized search was conducted using PubMed, Google Scholar between January 01, 2002, and August 31, 2013. The Medline search uses a complex search strategy, including Medical Subject Heading (MeSH). After initial evaluations, main keywords (knowledge, attitude, behavior, practice, oral health, questionnaire, and 12-year-old children) in

both singular and combined types were determined by MESH system. "OR" between the synonyms and "AND" between the main keywords were used to import the keyword combinations in the reviewed sites.

In addition, a search was conducted on Google Scholar with the same keywords and relevant studies cited in the reference lists of the selected papers were considered. First, the exclusion criteria of the irrelevant articles were applied in the three steps of title, abstract, and the full-text. The full-texts of all the related studies were assessed by one author. If there was any doubt or question, there was consultation with corresponding author/epidemiologist and statistical advisor. After that all the full-texts of the relevant articles were reviewed carefully, and the references of all selected articles were reviewed to identify any additional studies. A guideline for conducting a KAP study was used to evaluate the quality of articles.¹⁰ Based on this guideline, steps in the preparation of a KAP study are: (1) domain identification, (2) question preparation, and (3) validation of questions. Validation of questions should be aimed at assessing their ease of comprehension, relevance to their intended topics, effectiveness in providing useful information, and the degree to which the questions are interpreted and understood by different individuals.

Unpublished studies and abstracts were not considered for inclusion in this systematic review. Full-text of three studies was not available and in spite of attempts to contact the authors, this not achieved, so these studies were excluded from the study. Finally, 29 out of 496 articles remained at the end of these steps (Figure 1).

All items in knowledge, practice, and attitudes areas were determined, and the number of items and format of questions and responses were reviewed. Finally, the questions were categorized based on different items in three areas of knowledge, practice or behavior, and attitudes.




Figure 1. Flow diagram of studies of knowledge, attitude, and practice on oral health among 12-year-old children consisted for inclusion

Results

First, we identified 496 citations meeting our search criteria. From these articles, 29 evaluated oral health in 12-year-old children by structured questionnaires.^{3,4,6,8,11-35} Table 1 shows information about the name of the first author, country, year of publication, the age of subjects, and sample size of each study. In the most studies, the survey instrument was an administered pre-tested questionnaire. Knowledge questions have been asked in 23 of 29 articles. The characteristics of questionnaires have been shown by figure 2. Figure 2 indicates that only 48.3% of articles have reported on the validity and reliability of structured questionnaires, and the number and form of questions and responses were highly variable.


Knowledge area

The most important questions on knowledge included items on the importance of preserving natural teeth, effects of brushing and using fluoridated toothpaste, the meaning of bleeding gums and caries and how to protect against it, and the effects of sweets and soft drinks on the dentition. Table 2 shows the importance of keeping natural teeth of knowledge area (7 different formats) and knowledge of sweet and soft drink effects on the dentition (3 different formats).

One important and common aspect of knowledge questions was about the role of fluoride in oral health, which were addressed in five formats, but the dominant one was: "The use of fluoride prevents tooth decay; true/false/do not know."

Table 1. Characteristics of studies included in review

First author	Country	Age (year)	Sample size (n)
Rajab et al. ²²	Jordan	6-16	1556
de Almeida et al. ¹¹	Portugal	6 and 12	800 (12-year-old)
Zhu et al. ⁴	China	12-18	4400
Varenne et al. ³¹	Burkina Faso	12	505
Al-Omiri et al. ³⁵	North Jordan	10-16	557
Ahmed et al. ³⁴	Iraq	12	392
Grewal and Kaur ¹⁵	India	11-16	200
Singh ¹⁷	India	5-12	377
Smyth et al. ¹²	Spain	12	1105
Harikiran et al. ⁸	India	11-12	212
Kawamura et al. ²⁴	Japan	4-12	1584
Ouma and Martha ³²	Soweto	6-12	336
Petersen et al. ²¹	Chin	11-15	2662
Jürgensen and Petersen ²⁸	Laos	12	621
Granville-Garcia et al. ²⁷	Brazil	10-19	679
Hysi et al. ²⁵	Albania	12	372
Lian et al. ²³	Sarawak	12	209
Prasad et al. ³⁰	India	12-15	652
Shenoy and Sequeira ¹⁸	India	12-13	415
Diwan et al. ¹³	India	3-19	798
Vakani et al. ¹⁶	Pakistan	11-12	300
Gathecha et al. ²⁶	Kenya	12	639
Haleem et al. ¹⁴	Pakistan	10-12	1517
Mehta and Kaur ⁶	India	12	440
Mafuvadze et al. ³³	Zimbabwe	12	205
Prasai et al. ¹⁹	Nepal	5-16	361
Sharada et al. ²⁰	India	12-13	514
de Silva-Sanigorski et al. ²⁹	Australia	5-6 and 11-12	377
Suprabha et al. ³	India	11-13	858

**Figure 2.** The characteristics of questionnaires

CQ: Closed question; C and OQ: Closed and open question; MCQ: Multiple choice question; S and MCQ: Combination short and multiple choice question

Table 2. The item “importance of keeping natural teeth” and “effects of sweets and soft drinks on the dentition” of knowledge questions

Questions	Questionnaires
1. Keeping natural teeth is important for general well-being a. True b. False c. Do not know	India-Mangalore ³ Mofolo, Soweto ³¹ India-Mangalore ¹⁷
2. Do you think keeping your mouth clean and healthy is good for health? a. Yes b. No c. Do not know	Panchkula, India ⁶ China ⁴ Karachi ¹⁵ India ¹⁶
3. General body health has a relationship to oral health and dental diseases a. True b. False	Kuching, Sarawak ²² Jordan ³⁴
4. Keeping natural teeth is not that important a. Agree b. Disagree c. Do not know	India-Bangalore ⁸ Burkina Faso-Africa ³⁰
5. Teeth are an important part of your body a. Agree b. Disagree c. Do not know	Panchkula, India ⁶ Japan ²³
6. Natural teeth are better than false teeth a. True b. False c. Do not know	India-Mangalore ³
7. Keeping natural teeth is not that important a. Agree b. Disagree c. Do not know	Burkina Faso-Africa ³⁰
8. Sugar causes tooth decay a. A lot b. Quite a lot c. Not much d. Not at all	India-Mangalore ³ India-Mangalore ¹⁷ Kuching, Sarawak ²² Zimbabwe ³² Nepal ¹⁸ India ¹⁹ Jordan ³⁴
9. Eating and drinking sweet food does not cause decay a. True b. False c. Do not know	India-Mangalore ³ India-Bangalore ⁸ Panchkula, India ⁶
10. Do you know which of the following food items is/are the main cause of tooth decay? a. Sweets/chocolates/biscuits/cakes/chips/waffers b. Fresh fruits c. Raw vegetables d. Do not know	Karachi ¹⁵ India ¹⁶ Australia ²⁸

Attitude area

Attitude questions had been asked by 10 articles. The attitudes toward oral health have been evaluated by questions on the importance of natural teeth (Table 3), fear of dental treatment, feelings regarding the treatment, thoughts about involvement in the dental treatment, opinion about and attitudes toward dentists

and dental care and regular dental visits.

Questions on the importance of natural teeth were asked in several ways. The most common questions in this item were:

1. Decay makes my teeth look bad:
 - a. Fully agree
 - b. Agree
 - c. Disagree

Table 3. The items of attitude question about importance of natural teeth

Questions	Questionnaires
1. It is important to take care of my teeth a. Fully agree b. Agree c. Disagree d. Fully disagree e. No opinion	Spain ¹¹
2. You care about your teeth as much as any part of your body a. Yes/no	Jordan ³⁴
3. Decay makes my teeth look bad a. Fully agree b. Agree c. Disagree d. Fully disagree e. No opinion	India-Bangalore ⁸ Spain ¹¹ India ¹⁶ 13-Jordan ³⁴
4. Dental problems can cause other health problems a. Fully agree b. Agree c. Disagree d. Fully disagree e. No opinion	Japan ²³
5. Dental problem can be serious a. Fully agree b. Agree c. Disagree d. Fully disagree e. No opinion	Japan ²³
6. Dental disease is less important than other health problems a. Fully agree b. Agree c. Disagree d. Fully disagree e. No opinion	Japan ²³
7. It is natural for people to lose all their teeth in old age a. Fully agree b. Agree c. Disagree d. Fully disagree e. No opinion	Japan ²³
8. Milk teeth are not important because they fall out soon a. Fully agree b. Agree c. Disagree d. Fully disagree e. No opinion	Japan ²³
9. I am able to prevent my teeth from decaying a. Fully agree b. Agree c. Disagree d. Fully disagree e. No opinion	Japan ²³

- d. Fully disagree
- e. No opinion.

Practice area

The practice of 12-year-old children has been assessed by 28 articles. The most common questions about the practice were tooth brushing habits (TBH) (such as frequency, duration, and brushing aides) (Table 4), the frequency of sugar consumption and drinking soft drinks, and dental visits (such as regularity, reasons behind the visits and the effect of pain on the first visit). Questions in some of the items had almost the same form. Table 3 shows the item TBH of practice

questions. In this item although questions were asked almost the same format, but the number of questions was different. In some studies, the details of brushing activity such as time of brushing, type of toothpaste, and frequency in a change of toothbrush were determined. The most common question in this item was:

2. How often do you brush your teeth?
 - a. Less than once a week
 - b. Once a week
 - c. 2-3 times a week
 - d. Twice or more a day
- Every day do not know/no answer.

Table 4. The item of brushing activity of practice questions

Questions	Questionnaires
1. How often you brush your teeth?	China ⁴
a. Less than once a week	Panchkula, India ⁶
b. Once a week	India-Bangalore ⁸
c. 2-3 times a week	Uttarakhand-India ¹²
d. Every day	Amristar, India ¹⁴
e. Twice or more a day	Kuching, Sarawak ²²
f. Do not know/no answer	Albania ²⁴
	Kenya ²⁵
	Laos ²⁷
	Australia ²⁸
	Mofolo, Soweto ³¹
	Jordan ³⁴
2. How many times you brush daily?	India-Mangalore ³
a. Less than once per day	Portugal ¹⁰
b. Once per day	Pakistan ¹³
c. Twice per day	Karachi ¹⁵
d. More than twice per day	India ¹⁶
	India-Mangalore ¹⁷
	Nepal ¹⁸
	Brazil ²⁶
	Zimbabwe ³²
	Iraq ³³
	Spain ¹¹
3. The last time I brushed my teeth was	
a. Today/yesterday/day	
b. Before yesterday/more than 2 days ago	
4. Type of brush	India-Mangalore ³
a. Plastic toothbrush	Uttarakhand-India ¹²
b. Charcoal/chew sticks	Thiruchengode-India ²⁹
c. Miswak/wooden	
d. Plastic toothpicks/finger/brush	
e. Finger/do not know/no	

Table 4. The item of brushing activity of practice questions (Continue)

Questions	Questionnaires
5. Which of the following do you use for cleaning your teeth? a. Toothbrush b. Finger c. Chew stick/miswak	India-Mangalore ³ Panchkula- India ⁶ India-Bangalore ⁸ Uttarakhand-India ¹² Pakistan ¹³ Amristar, India ¹⁴ India ¹⁶ Nepal ¹⁸ Kenya ²⁵ Mofolo, Soweto ³¹ Jordan ³⁴
6. Type of toothpaste a. Fluoridated b. Non-fluoridated c. Do not know	India-Mangalore ³ China ⁴ India-Bangalore ⁸ Amristar, India ¹⁴ China ⁴
Started brushing teeth: Before schooling when attending primary school after completing primary school	Mofolo, Soweto ³¹ India ¹⁶
7. Rinse your mouth after each meal a. Yes b. No	India-Mangalore ¹⁷ Thiruchengode-India ²⁹
8. Time of brushing a. Morning only b. Morning and night c. Night only d. After every meals	India-Mangalore ³ China ⁴ Jordan ³⁴
9. When will you change your toothbrush? a. 1-3 months b. 4-6months c. 1 year d. Above/after bristles e. Splayed/do not know	India-Mangalore ³ India ¹⁶ India-Mangalore ¹⁷ Thiruchengode-India ²⁹
10. Do you use any other oral hygiene aids? a. Dental floss b. Interdental brush c. Toothpicks d. Mouthwash e. None f. Do not know	Amristar, India ¹⁴ India ¹⁶

Discussion

Based on strong evidence, in developing countries, oral hygiene is one of the main health considerations.^{4,36} Therefore, repeated KAP studies is highly recommended to monitor trends; however, a valid and culturally adapted questionnaire is one of the main pre-requisites. This systematic

review showed that despite the importance of the 12-year age group, there is not a standardized questionnaire for evaluating oral health knowledge, attitude, and behavior in these children.

In the majority of studies (51.7%), there was not a report on validity and reliability of questionnaires. Furthermore, these tools vary

widely in terms of the number of items and format of questions and responses. However, there was the high similarity between the content of questionnaires. The most similarity was between practice questions, and the least one was between attitude questions. The most common format of questions was a combination of multiple choices and yes-no answers, and the most of the questions were asked in closed-ended format. The number of questions had a wide range. The most common number of questions was 10-14 and 15-19. Perhaps the reason is some studies that have assessed only one or two areas. It seems that an appropriate and comprehensive questionnaire that evaluate KAP (different domains such as brushing habits, diet, and dental visits), should contain 30 questions. The further questions could be boring for children, and they may not respond properly and accurately to all questions and with fewer questions may not adequately assess all aspects of KAP on oral health. In most studies, knowledge questions were asked before the others,^{16,17} but in some studies, there was not a regular sequence to ask questions.³² It seems reasonable that the level of knowledge was measured first and then the practice and attitude on oral health.

In a suitable questionnaire, the knowledge questions should be included some items on the effects of sweets and soft drinks on the dentition, the importance of preserving natural teeth, the effect of brushing, using fluoridated toothpaste, and the importance of flossing. It is better to response the knowledge questions with both short (yes or no or true or false) and multiple choice forms depending on the type of question. The results of this review showed that despite the importance of flossing to prevent tooth decay, only a limited number of studies have asked about knowledge on using dental floss. Furthermore, in some of studies, knowledge and attitude questions were mixed together without logical procedure.³²

As mentioned earlier in this review,

knowledge questions were asked in several ways. The most important questions were about knowledge of diet effect on tooth caries. It seems that the best form of question was:

3. Do you know which of the following food items is/are the main cause of tooth decay?

- a. Sweets/chocolates/biscuits/cakes/chips/wafers
- b. Fresh fruits
- c. Raw vegetables
- d. Do not know

Although this form of the question was only used by three studies, but because the answers were multiple choice and include a variety of snacks, it appears to be more acceptable for children.

It obvious that an appropriate questionnaire should be included more questions with multiple choice answers in the area of practice. Because this area measures different domains such as brushing habits, the frequency of sugar consumption and drinking soft drinks, and dental visits. In this systematic review, some of the articles only evaluated the practice of children (7 article), and the most of other studies have focused on the practice of oral health in comparison with knowledge and attitude. It seems that the reason of this emphasis on practice is that most of the studies have been shown that the practice is not fully explained by knowledge and attitude, and the oral healthcare practice is influenced by socioeconomic factors, especially mother's education level and location where the children live (urban and rural).^{3,12} In this review, the most studies measured the TBH with questions about the frequency, duration, and brushing aides; in only two articles type of cleaning of teeth and type of brushing have been evaluated. The most important question about TBH was about the correct number of brushing. It seems that the best form of question about this subject was:

4. How often do you brush your teeth? (13 studies)

- a. Less than once a week
- b. Once a week
- c. 2-3 times a week
- d. Every day
- e. Twice or more a day
- f. Do not know/no answer.

Answers of this question were more complete and included a wider range of time than other forms of questions.

Furthermore, in a perfect questionnaire, the attitude questions should be included items about the importance of natural teeth, and fear of dental treatment and opinions about and attitudes toward dentists and dental care. It is better to response these questions as three Likert-scale questions (agree, disagree, no opinion). In this review, the attitudes of 12-year-old have been evaluated in 10 articles. It seems that although attitude about fear of the dentist and experience on a dental visit could be effective on oral health practice, less emphasis has been placed on this topic. This, may be due to the fact that, oral health attitude does not necessarily relate to better health behavior.^{12,13}

Although there are studies in Portugal, Japan, and Spain, but most studies have been conducted in developing countries. However, it can be due to effective networking within the schools, and appropriate and comprehensive school-based educational programs in developed countries. In developing countries, it appears that even in a community, for example, India that many studies have been conducted about this subject,^{3,6,8,17,19} there was not a standard questionnaire and researchers collected data by self-administered structured questionnaires.

According to the above stated, the importance of knowledge on oral health and the role of improving knowledge as a prerequisite for oral health perception and behavior, a standard questionnaire, particularly in the area of knowledge and attitude, is an urgent need. This questionnaire must be comprehensive and

should include different items in each area, but it appears that this issue has not been considered by researchers even in the different area of one country.

Therefore, it is recommended that further countrywide intensive studies should be carried out to compare national and international oral health knowledge, practice, and attitude in this age group. These studies should use an appropriate and accurate questionnaire-based on the culture of their country. Results of this systematic review can help researchers to select the best version of questions in each item and to design an appropriate questionnaire for this group.

Before drawing any conclusion, limitations of this study are worth mentioning. In this study, there was the lack of access to some of the full-text articles and in spite of attempts to contact the authors, this not achieved, so these studies were excluded from the study. Moreover, in some full-text articles, the questionnaires were not available, and the results were presented in the tables. We tried to contact the authors and get the questionnaires but did not receive a response, so the questions were extracted from the tables. Another limitation of this study was that we just searched for studies in English, and we did not have access to non-English studies.

Conclusion

It seems that, despite the importance of promotion of oral health by improving KAP in 12-year-old children, there is not a standard questionnaire. Therefore, it is absolutely necessary to design a standard questionnaire to evaluate KAP in this age group of children.

Conflict of Interests

Authors have no conflict of interest.

Acknowledgments

This work was supported by Oral and Dental Diseases Research Center AND Kerman

Social Determinants on Oral Health
Research Center, Kerman University of

Medical Sciences, Kerman, Iran.

References

1. Petersen PE, Bourgeois D, Ogawa H, Estupinan-Day S, Ndiaye C. The global burden of oral diseases and risks to oral health. *Bull World Health Organ* 2005; 83(9): 661-9.
2. Petersen PE. The World Oral Health Report 2003: continuous improvement of oral health in the 21st century--the approach of the WHO Global Oral Health Programme. *Community Dent Oral Epidemiol* 2003; 31(Suppl 1): 3-23.
3. Suprabha BS, Rao A, Shenoy R, Khanal S. Utility of knowledge, attitude, and practice survey, and prevalence of dental caries among 11- to 13-year-old children in an urban community in India. *Glob Health Action* 2013; 6: 20750.
4. Zhu L, Petersen PE, Wang HY, Bian JY, Zhang BX. Oral health knowledge, attitudes and behaviour of children and adolescents in China. *Int Dent J* 2003; 53(5): 289-98.
5. Bagramian RA, Garcia-Godoy F, Volpe AR. The global increase in dental caries. A pending public health crisis. *Am J Dent* 2009; 22(1): 3-8.
6. Mehta A, Kaur G. Oral health-related knowledge, attitude, and practices among 12-year-old schoolchildren studying in rural areas of Panchkula, India. *Indian J Dent Res* 2012; 23(2): 293.
7. World Health Organization. Oral health surveys: basic methods. Geneva, Switzerland: World Health Organization; 1997.
8. Harikiran AG, Pallavi SK, Hariprakash S, Nagesh KS. Oral health-related KAP among 11- to 12-year-old school children in a government-aided missionary school of Bangalore city. *Indian J Dent Res* 2008; 19(3): 236-42.
9. Pakshir HR. Oral health in Iran. *Int Dent J* 2004; 54(6 Suppl 1): 367-72.
10. Kaliyaperumal K. Guideline for conducting a knowledge, attitude and practice (KAP) study. *AECS Illumination* 2004; 4(1): 7-9.
11. de Almeida CM, Petersen PE, Andre SJ, Toscano A. Changing oral health status of 6- and 12-year-old schoolchildren in Portugal. *Community Dent Health* 2003; 20(4): 211-6.
12. Smyth E, Caamano F, Fernandez-Riveiro P. Oral health knowledge, attitudes and practice in 12-year-old schoolchildren. *Med Oral Patol Oral Cir Bucal* 2007; 12(8): E614-E620.
13. Diwan S, Saxena V, Bansal S, Kandpal SD, Gupta N. Oral health: knowledge and practices in rural community. *Indian Journal of Community Health* 2011; 23(1): 29-1.
14. Haleem A, Siddiqui MI, Khan AA. School-based strategies for oral health education of adolescents-a cluster randomized controlled trial. *BMC Oral Health* 2012; 12: 54.
15. Grewal N, Kaur M. Status of oral health awareness in Indian children as compared to Western children: a thought provoking situation (a pilot study). *J Indian Soc Pedod Prev Dent* 2007; 25(1): 15-9.
16. Vakani F, Basaria N, Katpar S. Oral hygiene KAP assessment and DMFT scoring among children aged 11-12 years in an urban school of Karachi. *J Coll Physicians Surg Pak* 2011; 21(4): 223-6.
17. Postgraduate Institute of Dental Sciences. Atraumatic restorative treatment to the Rural population [Online]. [cited 2006]; Available from: URL: http://new.paho.org/hq/dmdocuments/2009/OH_top_PT_rl.pdf
18. Shenoy RP, Sequeira PS. Effectiveness of a school dental education program in improving oral health knowledge and oral hygiene practices and status of 12- to 13-year-old school children. *Indian J Dent Res* 2010; 21(2): 253-9.
19. Prasai DL, Shakya A, Shrestha M, Shrestha A. Dental caries prevalence, oral health knowledge and practice among indigenous Chepang school children of Nepal. *BMC Oral Health* 2013; 13: 20.
20. Sharada A, Shetty S, Ramesh N, Sharda J, Bhat N, Asawa K. Oral Health Awareness and Attitude among 12-13 Year Old School Children in Udaipur, India. *International Journal of Dental Clinics* 2011; 3(4): 16-9.
21. Petersen PE, Jiang H, Peng B, Tai BJ, Bian Z. Oral and general health behaviours among Chinese urban adolescents. *Community Dent Oral Epidemiol* 2008; 36(1): 76-84.
22. Rajab LD, Petersen PE, Bakaeen G, Hamdan MA. Oral health behaviour of schoolchildren and parents in Jordan. *International Journal of Paediatric Dentistry* 2002; 12(3): 168-76.
23. Lian CW, Phing TS, Chat CS, Shin BC, Baharuddin LH, Che'Jalil ZB. Oral health knowledge, attitude and practice among secondary school students in Kuching, Sarawak. *Archives of Orofacial Sciences* 2010; 5(1): 9-16.
24. Kawamura M, Takase N, Sasahara H, Okada M. Teenagers' oral health attitudes and behavior in Japan: comparison by sex and age group. *J Oral Sci* 2008; 50(2): 167-74.
25. Hysi D, Droboniku E, Toti C, Xhemnica L, Petrela E. Dental caries experience and oral health behaviour among 12-year-olds in the city of Tirana, Albania. *OHDMBSC* 2010; 9(4): 229-34.
26. Gathecha G, Makokha A, Wanzala P, Omolo J, Smith P. Dental caries and oral health practices among 12 year old

- children in Nairobi West and Mathira West Districts, Kenya. *Pan Afr Med J* 2012; 12: 42.
27. Granville-Garcia AF, Fernandes LV, de Farias TSS, D'Ávila S, Cavalcanti AL, Menezes VA. Adolescents' knowledge of oral health: a population-based study. *Rev Odonto Ciênc* 2010; 25(4): 361-6.
 28. Jürgensen N, Petersen PE. Oral health and the impact of socio-behavioural factors in a cross sectional survey of 12-year old school children in Laos. *BMC Oral Health* 2009; 9: 29.
 29. de Silva-Sanigorski A, Ashbolt R, Green J, Calache H, Keith B, Riggs E, et al. Parental self-efficacy and oral health-related knowledge are associated with parent and child oral health behaviors and self-reported oral health status. *Community Dent Oral Epidemiol* 2013; 41(4): 345-52.
 30. Prasad AK, Shankar S, Sowmya J, Priyaa CV. Oral health knowledge attitude practice of school students of KSR Matriculation School, Thiruchengode. *J Indian Acad Dent Specialists* 2010; 1(1): 5-11.
 31. Varenne B, Petersen PE, Ouattara S. Oral health behaviour of children and adults in urban and rural areas of Burkina Faso, Africa. *Int Dent J* 2006; 56(2): 61-70.
 32. Ouma N, Martha A. Oral health promotion in primary schools in Mofolo, Soweto [Online]. [cited 2008 Oct 23]; Available from: URL: <http://wiredspace.wits.ac.za/jspui/handle/10539/5803>
 33. Mafuvadze BT, Mahachi L, Mafuvadze B. Dental caries and oral health practice among 12 year old school children from low socio-economic status background in Zimbabwe. *Pan Afr Med J* 2013; 14: 164.
 34. Ahmed NA, Astrom AN, Skaug N, Petersen PE. Dental caries prevalence and risk factors among 12-year old schoolchildren from Baghdad, Iraq: a post-war survey. *Int Dent J* 2007; 57(1): 36-44.
 35. Al-Omiri MK, Al-Wahadni AM, Saeed KN. Oral health attitudes, knowledge, and behavior among school children in North Jordan. *J Dent Educ* 2006; 70(2): 179-87.
 36. Petersen PE, Kwan S. Evaluation of community-based oral health promotion and oral disease prevention-WHO recommendations for improved evidence in public health practice. *Community Dent Health* 2004; 21(4 Suppl): 319-29.