



Practice and Pattern of Antenatal and Postnatal Exercise among Nigerian Women: A Cross-Sectional Study

Chidozie Emmanuel Mbada^{1*}, Olubukayomi Egunoluwa Adebayo¹, Taofeek Oluwole Awotidebe¹, Funmilola Adenike Faremi², Monisola Omoyemi Oginni², Abiola Oladele Ogundele³, Anne Antonette I. Emechete¹

Abstract

Objectives: The African culture seems to play a major prohibiting role in physical exercise during pregnancy and immediate postpartum. This study was designed to assess practice and pattern of antenatal and postnatal exercise among Nigerian women.

Materials and Methods: Three hundred and sixty-five women (189 pregnant women and 179 nursing mothers) from six selected hospitals in south-west Nigeria participated in this cross-sectional study. Data were obtained on socio-demographic and obstetric characteristics, and practice of antenatal and postnatal exercise. Descriptive and inferential statistics were used to analyze data at $P < 0.05$ alpha level.

Results: The mean age of all the respondents was 28.9 ± 4.63 years. A majority of the pregnant women responders were nulliparous (39.2%) while the nursing mother responders were mostly multiparous (54.5%). Prevalence for engagement in physical exercise was 81.9% and mostly based on self-prescription (35.5%) and prescription by nurses (28.8%). Antenatal and postnatal exercise practice was 84.7% (160/189) and 79.0% (139/176), respectively. Aerobic (43.5%) and stretching (33.4%) were the most common type of physical exercises. Exercise frequency was mostly 1-2 times per week (40.8%). There was significant association between exercise practice and education level ($\chi^2 = 18.795$; $P = 0.001$).

Conclusion: Engagement in antenatal and postnatal exercise among Nigerian pregnant and nursing women was high and mostly based on self-prescription. Education level significantly influence physical exercise practice among Nigerian women with aerobic and stretching as the predominant exercises in pregnancy and postpartum.

Keywords: Exercise, Nigeria, Pregnancy, Postpartum

Introduction

Exercise has become an essential aspect of antenatal and postnatal care in most settings (1-3). Pervading skepticism about safety and benefit of exercises in pregnancy and postpartum seems to be giving way to populating evidence that relevant exercise are safe and of significant benefit to the mother and child (1,3-5). Explicitly, empirical evidence revealed the benefits of exercise to include preservation and promotion of aerobic and musculoskeletal fitness levels, improvement in posture, coordination and balance, increase in endurance and stamina, deterrence of extreme maternal weight gain, gestational glucose control, improved psychological adjustment to changes in pregnancy and postpartum, facilitation of labour and more rapid postnatal recovery (4-9).

Consequent to the foregoing, the American College of Obstetricians and Gynecologists recommended low-impact, moderate-intensity and regular exercises for pregnant women irrespective of their physical fitness level for approximately 30 minutes on most days of the week.

However, there are empirical reports that women are yet to attain these recommendations (10,11). Rather, there are quantity of data reporting decreased physical activity in early pregnancy (12-15), late pregnancy (14-16) and during the postpartum period (17-19). Lack of engagement in physical exercise during pregnancy and postpartum period have been implicated on a gamut of factors which may not be mutually exclusive. There is still some conservative view and myths that exercise is too delicate and unsafe for pregnant women. On the other hand, women are afraid to participate in physical exercise owing to fear that it will negatively impact on their health (5,20). In addition, increase caring responsibility associated with pregnancy and postpartum and cultural perceptions are determinants of physical exercise in women. Previous investigators have reported prevalence and pattern of physical exercise among pregnant women from the developed countries (12,13,21-24). Anecdotally, the African culture seems to play a prohibiting role in physical exercises during pregnancy and immediate postpartum by man-

Received 9 December 2014, Accepted 27 January 2015, Available online 20 February 2015

¹Department of Medical Rehabilitation, College of Health Sciences, Obafemi Awolowo University, Ile-Ife, Nigeria. ²Department of Nursing Science, College of Health Sciences, Obafemi Awolowo University, Ile-Ife, Nigeria. ³Department of Physiotherapy, Obafemi Awolowo University Teaching Hospitals Complex, Ile-Ife, Nigeria.

*Corresponding author: Chidozie E. Mbada, Department of Medical Rehabilitation, College of Health Sciences, Obafemi Awolowo University, Ile-Ife, Nigeria. Tel: +234802825254, Email: doziembada@yahoo.com



dating confinement periods that varies across different tribes. However, there is an apparent dearth of such studies among African women (25). This study was designed to assess the practice and pattern of antenatal and postnatal exercise among Nigerian women.

Materials and Methods

A total of 365 (189 pregnant women and 179 nursing mothers) out of 400 women that were approached, responded in this cross-sectional study, therefore yielding a response rate of 91.3%. Consecutive sampling was used to recruit respondents from six selected hospitals namely Urban Comprehensive Health Centre (UCHC), Enuwa Primary Health centre, Comprehensive Health centre Aderemi, Obafemi Awolowo University (OAU) Health Centre, OAU Teaching Hospitals Complex (OAUTHC) and Seventh Day Adventist Hospital in Ile-Ife, Osun state, south west Nigeria. Informed consent of all respondents was required for participation in the study.

A structured Knowledge, Attitude and Practice (KAP) questionnaire containing pre-coded and open questions by Walsh et al. (26) was adapted and used for data collection in this study. The content validity of the adapted version of the questionnaire was subjected to expert reviews. The reliability of the Yoruba version of the questionnaire was assessed in a pilot study by a test-retest method among 10 respondents attending the UCHC of the OAUTHC, observing 7 days between test and re-test. The summation of all the checked items on the questionnaire at test and re-test were compared. The questionnaire items yielded an agreement percentage that ranged from 87.4% to 99.6%, the intra-class coefficient was 0.985 and the confidence interval ranged from 0.94% to 0.996. Respondents who were not literate in either English or Yoruba were excluded from the study. Yoruba is the local language spoken in the area where the study was conducted. This self-administered questionnaire sought information on socio-demographic, maternal infant characteristics and practice of antenatal and postnatal exercise. Respondents were asked to recall on their current antenatal or postnatal exercise pattern. Between 10 and 15 minutes was required to complete the questionnaire.

Data Analysis

Descriptive statistics of mean, standard deviation, frequency distribution were used to summarize data. Inferential statistics of Chi-square test was used to test the association between exercise practice (antenatal and postnatal) and individual factors. Alpha level was set at 0.05. Data was analyzed using SPSS version 16.0 (SPSS Inc., Chicago, USA).

Results

The mean age of all the respondents was 28.9±4.63 years. The socio-demographic characteristics of respondents are presented in Table 1. A majority of the respondents were Christians (76.4%), business women (39.9%), had tertiary education (67.9%) and were within the level of income of

\$100 to \$200 per month (29.6%).

The maternal and obstetrics characteristics of respondents are presented in Table 2. A majority of the pregnant women responders were nulliparous (39.2%) while the nursing mother responders were mostly multiparous (54.5%). Spontaneous vaginal delivery was the most reported mode of delivery among the pregnant (57.7%) and nursing (92.6%) women, respectively. The parous respondents mostly utilized hospitals as place of delivery (68.0%) and mostly commenced antenatal care between 3 and 6 months of pregnancy (32.1%).

Prevalence, frequency and prescriptors of antenatal and postnatal exercise are presented in Table 3. The total prevalence for engagement in physical exercise among the respondents was 81.9%. The prevalence of antenatal and postnatal exercise practice was 84.7% and 79.0%, respectively. Exercise frequency was mostly 1-2 times per week (40.8%). Physical exercise engagement based on self-prescription (35.5%) (i.e. engagement in exercise for therapeutic purpose without medical advice) and nurse-prescription (prescribing of exercises by nurses) by (28.8%) were mostly reported.

The types of exercise practiced by respondents are presented in Table 4. Aerobic (43.5%) and stretching (33.4%) were the most common type of physical exercise practiced by the respondents. On the other hand, cycling (4.3%) and swimming (4.7%) were scarcely practiced among the re-

Table 1. Socio-demographic characteristics of the respondents

Variable	ANR (n=189) No. (%)	PNR (n=176) No. (%)	Total (N=365) No. (%)
Religion			
Christianity	145 (76.7)	134 (76.1)	279 (76.4)
Islam	43 (22.8)	42 (23.9)	85 (23.3)
Traditional religion	1 (0.5)	0 (0)	1 (0.3)
Occupation			
Home maker	23 (12.2)	23 (13.1)	46 (12.6)
Trading/business	103 (54.5)	79 (44.9)	184 (39.9)
Civil servant	39 (20.6)	44 (25.0)	83 (22.7)
Student	20 (10.6)	21 (11.9)	41 (11.2)
Not specified	4 (2.1)	9 (5.1)	13 (3.6)
Educational qualification			
Primary	9 (4.7)	12 (6.8)	21 (5.8)
Secondary	49 (25.9)	47 (26.7)	96 (26.3)
Tertiary	131 (69.3)	117 (66.5)	248 (67.9)
Family Setting			
Polygamy	38 (20.1)	46 (26.1)	84 (23.0)
Monogamy	125 (66.1)	117 (66.5)	242 (66.3)
Single Parenting	26 (13.8)	13 (7.4)	39 (10.7)
Income			
Less than \$100	32 (16.9)	27 (15.3)	59 (16.2)
\$100-\$200	51 (27.0)	57 (32.4)	108 (29.6)
\$200-\$300	38 (20.1)	45 (25.6)	83 (22.7)
\$300-\$500	13 (6.9)	15 (8.5)	28 (7.7)
\$500-\$1000	10 (5.3)	4 (2.3)	14 (3.8)
Greater than \$1000	10 (5.3)	5 (2.8)	15 (4.1)
Non Disclosure	25 (13.2)	33 (18.8)	58 (15.9)

ANR: Antenatal Respondents; PNR: Postnatal Respondents

Table 2. Maternal and obstetric characteristics of the respondents

Variable	ANR (n=189) No. (%)	PNR (n=176) No. (%)	Total (N=365) No. (%)
Parity			
Nulliparous	74 (39.2)	0 (0)	74 (20.3)
Primiparous	53 (28.0)	80 (45.5)	133 (36.4)
Multiparous	62 (32.8)	96 (54.5)	158 (43.3)
Mode of delivery			
Vaginal	109 (57.7)	163 (92.6)	272 (74.5)
Caesarean Section	6 (3.2)	13 (7.4)	19 (5.2)
Not Applicable	74 (39.2)	0 (0)	74 (20.3)
Place of delivery			
Hospital	93 (49.2)	155 (88.0)	248 (68.0)
Home	7 (3.7)	5 (2.8)	12 (3.3)
Mission	15 (7.9)	16 (9.2)	31 (8.5)
Not applicable	74 (39.2)	0 (0.0)	74 (20.2)
Previous antenatal care start time			
< 1 month	5 (2.7)	25 (14.2)	30 (8.2)
1–3 months	54 (28.6)	49 (27.8)	103 (28.2)
3–6 months	44 (23.2)	73 (41.5)	117 (32.1)
6–9 months	12 (6.3)	29 (16.5)	41 (11.2)
Not applicable	74 (39.2)	0 (0.0)	74 (20.3)
Present antenatal care duration			
< 1 month	75 (39.7)	0 (0.0)	75 (20.5)
1–3 months	57 (30.2)	0 (0.0)	57 (15.6)
3–6 months	44 (23.3)	0 (0.0)	44 (12.1)
6–9 months	13 (6.9)	0 (0.0)	13 (3.6)
Not applicable	0 (0)	176 (100)	176 (48.2)

ANR: Antenatal Respondents; PNR: Postnatal Respondents

Table 3. Prevalence, frequency and prescriptors of antenatal and postnatal exercise

Variable	Antenatal No. (%)	Postnatal No. (%)	Total No. (%)
Prevalence of exercise			
Yes	160 (84.7)	139 (79.0)	299 (81.9)
No	29 (15.3)	37 (21.0)	66 (18.1)
Frequency of exercise practice/week			
1–2 times	65 (40.6)	57 (41.0)	122 (40.8)
3–4 times	49 (30.6)	52 (37.4)	101 (33.8)
> 4 times	46 (28.8)	30 (21.6)	76 (25.4)
Prescriptor of exercise			
Doctor			
Yes	26 (16.3)	23 (16.5)	49 (16.4)
No	134 (83.7)	116 (83.5)	250 (83.6)
Nurse			
Yes	45 (28.1)	41 (29.5)	86 (28.8)
No	115 (71.9)	98 (70.5)	213 (71.2)
Physical therapist			
Yes	56 (35.0)	27 (19.4)	83 (27.8)
No	104 (65.0)	112 (80.6)	216 (72.2)
Spouse			
Yes	7 (4.4)	10 (7.2)	17 (5.7)
No	153 (95.6)	129 (92.8)	282 (94.3)
Self			
Yes	52 (32.5)	54 (38.8)	106 (35.5)
No	108 (67.5)	85 (61.2)	193 (64.5)

Table 4. Types of antenatal and postnatal exercise practiced by respondents (n=299)

Variable	ANR (n=160) No. (%)	PNR (n=139) No. (%)	Total (N=299) No. (%)
Aerobics			
Yes	65(40.6)	65(46.8)	130(43.5)
No	95(59.4)	74(53.2)	169(56.5)
Abdominal exercise			
Yes	40(25.0)	23(16.5)	63(21.1)
No	120(75.0)	116(83.5)	236(78.9)
Pelvic floor exercise			
Yes	22(13.7)	33(23.7)	55(18.4)
No	138(86.3)	106(76.3)	244(81.6)
Back care exercise			
Yes	28(17.5)	20(12.5)	48(16.1)
No	132(82.5)	119(87.5)	251(83.9)
Swimming			
Yes	8(5.0)	6(4.3)	14(4.7)
No	152(95.0)	133(95.7)	285(95.3)
Cycling			
Yes	5(3.1)	8(5.8)	13(4.3)
No	155(96.9)	131(94.2)	286(95.7)
Stretching exercise			
Yes	57(35.6)	43(30.9)	100(33.4)
No	103(64.4)	96(69.1)	199(66.6)
Relaxation and breathing exercise			
Yes	42(26.3)	30(21.6)	72(24.1)
No	118(73.7)	109(78.4)	227(75.9)
Muscle strengthening exercise			
Yes	36(22.5)	40(28.8)	76(25.4)
No	124(77.5)	99(71.2)	223(74.6)

ANR: Antenatal Respondents; PNR: Postnatal Respondents

spondents.

Chi-square test of association between exercise practice and socio-demographic and obstetric characteristics is presented in Table 5. The result showed a significant direct association between exercise and education level ($\chi^2=18.795$; $P=0.001$) with respondents with tertiary education level reporting higher exercise practice (72.9%).

Discussion

This study assessed practice and pattern of antenatal and postnatal exercise among Nigerian women. The study found a high prevalence of engagement in physical exercise among Nigerian pregnant and nursing women. However, comparison of this study's finding is limited by apparent dearth of data on level of engagement in exercise among Nigerian pregnant and nursing women over time. Nonetheless, this finding may be an indication of a paradigm shift in attitude towards exercise in pregnancy and postpartum period. Anecdotal observation in the study environment is in agreement with previous submissions that acceptance of modern antenatal care, engagement in exercise and delivery care among pregnant women from Africa is inhibited by cultural myths, traditional beliefs and misperceptions (27-30).

Many cultures in Africa, based on fatalistic views observe

Table 5. Chi-square test of association between exercise practice and socio-demographic and obstetric characteristics

Variable	Exercise practice		χ^2	P-value
	Yes, Number (%)	No, Number (%)		
Age group			1.577	0.209
<30 years	179 (59.9)	45 (68.2)		
≥30 years	120 (40.1)	21 (31.8)		
Education			18.795	0.001
Primary	15 (5)	6 (9.1)		
Secondary	66 (22.1)	30 (45.5)		
Tertiary	218 (72.9)	30 (45.4)		
Occupation			9.134	0.058
Home maker	37 (12.4)	9 (13.6)		
Trading/business	148 (49.5)	34 (51.5)		
Civil servant	74 (24.7)	9 (13.6)		
Student	28 (9.4)	13 (19.7)		
Not specified	12 (4)	1 (1.5)		
Parity			0.157	0.925
Nulliparous	61 (20.4)	13 (19.7)		
Primiparous	110 (36.8)	23 (34.8)		
Multiparous	128 (42.8)	30 (45.5)		

* Significance at $\alpha = 0.05$

taboos during pregnancy and postpartum. However, socio-economic conditions seem to have diminished the practice of seclusion and confinement during pregnancy. Nonetheless, during postpartum, confinement periods up to 40 days or more are still enforced by mother-in-law and elderly women in the community to protect the mother and infant from disease and evil spirit. During the confinement periods, mothers are often put on long bed rest and feeding on certain diets in order to restore the contraction of the uterus, dry the womb, reduce postpartum haemorrhage, establish milk production and healing of the perineum or caesarean section incision. However, these mothers are restrained from physical activity and exercises based on beliefs that exercises are contra-indicated and alter the formation of new blood. In contrast, empirical finding from this study revealed high physical exercise practice rate among the women. However, the finding on prevalence of engagement in physical exercise among women in this study should be treated with caution because the respondents in study comprised of women attending antenatal or postnatal clinics who may have imbibed health promotion and education on importance of exercises as component of antenatal and postnatal care. It is therefore advocated that a community-based survey may be needed to validate the finding of this hospital-based survey.

This study found that most of the women engaged in physical exercise based on self-prescription. Self-prescription of exercise in pregnancy and postpartum portend a negative dimension to a positive development. Self-prescription of exercise in pregnancy and postpartum may be associated with the availability of wide range of information emanating from family, friends, print materials, internet and the social media. Most of this information is confusing, unscientific and lacking in specificity. On the other hand, self-prescription of exercise in pregnancy and post-

partum may be as a result of non-availability of competent and qualified exercise experts such as a women's health or obstetric physical therapist. Furthermore, the result of this study showed that more women engaged in physical exercise based on prescription by nurses than physical therapists. This finding confirms the non-availability of specialist physical therapists in most of the selected hospitals where this study was conducted. Nurses and midwives have been reported as the commonest information sources on exercise in pregnancy and postpartum in other studies (31,32). Specifically, Whitford et al. (33) in a study in north-east Scotland, found that midwives were the health professionals most likely to give information on pelvic floor exercises. Although, physicians, nurses, midwives and community health aides often prescribe exercises in some settings, may not possess the requisite knowledge of the principles, prescription, indications and contraindications to exercise in pregnancy and postpartum. Competent supervision is significant for maintaining quality of exercise. Studies have shown that physical therapists' supervised exercise in pregnancy and postpartum resulted in improved outcomes (33-35). However, healthcare professionals prescribing exercise in pregnancy and postpartum should seek knowledge of the potential risks and basic tenets of exercise prescription in health and disease.

This study found aerobic and stretching as the predominant physical exercise types in pregnancy and postpartum. This study corroborates previous findings that have found aerobic and stretching exercises as simple, easy to do and adequate in pregnancy (4,36-39). Although, cycling and swimming have quantity of support in literature as ideal exercises in pregnancy and postpartum, they were scarcely practiced among the women in this study. Swimming is far remote as a way of life of the people in the study environment and there is lack of swimming skills and limited availability of swimming pools. This is further complicat-

ed by prevalent hydrophobia and superstitions that make swimming a taboo for pregnant and nursing women. Furthermore, non-availability or non-affordability of bicycle ergometer for exercise may have contributed to low rate of cycling exercise among the women in this study.

Lastly, this study found significant association between exercise practice and education level. Women with higher level of education were more involved in exercise in pregnancy and postpartum. This finding is consistent with some previous reports that have associated higher participation in physical activity or exercise with higher education (12,14,24,40,41). Petersen et al. (42) in a study on correlates of physical activity among pregnant women in the United States found that pregnant women meeting the moderate or vigorous physical activity recommendations were more likely to be younger, non-Hispanic white, more educated, not married, nonsmokers, and to have higher incomes. Gaston and Cramp (43) in a review on exercise in pregnancy found higher education and income as predictors of higher exercise participation during pregnancy. The psychometric properties of the original instrument from which the questionnaire in this study was adapted was not reported by the authors. Therefore, lack of information on validity and reliability of the original instrument and the use of the same questionnaire for a heterogeneous sample is a potential limitation of this present study. However, the findings of this study were presented under the pregnant and nursing women dichotomy and thus reduce possible influence of homogeneity of sample.

Conclusion

Nigerian pregnant and nursing women demonstrate high engagement in physical exercises which were usually by self-prescription. Higher education is a significant determinant of exercise participation during pregnancy among Nigerian women. The predominant exercise types in pregnancy and postpartum were aerobic and stretching.

Ethical issues

Ethical approval for the study (IPHOAU/12/13) was obtained from the Health Research Ethics Committee of the Institute of Public Health, OAU, Ile-Ife, Nigeria.

Financial support

No grant was received to conduct this study. However, the ACU Early Career Travel Grant was received towards dissemination of the findings of this study.

Conflict of interests

Authors declare no conflict of interests.

Acknowledgments

The authors gratefully acknowledge all respondents, and the administrative and nursing staff of the different hospitals selected for this survey. The Early Careers Grants offered by Association of Commonwealth Universities (ACU) towards the presentation of part of the findings of this study is immensely appreciated.

References

1. ACOG Committee Obstetric Practice. ACOG Committee opinion. Number 267, January 2002: exercise during pregnancy and the postpartum period. *Obstet Gynecol* 2002; 99(1):171-3.
2. Davies GA, Wolfe LA, Mottola MF, MacKinnon C, Arsenault MY, Bartellas E, et al. Exercise in pregnancy and the postpartum period. *J Obstet Gynaecol Can* 2003;25(6):516-29.
3. Wadsworth P. The benefits of exercise in pregnancy. *J Nurse Practitioners* 2007;3(5):333-9.
4. Artal R, O'Toole M. Guidelines of the American College of Obstetricians and Gynecologists for exercise during pregnancy and the postpartum period. *Br J Sports Med* 2003;37(1):6-12.
5. Shelby S. Medical report: Exercise in the postpartum period. *ACSM'S Health Fitness J* 2006;10(4):40-1.
6. Artal R, Platt LD, Sperling M, Kammula RK, Jilek J, Nakamura R. Maternal cardiovascular and metabolic responses in normal pregnancy. *Am J Obstet Gynecol* 1981;140(2):123-7.
7. Clapp JF III. Exercise in pregnancy: a brief clinical review. *Fetal Med Rev* 1990;161:1464-9.
8. Lokey EA, Tran ZV, Wells CL, Myers BC, Tran AC. Effects of physical exercise on pregnancy outcomes: a meta-analytic review. *Med Sci Sports Exerc* 1991;23(11):1234-9.
9. Sternfeld B, Quesenberry CP Jr, Eskenazi B, Newman LA. Exercise during pregnancy and pregnancy outcome. *Med Sci Sports Exerc* 1995;27(5):634-40.
10. Soultanakis HN, Artal R, Wiswell RA. Prolonged exercise in pregnancy: glucose homeostasis, ventilatory and cardiovascular responses. *Semin Perinatol* 1996;20(4):315-27.
11. Artal R, Sherman C. Exercise during pregnancy. Safe and beneficial for most. *Phys Sportsmed* 1999; 27(8):5-60. doi: 10.3810/psm.1999.08.947.
12. Evenson KR, Savitz DA, Huston SL. Leisure-time physical activity among pregnant women in the US. *Paediatr Perinat Epidemiol* 2004;18(6):400-7.
13. Pivarnik JM, Chambliss HO, Clapp JF, Dugan SA, Hatch MC, Lovelady CA, et al. Impact of physical activity during pregnancy and postpartum on chronic disease risk. *Med Sci Sports Exerc* 2006;38(5):989-1006.
14. Mottola MF, Campbell MK. Activity patterns during pregnancy. *Can J Appl Physiol* 2003;28(4):642-53.
15. Pereira MA, Rifas-Shiman SL, Kleinman KP, Rich-Edwards JW, Peterson KE, Gillman MW. Predictors of change in physical activity during and after pregnancy. *Am J Prev Med* 2007;32(4):312-9.
16. Borodulin KM, Evenson KR, Wen F, Herring AH, Benson AM. Physical activity patterns during pregnancy. *Med Sci Sports Exerc* 2008;40(11):1901-8. doi: 10.1249/MSS.0b013e31817f1957.
17. Fell DB, Joseph KS, Armson BA, Dodds L. The impact of pregnancy on physical activity level. *Maternal Child Health J* 2009;13(5):597-603. doi: 10.1007/s10995-008-0404-7.
18. Schmidt MD, Pekow P, Freedson PS, Markenson G, Chasan-Taber L. Physical activity patterns during

- pregnancy in a diverse population of women. *J Women Health* 2006;15(8):909-18.
19. van Raaij JM1, Schonk CM, Vermaat-Miedema SH, Peek ME, Hautvast JG. Energy cost of physical activity throughout pregnancy and the first year postpartum in Dutch women with sedentary lifestyles. *Am J Clin Nutr* 1990;52(2):234-9.
 20. Blum JW, Beaudoin CM, Caton-Lemos L. Physical activity patterns and maternal well-being in postpartum women. *Matern Child Health J* 2004; 8(3):163-9.
 21. Treuth MS, Butte NF, Puyau M. Pregnancy-related changes in physical activity, fitness, and strength. *Med Sci Sports Exerc* 2005;37(5):832-7.
 22. Women and Physical Activity [database on the Internet]. 2010 [cited 24/01/2014]. Available from: <http://www.whv.org.au>
 23. Hegaard HK, Damm P, Hedegaard M, Henriksen TB, Ottesen B, Dykes AK, et al. Sports and leisure time physical activity during pregnancy in nulliparous women. *Matern Child Health J* 2011;15(6):806-13. doi: 10.1007/s10995-010-0647-y.
 24. Domingues MR, Barros AJ. Leisure-time physical activity during pregnancy in the 2004 Pelotas Birth Cohort Study. *Rev Saude Publica* 2007;41(2):173-80.
 25. Liu J, Blair SN, Teng Y, Ness AR, Lawlor DA, Riddoch C. Physical activity during pregnancy in a prospective cohort of British women: results from the Avon longitudinal study of parents and children. *Eur J Epidemiol* 2011;26(3):237-47. doi: 10.1007/s10654-010-9538-1.
 26. Walsh JM, McGowan C, Byrne J, McAuliffe FM. Prevalence of physical activity among healthy pregnant women in Ireland. *Int J Gynaecol Obstet* 2011;114(2):154-5. doi: 10.1016/j.ijgo.2011.02.016.
 27. Adeniyi AF, Ogwumike OO, Bamikefa TR. Postpartum exercise among Nigerian women: Issues relating to exercise performance and self-efficacy. *ISRN Obstet Gynecol* 2013;2013:294518. doi: 10.1155/2013/294518.
 28. Ribeiro CP, Milanez H. Knowledge, attitude and practice of women in Campinas, São Paulo, Brazil with respect to physical exercise in pregnancy: a descriptive study. *Reprod Health* 2011;8:31. doi: 10.1186/1742-4755-8-31.
 29. Logan C. Traditional beliefs and practices of pregnancy and childbirth. *Botsw Natl Health Bull* 1985;1(1):64-75.
 30. Krans EE, Chang JC. Low-income African American women's beliefs regarding exercise during pregnancy. *Matern Child Health J* 2012;16:1180-7.
 31. Igbere GO. Harmful cultural practices and reproductive health in Nigeria. *Continental J Tropic Med* 2012;6(1):27-33.
 32. Omobola OC. An overview of taboo and superstition among the Yoruba of Southwest of Nigeria. *Mediterranean Journal of Social Sciences* 2013;4(2):221-6.
 33. Whitford HM, Alder B, Jones M. A cross-sectional study of knowledge and practice of pelvic floor exercises during pregnancy and associated symptoms of stress urinary incontinence in North-East Scotland. *Midwifery* 2006;23(2):204-17.
 34. Hermansen IL, O'Connell BO, Gaskin CJ. Women's explanations for urinary Incontinence, their management strategies, and their quality of life during the postpartum period. *J Wound Ostomy Continence Nurs* 2010;37(2):187-92. doi: 10.1097/WON.0b013e3181cf7946.
 35. Abdulaziz K, Hasan T. Role of pelvic floor muscle therapy in obese perimenopausal females with stress incontinence: A randomized control trial. *Internet J Radiology* 2012;14 (2). Available from: <https://ispub.com/IJRA/14/2/14430>
 36. Robledo-Colonia AF, Sandoval-Restrepo N, Mosquera-Valderrama YF, Escobar-Hurtado C, Ramírez-Vélez R. Aerobic exercise training during pregnancy reduces depressive symptoms in nulliparous women: a randomised trial. *J Physiother* 2012;58(1):9-15. doi: 10.1016/S1836-9553(12)70067-X.
 37. Kasawara KT, Burgos CS, do Nascimento SL, Ferreira NO, Surita FG, Pinto E Silva JL. Maternal and perinatal outcomes of exercise in pregnant women with chronic hypertension and/or previous preeclampsia: A randomized controlled trial. *ISRN Obstet Gynecol* 2013;2013:857047. doi: 10.1155/2013/857047.
 38. Wolfe LA, Hall P, Webb KA, Goodman L, Monga M, McGrath MJ. Prescription of aerobic exercise during pregnancy. *Sports Med* 1989;8(5):273-301.
 39. Koltyn KF, Schultes SS. Psychological effects of an aerobic exercise session and a rest session following pregnancy. *J of Sports Med Phys Fitness* 1997;37(4):287-91.
 40. Larson-Meyer DE. Effect of postpartum exercise on mothers and their offspring: a review of the literature. *Obes Res* 2002;10(8):841-53.
 41. Davies GA, Wolfe LA, Mottola MF, MacKinnon C, Society of Obstetricians and gynecologists of Canada, SOGC Clinical Practice Obstetrics Committee. Joint SOGC/CSEP clinical practice guideline: exercise in pregnancy and the postpartum period. *Can J Appl Physiol* 2003;28(3):330-41.
 42. Petersen AM, Leet TL, Brownson RC. Correlates of physical activity among pregnant women in the United States. *Med Sci Sports Exerc* 2005;37(10):1748-53.
 43. Gaston A, Cramp A. Exercise during pregnancy. A review of patterns and determinants. *J Sci Med Sport* 2011;14(4):299-305.

Copyright © 2015 The Author(s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.