



Socioeconomic Determinants of Exposure to Secondhand Smoke Among Pregnant Women

Ahmad Saeed Alghamdi¹, Hazem Faisal Jokhadar¹, Ibraheem Mohammed Alghamdi^{1*}, Saleh Abdullah Alsohibani¹, Odai Jamaan Alqahtani¹, Hayfaa Abdelmageed Wahabi²

Abstract

Objectives: Exposure to secondhand smoke (SHS) constitutes major health problems that threaten human health around the world; its adverse effects can lead to serious illnesses in both the pregnant women and the growing fetus. The purpose of this study was to determine the prevalence of SHS among Saudi pregnant women in King Khalid University Hospital (KKUH) and to determine the association between the socioeconomic characteristics of the pregnant women and exposure to SHS.

Materials and Methods: This was a cross-sectional study conducted in the antenatal clinic, KKUH. Data were collected from consecutive women using a predesigned questionnaire addressing the participants' demographics and exposure to SHS status.

Results: The prevalence of SHS exposure among pregnant women was 24%. Low maternal education level and being unemployed were associated with more exposure to SHS ($P=0.039$) and ($P=0.02$), respectively. There was no association between maternal age, parity and monthly income and exposure to SHS ($P=0.697$), ($P=0.420$) and ($P=0.543$), respectively.

Conclusion: Maternal educational levels and unemployment are associated risk factors for SHS exposure during pregnancy.

Keywords: Passive smoking, Pregnant women, Saudi Arabia, Secondhand smoke, Socioeconomic factors

Introduction

Exposure to secondhand smoke (SHS) constitutes major health problems that threaten human health around the world (1). It is a preventable cause of morbidity and mortality (1). Its adverse effects can lead to serious illnesses in both the pregnant women and the growing fetus (1). During pregnancy SHS exposure can lead to long lasting effects on the infant's respiratory, cardiovascular, central nervous system and other systemic abnormalities (1).

More than 600 000 premature deaths every year are caused by exposure to SHS (1). About 40% of all children, 33% of male non-smokers and 35% female non-smokers are exposed to SHS indoors (1). 31% of the deaths caused by SHS occur in children (1).

In Saudi Arabia, the prevalence of active smoking among males and females ranges according to the age groups with median of 26.5% and 9%, respectively (2). Although according to previous study, active smoking among pregnant women was low 0.08%, exposure to SHS is reported to be as high as 31% (3).

The demographic profile of pregnant women exposed to SHS showed that the exposed participants were younger in age, of low parity and lower level of education than non-exposed (3). However, the study did not show the socioeconomic determinants of SHS exposure among Saudi pregnant women (3).

The aims of this study were to determine the prevalence of SHS exposure among pregnant women and to explore

the socio-demographic determinants of exposure to SHS among Saudi pregnant women attending the antenatal care clinic in King Khalid University Hospital (KKUH).

Materials and Methods

This was prospective cross-sectional study designed to investigate the socioeconomic determinants of SHS exposure during pregnancy and to evaluate the pregnant women's knowledge about the harmful effects of SHS.

The study was done in an antenatal outpatient clinic at KKUH, Riyadh, Saudi Arabia. KKUH is a tertiary referral center and includes a neonatal intensive care unit (NICU) and an in vitro fertilization unit. The obstetrics department provides services for 3500 to 4000 deliveries per year. For this study, we invited pregnant women from KKUH antenatal care clinic from October 2013 to March 2014 to participate. The women were invited irrespective of parity or gestational age.

The prevalence rate for SHS exposure during pregnancy was estimated to be 31% from a recently published study (3). Based on that prevalence the sample size was calculated with 95% CI and 5% precision. The minimum sample size needed was 328 participants.

A consecutive sampling technique was used. The inclusion criteria were: pregnant women aged 18 years and older who consented to participate in the study. Exclusion criteria were women who actively smoked and those who declined to participate in the study.

Received 20 December 2015, Accepted 23 February 2016, Available online 10 March 2016

¹College of Medicine, King Saud University, Riyadh, Kingdom of Saudi Arabia. ²Department of Family and Community Medicine, King Saud University, Riyadh, Kingdom of Saudi Arabia.

*Corresponding author: Ibraheem Mohammed Alghamdi, College of Medicine, King Saud University, Riyadh, Kingdom of Saudi Arabia. Tel: +966560460516, Email: ibraheemalghamdi92@gmail.com



A questionnaire was used for data collection. The questionnaire was designed to assess the demographic characteristics and SHS exposure of the participants (Appendix 1). The first section of the questionnaire concerned the demographic characteristics of the participants. In this section, closed-ended questions were used in addition to open-ended ones. For instance, open-ended questions were used when asking about age and nationality, while closed-ended questions were used for educational level, marital status, parity, income, and employment. The next section was designed to determine the prevalence of SHS. We calculated the number of women who were exposed to SHS as a percentage from the total study population. We then determined, as a secondary outcome, the prevalence of active smoking among pregnant women in addition to quantifying the amount of SHS according to the hours of exposure and the number of smokers in the house.

Statistical Analysis

The statistical analysis for this study was performed using SPSS, version 21.0 (SPSS Inc., Chicago, IL, USA). Descriptive statistics were computed to assess the exposure to SHS during pregnancy. Univariate analysis was applied to assess the association between exposure to SHS and the socio-demographic characteristics by comparing these characteristics on exposed and non-exposed participants in the following; age, parity, educational level, employment, and monthly income. Chi-squared test used to compare dichotomous outcomes (unemployment, monthly income, educational level and parity), while independent *t* test used to compare continuous variables (age). Statistical significance was determined with a *P* value of <0.05.

Results

Three hundred sixty-three pregnant women were invited to participate in this study; 328 of whom were found to be eligible, 33 declined to participate and 2 were excluded according to our exclusion criteria.

The mean age of the participants was 29± 7.39 years, and most were Saudis 303 (92.4%). Of the participants 234 (72.3%) were multiparous and 139 (42.8%) had a bachelor's degree, while only 2 (0.6%) were illiterate. 113 (34.9%) were employees. The main mode of transportation was in a private car 165 (87.3%). The majority of the women's families fell under moderate- to high-income levels 243 (81.5%). The demographic characteristics are shown in Table 1.

Participants were asked to determine the hours of SHS exposure at different locations. The results presented in Table 2. It shows the number of total responses to questions related to SHS exposure at home, at work and while in their transportation vehicle. While 122 (37.3%) of the respondents reported having active smokers in their residence, only 79 (24%) were actually exposed, as shown in Table 2. Thirty-four (10.5%) of the exposed participants were exposed to 5-20 cigarettes per day at home and only 15 (4.6%) were exposed at work. There were 40 (12.3%)

that were exposed to at least once or twice a day and 31 (9.5%) were exposed in their transportation vehicle.

The 122 pregnant women who reported having active smokers at home were compared to the women who were not exposed to SHS. There was no significant association between SHS exposure and age (*P*=0.697), parity (*P*=0.420) and monthly income of the family in the study sample (*P*=0.543).

However, the study showed evidence of association between the exposure of pregnant women to SHS and low educational levels (*P*=0.039). Employed women were less likely to be exposed to SHS than unemployed women (*P*=0.02) as shown in Table 3.

Discussion

We found the prevalence of exposure to SHS among the study participants to be 24%, though 37% of them have active smokers at home. This prevalence constitutes a health problems to the pregnant women generally and specifically to the growing fetus. According to the literature SHS exposure has been approved to be a carcinogen, it can cause lung cancer and cardiovascular diseases

Table 1. Demographic Characteristics of the Study Participants

Characteristic	Saudi Arabia n = 328 (%)
Age	305 ^a
Mean (standard deviation)	29.86 years (7.4)
Nationality	328
Saudi	303 (92.4)
Non-Saudi	24 (7.3)
Marital status	328
Married	322 (98.2)
Divorced	6 (1.8)
Number of previous pregnancies	324 ^a
Zero	90 (20.8)
1-5 times	204 (63.0)
More than 5 times	30 (9.3)
Level of education	325 ^a
Primary school	12 (3.7)
Intermediate school	19 (5.8)
High school	79 (24.3)
Undergraduate	48 (14.8)
Bachelor	139 (42.8)
Postgraduate	26 (8.0)
Illiterate	2 (0.6)
Employment	324 ^a
Yes	113 (34.9)
No	211 (65.1)
Transportation ^b	189 ^a
Private car	165 (87.3)
Taxi	14 (7.41)
Bus	10 (5.29)
Monthly income	298 ^a
Less than 4999 Saudi Riyal	55 (18.5)
5000–9999 Saudi Riyal	142 (47.7)
10000–14999 Saudi Riyal	76 (25.5)
≥15000 Saudi Riyal	25 (8.4)

^aTotal number in the variable excluding missing data.

^bTransportation to work, college, school, or family transportation.

Table 2. Exposure to SHS at Home, at Work, and in Transportation

Variable	Saudi Arabia n = 328 (%)
Any smokers at home	327 ^a
Yes	122 (37.3)
No	205 (62.5)
Number of smokers at home	326 ^a
1-2	114 (35)
3-5	5 (1.5)
None	207 (63.5)
Number of cigarettes smoked per day	125 ^a
Less than 5	25 (7.7)
5-20	34 (10.5)
More than 20	8 (2.5)
I do not know	58 (17.8)
Number of hours of SHS exposure at home	325 ^a
Less than 1 hour	38 (11.7)
1-2 hours	18 (5.5)
More than 2 hours	23 (7.1)
None	246 (75.7)
Any smokers at work	220 ^a
Yes	12 (4)
No	208 (64.6)
Number of smokers at work	324 ^a
1-2	8 (2.5)
3-5	7 (2.2)
None	309 (95.4)
Number of hours of SHS exposure at work	323 ^a
Less than 1 hour	12 (3.7)
More than 2 hours	3 (0.9)
None	304 (94.1)
Frequency of SHS exposure at work per day	326 ^a
1-2 times	40 (12.3)
3-5 times	5 (1.5)
More than 5 times	11 (3.4)
None	270 (82.8)
Frequency of SHS exposure at work per week	323 ^a
1-2 days	10 (3.1)
3-5 days	6 (1.9)
More than 5 days	1 (0.3)
None	306 (94.7)
SHS exposure on the transportation vehicle	326 ^a
Yes	31 (9.5)
No	295 (90.5)

Abbreviation: SHS, Secondhand smoke.

^aTotal number in the variable excluding missing data.

in adults and serious health conditions, such as middle ear infections, pneumonia and bronchitis in infants and children (4). SHS exposure can lead to abnormal newborn length and birth weight (3). The negative effect of the nicotine and cotinine on the development of the placenta and on the function of the oxygen transformation to the fetus is believed to be the cause of such newborn deformities (5-7).

Out of the 37% (122) who reported presence of active smokers at home, 35.5% (114) have 1-2 smokers and the rest 3-5 smokers. Our study shows that even though there is high number of consumed cigarettes per day, most of the exposed pregnant reported exposure time of less than

Table 3. Association Between Demographic Characteristics and SHS Exposure

Characteristics	Exposure to SHS		P value*
	Yes n (%)	No n (%)	
Maternal Age (Years)	29.76 ± 7.55	29.24 ± 5.45	0.697
Parity			
Primigravida	20 (58.8)	14 (41.2)	0.420
Multigravida	56 (66.7)	28 (33.3)	
Educational Level	119 ^b		
High	44 (57.9)	32 (42.1)	0.039
Low	33 (76.7)	10 (23.3)	
Employment			
Yes	24 (52.2)	22 (47.8)	0.020
No	54 (73.0)	20 (27.0)	
Monthly Income	111 ^a		
Low ^c	19 (67.9)	9 (32.1)	
Middle-high ^d	51 (61.4)	32 (38.6)	0.543

Abbreviation: SHS, Secondhand smoke.

*P value <0.05 significant.

^bTotal number in the variable excluding missing data.

^cMiddle to high income (more than 5000 Saudi Riyal).

^dLow income (less than 5000 Saudi Riyal).

Statistical tests included chi-square tests and independent t tests.

one hour, which is inconsistent with other studies that shows the more cigarettes consumed at home, the more hours of exposure reported (8,9). This can be due to either the participants and the active smokers have good knowledge about the harms of SHS exposure and they follow measures to avoid it or it could be due to under reporting by the participants of the real timing of exposure.

Significant association between the low level of education of the pregnant women in the study and the more exposure to SHS is found in this study, which has been showed in the literature (3). Being an unemployed is significantly associated with more exposure to SHS among the participants and this could be due to the less chance to be an employed with low level of education which takes us back to the point of the low level of education is associated with more exposure or it could be due to that the unemployed participants stay at home for longer time which increases the risk of more exposure.

Our study did not find significant association between age, parity and level of monthly income and more exposure to SHS, which is inconsistent with the literature that shows pregnant women with younger age and low parity number are more likely to be exposed to SHS (3).

The study did not measure cotinine level in the pregnant women blood, to see if they were telling the truth about their exposure to SHS. Another limitation was we did not specify whether the participants lived in rural or urban areas, as differences in the cultures and the socioeconomic characteristics have been confirmed in the literature (8-10). Also we did not assess the knowledge, attitude and behavior of the participants toward SHS exposure.

Conclusion

One fourth of pregnant women participating in the study were exposed to SHS which constitutes a major public

health problem. Low educational levels and unemployment are associated with more exposure to SHS. There is urgent need to employ evidence based measures to reduce active smoking and exposure to SHS.

Ethical Issues

The research was approved by the Institutional Review Board (IRB) of KKHU and approval was granted before starting the study. All participants were asked to sign an informed written consent before the start of data collection.

Conflict of Interests

Authors declare that there is no conflict of interests.

Financial Support

Researchers received no financial support.

Acknowledgments

The authors acknowledge the nurses team in OB/GYN clinic, King Khaled University Hospital, Riyadh, for their aid to interview the pregnant women. In addition, the authors acknowledge Dr. Shaik Shaffi Ahamed, Associate Professor Department of Family & Community Medicine, College of Medicine, King Saud University, for his aid to do the statistical analysis using SPSS.

References

- Worldwide burden of disease from exposure to second-hand smoke. http://www.who.int/quantifying_ehimpacts/publications/shsarticle2010/en/
- Bassiony MM. Smoking in Saudi Arabia. *Saudi Med J*. 2009;30(7):876-881.
- Wahabi HA, Alzeidan RA, Fayed AA, Mandil A, Al-Shaikh G, Esmail SA. Effects of secondhand smoke on the birth weight of term infants and the demographic profile of Saudi exposed women. *BMC Public Health*. 2013;13:341. doi:10.1186/1471-2458-13-341.
- Bloch M, Althabe F, Onyamboko M, et al. Tobacco use and secondhand smoke exposure during pregnancy: an investigative survey of women in 9 developing nations. *Am J Public Health* 2008;98(10):1833-1840. doi: 10.2105/AJPH.2007.117887.
- Vardavas CI, Fthenou E, Patelarou E, et al. Exposure to different sources of second-hand smoke during pregnancy and its effect on urinary cotinine and tobacco-specific nitrosamine (NNAL) concentrations. *Tobacco Control*. 2013;22(3):194-200. doi:10.1136/tobaccocontrol-2011-050144.
- Romani FE, Lanzone AN, Tropea AN, Tiberi FE, Catino S, Apa RO. Nicotine and cotinine affect the release of vasoactive factors by trophoblast cells and human umbilical vein endothelial cells. *Placenta*. 2011;32(2):153-160.
- Zdravkovic T, Genbacev O, McMaster MT, Fisher SJ. The adverse effects of maternal smoking on the human placenta: a review. *Placenta*. 2005;26(suppl A):81-6.
- Yang L, Tong EK, Mao Z, Hu T-w. Exposure to secondhand smoke and associated factors among non-smoking pregnant women with smoking husbands in Sichuan province, China. *Acta Obstetrica et Gynecologica Scandinavica*. 2010;89(4):549-557. doi:10.3109/00016341003713851.
- Zhang L, Hsia J, Tu X, et al, Stanton B. Peer Reviewed: Exposure to secondhand tobacco smoke and interventions among pregnant women in China: A systematic review. *Preventing Chronic Disease*. 2015;12. doi:10.5888/pcd12.140377
- Bhanji S, Andrades M, Taj F, Khuwaja AK. Factors related to knowledge and perception of women about smoking: a cross sectional study from a developing country. *BMC Womens Health* 2011;11:16. doi:10.1186/1472-6874-11-16.

Copyright © 2016 The Author(s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Appendix 1

Demographic characteristics: This section asks about demographic profile.

1-How old are you?

2-What is your nationality?

- Saudi
 Not Saudi

3-What is your marital status?

- Married
 Divorced
 Widow

4-number of previous pregnancies?

- zero
 1-5 times
 more than 5 times

5-What is your highest degree of education?

- Primary school
 Intermediate school
 High school
 Undergraduate
 Bachelor
 Postgraduate
 Illiterate

6-Do you work?

- Yes
 No

7-What do you use for transportation?

- Private car
 Taxi
 Bus

8- What is the range of your household income?

- Less than 4,999SR
 5,000-9,999 SR
 10,000-14,999 SR
 More than 15,000 SR

9-do you smoke? Which type?

- Cigarettes
 Hookah
 Chewing tobacco
 I don't smoke

Exposure to SHS: This section asks about Participant's exposure to smoking during daily/weekly/monthly routine.

10-Is there any smoker at your home?

- Yes
 No

11-How many smokers are there?

- 1-2
 3-5
 None

12-How many cigarettes do they smoke per day at home?

- Less than 5
 5-20
 More than 20
 Don't know

13- How many hours approximately do you expose to cigarettes smoke per day at home?

- Less than one hour
 1-2 hours
 More than two hours
 None

14- Do you expose to cigarettes smoke inside the home or not?

- Yes
 No

15-Is there any smoker in your work place?

- Yes
 No

16- How many smokers are there?

- 1-2
 3-5
 None

17-How many hours approximately do you expose to cigarettes smoke per day at work?

- Less than one hour
 More than two hours
 None

18- How many times do you expose to cigarettes smoke per day?

- 1-2 times
 3-5 times
 more than 5 times
 None

19-How many days do you expose to cigarettes smoke per week at work?

- 1-2 day/s
 3-5 days
 More than 5 days
 None

20- Do you expose to cigarettes smoke in transportation vehicle?

- Yes
 No