

A checklist of Iranian Eupodoidea (Acari: Prostigmata)

Maryam Darbemamieh¹, Hamidreza Hajiqanbar^{1*} and Mohammad Khanjani²

1. Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, 14115-336, Tehran, Iran.

2. Department of Plant Protection, College of Agriculture, Bu Ali-Sina University, Hamedan, Iran.

Abstract: The present checklist is a compilation of the eupodoid mites of Iran using published records and original data from recent researches. It contains 19 species belonging to 13 genera and five families. Family Cocceupodidae Jesionowska, 2010 (because of moving the genera to a new family) and two species i.e. *Foveacheles (Foveacheles) cegetensis* Zacharda, 1983 and *Linopodes antennaepes* Banks, 1894 are new records for Iranian mite fauna. In addition to some corrections to specific identities which have been previously reported in Iranian literature, we report here the known geographical distribution and habitats in Iran and distribution in the world as well.

Keywords: Eupodoid mites, checklist, new record, *Foveacheles (Foveacheles) cegetensis*, *Linopodes antennaepes*, Iran

Introduction

According to Zhang *et al.*, 2011, nine families have been distinguished in the superfamily Eupodoidea Koch, 1842. These families have been listed as Eupodidae Koch, 1842 (10 genera, 69 species), Penthaleidae Oudemans, 1931 (five genera, 16 species), Penthalodidae Thor, 1933 (six genera, 35 species), Rhagidiidae Oudemans, 1922 (28 genera, 157 species), Strandmanniidae Zacharda, 1979 (one genus, two species), Eriorhynchidae Qin & Halliday, 1997 (one genus, five species), Pentapalpidae Olivier & Theron, 2000 (one genus, one species), Dendrochaetidae Olivier, 2008 (one genus, one species) and Cocceupodidae Jesionowska, 2010 (three genera, 23 species). Eupodoids are worldwide in distribution and often are found in extreme environments including high alpine and polar habitats. They are predaceous, fungivorous, bryophagous or phytophagous and not known to

be symbiotic on other animals (Krantz and Walter, 2009; Qin, 1996).

The present checklist is a survey to collect the results of all identified eupodoid mites in Iran and to indicate their taxonomic status, habitats and distribution. Some name changes, new records and/or new location reports are added to previous reports. Distribution in the world is added as much as possible.

Materials and Methods

The authors checked all available papers including new taxa collected from Iran till 2012, and new reports of the families, genera and species. Collected samples from soil, mushroom, stored products, litter and moss were extracted using Berlese-Tullgrens over a period of 48 hours. Some species were removed directly using a stereomicroscope. Specimens were cleared in lactic acid and mounted on permanent slides using Hoyer's medium (Krantz and Walter, 2009). Identification of rhagidiid species was verified by Dr Miloslaw Zacharda. The used taxonomy method, followed that of Qin (1996), Zacharda (1980) and Jesionowska (2010 b). All mounted specimens of new records are deposited

Handling Editor: Dr. Ali Asghar Talebi

* **Corresponding author**, e-mail: hajiqanbar@modares.ac.ir
Received: 18 May 2013, Accepted: 25 August 2013

in the Acarology Museum, Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran.

Superfamily Eupodoidea Koch, 1842

Family Cocceupodidae Jesionowska, 2010

Cocceupodes Thor, 1934

Cocceupodes sp.

Distribution and habitat in Iran: Mazandaran, Citrus (Faraji and Kamali, 1993); Guilan, soil of tea fields (Nejadghanbar *et al.*, 2010).

World Distribution: Palaearctic [Austria, Britain, Belgium (Ducarme *et al.*, 2004), Denmark, Germany (Strandtmann and Goff, 1978), Ireland, Italy, Norwegian mainland, Poland (Jesionowska, 2010), Norway, Sweden, Switzerland], Nearctic (United States) (Strandtmann and Goff, 1978), Australian (New Zealand) (Qin, 1998), Afrotropical (Southern Africa) (Olivier and Theron, 2003), (Egypt) (Abou-Awad *et al.*, 2006).

Linopodes Koch, 1835

Linopodes sp.

Distribution and habitat in Iran: East Azerbaijan, Soil (Fathipour, 1994); Kermanshah, Soil (New location).

World Distribution: Palaearctic (Austria, Britain, Denmark, Hungary, Poland, Switzerland, Japan) (Banks, 1915; Shiba, 1976; Clancy, 1981); Nearctic (United States) (Banks, 1915), Australian (Australia, New Zealand) (Qin, 1998), Oriental region (Malaysia) (Shiba, 1976); Ethiopian region (Egypt, South Africa) (Abou-Awad *et al.*, 2006).

Linopodes antennaepes Banks, 1894

Distribution and habitat in Iran: Kermanshah and Kamyaran, Mushroom (New record for Iran).

World Distribution: Palaearctic (Italy, Britain), Nearctic (United States), Australian (Australia) (Banks, 1915; Clancy, 1981).

Linopodes cameronensis Shiba, 1976

Distribution and habitat in Iran: Gachsaran, Soil (Moradian and Ostovan, 2011)

World Distribution: Oriental region (Malaysia) (Shiba, 1976)

Family Eupodidae Koch, 1842

Eupodes Koch, 1835

Eupodes sp.

Distribution and habitat in Iran: East Azerbaijan (Miandoab), sugar beet soil (Haddad *et al.*, 2005).

World Distribution: Afro-tropical (North Africa) (Abou-Awad *et al.*, 2006), Nearctic (Mexico, USA) (Sepedap and Withford, 1990), Palaearctic (Austria, Britain, France, Germany, Iceland, Italy, Norway, Switzerland) (Strandtmann and Goff, 1978; Olivier, 1997).

Eupodes crozetensis Strandtmann & Davies, 1972.

Distribution and habitat in Iran: East Azerbaijan, Soil (Fathipour, 1994); Fars, Stored hay (Ostovan, 1993).

World distribution: Southern Indian Ocean (Possession Island, Crozet Islands) (Strandtmann and Davies, 1972).

Eupodes sigmoidensis Strandtmann & Goff, 1978.

Distribution and habitat in Iran: Hamedan, Bean (Khanjani, 1996); East Azerbaijan, Alfalfa Soil (Lotfollahi *et al.*, 2010)

World Distribution: West Africa (Ivory Coast, Lamto-Pakobosavanna), Nearctic (USA) (Strandtmann and Goff, 1978).

Eupodes voxencollinus Thor, 1934

Distribution and habitat in Iran: Hamedan, Bean (Khanjani, 1996).

World Distribution: Nearctic and Palaearctic (North Africa, Poland, Norway, Germany, Britain) (Fauna Europaea, 2012; Jesionowska, 2003; Strandtmann and Goff, 1978).

Benoinyssus sp.

Distribution and habitat in Iran: Miandoab, sugar beet soil (Haddad *et al.*, 2005).

World Distribution: Palaearctic (Britain, Germany, Poland) (Fauna Europaea, 2012), Nearctic (Mexico), Ethiopian (Southern Africa) (Sepedap and Withford, 1990; Olivier and Theron, 2003).

Claveupodes Strandtmann & Prasse, 1977

Claveupodes sp.

Distribution and habitat in Iran: East Azerbaijan, Alfalfa Soil (Lotfollahi *et al.*, 2010); East Azerbaijan (Miandoab), sugar beet soil (Haddad *et al.*, 2005).

World Distribution: Palaearctic (Belgium, Britain, Germany, Poland), Australian region

(Fauna Europaea, 2012; Russell *et al.*, 2010; Ducarme *et al.*, 2004).

Family Penthaleidae Oudemans, 1931

Penthaleus Koch, 1835

Penthaleus major (Duges, 1834)

Distribution and Habitat in Iran: Chahar Mahal & Bakhtiari, Cereals soil (Noorbakhsh, 1993), (Noorbakhsh and Kamali, 1995); East Azerbaijan, soil (Fathipour, 1994); Hamedan, pagoda tree (*Sophora alopecuroides* L.) soil, (Khanjani, 1996), (Khanjani and Kamali, 1993), Khorasan, saffron soil (Rahimi, 1991); *Acroptilon* sp., soil (Modarres Awal, 1997), Khuzestan, soil (Sadeghi Nameghi, 1990), (Sadeghi Nameghi and Kamali, 1993).

World Distribution: Worldwide (Umina *et al.*, 2004).

Penthaleus minor (Canestrini, 1886)

Distribution and habitat in Iran: East Azerbaijan, soil (Fathipour, 1994).

World Distribution: Australian (Australia) (Qin, 1998), Palaearctic (Britain, Germany, Italy, Switzerland), Nearctic (Qin, 1998, Fauna Europaea, 2012).

Family Penthelodidae Thor, 1933

Penthelodes Murray, 1877

Penthelodes polonicus Jesionowska, 2010 (correct name used in Jesionowska, 2010a).

Penthodes polonica, Jesionowska, 2010 (incorrect name used in Hajizadeh and Noei, 2012).

Distribution and habitat in Iran: Guilan (Fuman, Rud-khan castle, Shaft, chubar), unknown plant Soil (Hajizadeh and Noei, 2012).

World Distribution: Poland (Jesionowska, 2010a).

Family Rhagidiidae Thor & Willmann, 1941

Brevipalpia minima Zacharda, 1980

Distribution and habitat in Iran: Hamedan (Emamzade), sour cherry orchard (Zacharda *et al.*, 2012).

World Distribution: Czech Republic (Central Bohemia), Iran (Zacharda *et al.*, 2012).

Coccorhagidia Thor, 1934

Coccorhagidia clavifrons (Canestrini, 1886)

Distribution and habitat in Iran: Hamedan, Alfalfa (Khanjani, 1996); Hamedan-Shokriyeh, peach, sour cherry and apple orchard soil (Zacharda *et al.*, 2012); Tehran, soil (new location).

World Distribution: Holarctic (Italy, Czech Republic, Slovak Republic, Russia, Canada, Hawaii, USA) (Zacharda *et al.*, 2012; Fauna Europaea, 2012).

Coccorhagidia pittardi Strandtmann, 1971

Distribution and habitat in Iran: Ardabil, hazelnut orchard soil; Hamedan-Shokriyeh, Peach orchard soil (Zacharda *et al.*, 2012).

World Distribution: Holarctic (Russia, Austria, Slovak Republic, United States) (Zacharda *et al.*, 2012).

Foveacheles Zacharda, 1980

Foveacheles sp.

Distribution and habitat in Iran: Hamedan (Shokriyeh), peach orchard soil (Zacharda *et al.*, 2012); Esfahan, soil (new location).

World Distribution: Palaearctic (Austria, Belgium, Britain, Czech Republic, France, Germany, Hungary, Ireland, Norway, Poland, Slovakia, Spain, Sweden, Switzerland, Netherland), Nearctic (Zacharda *et al.*, 2012; Fauna Europaea, 2012).

Foveacheles (Foveacheles) brevichelae Zacharda, 1980

Distribution and habitat in Iran: Hamedan (Shokriyeh), peach orchard soil, (Zacharda *et al.*, 2012).

World Distribution: Holarctic region (Czech Republic; the Otztal Alps in Austria; Elbrus, Central Caucasus, Russia; Tuktoyaktuk, Northwest Territory, Canada) (Zacharda *et al.*, 2012).

Foveacheles (Foveacheles) cegetensis Zacharda, 1983.

Distribution and habitat in Iran: Tehran, unknown plant soil (New record for Iran).

World Distribution: Czech Republic (Zacharda, 1983; Fauna Europaea, 2012).

Foveacheles (Foveacheles) osloensis (Thor, 1934)

Syn.: *Rhagidia osloensis* Thor, 1934

Distribution and habitat in Iran: East Azerbaijan, soil (Fathipour, 1994).

World Distribution: Palaearctic (Austria, Czech Republic, Germany, Norway, Sweden) (Zacharda, 1980; Fauna Europaea, 2012).

Rhagidia Thorell, 1871

Rhagidia sp.

Distribution and habitat in Iran: East Azarbaijan (Miandoab), sugar beet soil (Haddad *et al.*, 2005); Hamedan (Shokriyeh), peach orchard soil (Zacharda *et al.*, 2012).

World Distribution: Palaearctic (Austria, Belgium, Britain, Czech Republic, Denmark, France, Germany, Italy, Poland, Romania, Slovakia, Norway, Sweden, Switzerland, The Netherlands), Nearctic, Oriental (Fauna Europaea, 2012; Zacharda *et al.*, 2012).

Rhagidia cf. breviseta Zacharda, 1995

Distribution and habitat in Iran: Hamedan (Absineh), apple orchard soil (Zacharda *et al.*, 2012).

World Distribution: Holarctic (Canada, Alaska, Slovakia, Finland, Russia) and prefers cooler habitats such as caves, peat bogs and moss pads (Zacharda *et al.*, 2012).

Robustocheles Zacharda, 1980

Robustocheles mucronata (Willmann, 1936)

Distribution and habitat in Iran: Fars, Bed of humid stores (Ostovan, 1993); Ardabil, hazelnut orchard soil (Zacharda *et al.*, 2012); Hamedan-Ganjnameh, plum orchard soil (Zacharda *et al.*, 2012); Hamedan (Heydareh), apple orchard soil (Zacharda *et al.*, 2012); Golestan, soil (New location).

World Distribution: worldwide distribution [Russia (Central Siberia, Elbrus mountain), Poland, Slovak Republic, Czech Republic (North Bohemia), Thailand, North Vietnam, Brazil, Alaska] (Zacharda *et al.*, 2012).

Robustocheles hamedanensis Zacharda, 2012

Distribution and habitat in Iran: Hamedan (Shokriyeh) litter in peach orchard (Zacharda *et al.*, 2012).

World Distribution: Iran (Zacharda *et al.*, 2012)

Shibaia Zacharda, 1980

Shibaia heteropoda (Berlese, 1910)

Distribution and habitat in Iran: East Azarbaijan (Miandoab), sugar beet soil (Haddad *et al.*, 2005).

World Distribution: Palaearctic and Nearctic (Zacharda, 1980).

Shibaia longisensilla (Shiba, 1969), correct name in Zacharda, 1980

Shibaia longicornis (Shiba, 1969), incorrect name in Iranian records.

Distribution and habitat in Iran: Hamedan, Alfalfa (Khanjani and Kamali, 1993), Sainfoin (*Onobrychis viciifolia*) (Modarres Awal, 1997).

World Distribution: Holarctic region (abundant in Czech Republic) (Zacharda, 1980).

Discussion

The mites of the genus *Cocceupodes* show substantial homogeneous structure, distinctly different from species of typical genus *Eupodes*, and other species recently regarded as belonging to the family Eupodidae (e.g. *Benoinyssus* or even *Protereunetes*). This is the reason why Jesionowska (2010b) classified them as a separate family, Cocceupodidae, including three genera: *Cocceupodes*, *Filieupodes* and *Linopodes*. Main differential features are rostral setae, *ro*, located just behind a naso and not on it (differently from those in other eupodoids), and just two pairs of circumanal setae, with total number of ten pairs of setae present on opisthosoma (excluding genital region setae). The family Cocceupodidae was derived from the family Eupodidae in 2010 and therefore, the reports from the genus *Linopodes* were classified as Cocceupodidae instead of the family Eupodidae. In this study, *Linopodes antennaepes* Banks, 1894 is reported as new record of this family from Iran. Banks (1894) first described it as a reddish or yellowish mite, sometimes with some pale marks. Legs are mostly yellowish, except distal half of leg I, which is hyaline. Body oblong, rounded in front and behind; broadest at hind margin of cephalothorax; cephalothorax semicircular, with a large shining eye on each side. A narrow, emarginated, smooth band is just behind cephalothorax, which gives it a median projection reaching to the tip of the abdomen, the whole forming a T. Dorsum has a few scattered hairs. Leg I more than twice as long as body, femur I longer than body; legs II and III slender, not quite as long as body; leg IV with femur enlarged. Chelicerae are short, forming a little cone; palpi a little longer than chelicera, joints subequal, the third being the longest. It differs from the European *L. motorius* in having tibia I nearly as long as the metatarsus, and the tarsus I being divided into three or four joints. The

body of the male is a little more globose than that of the female. This species lives on the ground, and is most common under pieces of wood, bark, etc. that have been on the ground for some time. The first pair of legs is used as feelers; ordinarily, it walks slowly but when disturbed can move very rapidly to the rear (Banks, 1894). In current study, *L. antennaeipes* was collected from cultivated and forest mushrooms in Kermanshah and Kamyaran. *Linopodes* species are present in mushroom cultivation but are not common (Hussey et al., 1969; Clancy, 1981). *L. antennaeipes* is the only species in the genus recorded from Australia by Halliday (2001).

Other new record in this paper is *Foveacheles* (*Foveacheles*) *cegetensis* Zacharda, 1983 from the family Rhagididae which is recorded for first time from Iran. Some diagnostic features for identification of this species are as below. Proximal cheliceral seta inserted just before joint of digitus mobilis and almost reaching basis of distal cheliceral seta. Rhagidial organ I and II consist of 4 separated rhagidial setae. Stellate seta is between 1st and 2nd proximal rhagidial setae. Length of the body 970-1150 µm (Zacharda, 1983). The epimeral formula 3-1-6-3 and the location of the solenidion on genu III is medioventral, proximad of the first pair of ventral setae (Zacharda, 1993).

Acknowledgement

The authors wish to thank Prof. M. Zacharda (Department of Biodiversity and Biomonitoring, Institute of Systems Biology and Ecology, Academy of Sciences of the Czech Republic) for help in identification of rhagidiid mites. Special thanks go to Dr. A. Zamany and S. Barzegar (Department of Agriculture, Razi University of Kermanshah, Iran) for providing some specimens.

References

- Abou-Awad, B. A., El-Sawaf, B. M. and Abdelkhalek, A. A. 2006. Four new species of eupodid mites from Egypt (Acari: Eupodoidea: Eupodidae). *Acarologia*, 46 (1-2): 43-52.
- Banks, N. 1894. Some new American Acarina. *Transactions of the American Entomological Society*, 21 (2): 209-222.
- Banks, N. 1915. The Acarina or mites. A review of the group for the use of economic entomologists. United States Department of Agriculture. Office of the Secretary, report 108: 1-153.
- Cepedap, J. and Withford, W. G. 1990. Microarthropodos edaphicos del desierto Chihuahuense, al norte de Mexico. *Folia Entomologica Mexicana*, 78: 257-272.
- Clancy, G. 1981. Observations of mites associated with the low yielding crops of cultivated *Agaricus bisporus* in Australia. *Mushroom Science*, 11: 233-244.
- Coineau, Y. 1976. La premiere prelarveconnue du genre *Eupodes*, *Eupodesstrandtmanni*, n. sp. *Acarologia*, 18: 56-64.
- Ducarme, X., Andre, H. M. Wauthy, G. and Lebrun, P. 2004. Are there real endogenic species in temperate forest mites? *Pedobiologia*, 48: 139-147.
- Faraji, F. and Kamali, K. 1993. *Mites associated with Citrus spp.* In eastern Mazandaran. Proceedings of 11th Iranian Plant Protection Congress, Gilan University, Rasht, p. 186.
- Fathipour, Y. 1994. Soil mites fauna in orchards of Tabriz and population fluctuation and abundance of important species. M. Sc. thesis, Tarbiat Modarres University, Tehran, 172 pp.
- Fauna Europaea, 2012. Fauna Europaea version 2.5. Available on: <http://www.faunaeur.org> (accessed August 20, 2013)
- Haddad Irani-Nejad, K., Hajiganbar, H. R. and Talebi Chaichi, P. 2005. An introduction of the prostigmatic mites in sugarbeet fields in Miandoab plain. *Iranian Journal of Agricultural Science*, 36 (1): 247-262.
- Hajizadeh, J. and Noei, J. 2012. *Report of a new family for the mite fauna of Iran: Penthalodidae (Acari, Prostigmata)*. 20th Iranian Plant Protection Congress, p. 486.

- Halliday, R. B. 2001. Acarina. Mites, in Australian Faunal Directory. Australian Biological Resources Study, Canberra. Available on: <http://www.deh.gov.au/biodiversity/abrs/online-resources/abif/fauna/index.html>. (accessed September 2004).
- Hussey, N. W. 1961. Identification of Phoridae (Diptera) attacking cultivated mushrooms in Britain. *Annals and Magazine of Natural History*, 13 (3): 599-603.
- Jesionowska, K. 2003. Observations on the morphology of some eupodoid and endeostigmatic gnathosomata (Actinotrichida, Actinedida, Eupodoidea and Endeostigmata). *Acta zoologica cracoviensia*, 46 (3): 257-268.
- Jesionowska, K. 2010 a. A morphological study of the genus *Penthalodes* (Acari, Prostigmata, Eupodoidea, Penthalodidae) with description of a new species. *Zootaxa*, 2672: 29-49.
- Jesionowska, K. 2010 b. Cocceupodidae, a new family of eupodoid mites, with description of a new genus and two new species from Poland. Part I. (Acari: Prostigmata: Eupodoidea). *Genus*, 21 (4): 637-658.
- Kamali, K., Ostovan, H. and Atamehr, A. 2001. A Catalogue of Mites and Ticks (Acari) of Iran. Islamic Azad University Scientific Publication Center. 192 pp.
- Khanjani, M. 1996. Mites (Acari) associated with Fabaceae plants in Hamedan province and functional responses of *Anystis baccharum* (L.) and *Erythraeus* sp. to developmental stages of *Tetranychus turkestanii* (U. & N.) Ph. D. dissertation, Tarbiat Modarres University, Tehran, Iran, 437 pp.
- Khanjani, M. and Kamali, K. 1993. *Mites (Acari: Actinedida) associated with Fabaceae plants in Hamedan*. Proceedings of 11th Iranian Plant Protection Congress, Guilan University, Rasht, p. 265.
- Krantz, G. W. and Walter, D. E. 2009. A manual of acarology. 3rd ed. Lubbock (TX), Texas Tech University Press: 816 p.
- Lotfollahi, P., Haddad Irani-Nejad, K., Bagheri, M. and Valizade, M. 2010. Prostigmatid soil mites of Alfalfa fields in northwest of Iran (East Azerbaijan province) with one genus, subgenus and four species as new records. *Munis Entomology & Zoology*, 5: 1001-1010.
- Modarres Awal, M. 1997. List of agricultural pests and their natural enemies in Iran (revised edition). Ferdowsi University Press, 429 pp.
- Moradian, H. and Ostovan, H. 2011. *First report of a genus and species of the family Eupodidae (Acari: Prostigmata) from Iran*. Proceedings of Global Conference on Entomology (GCE), P: 141.
- Nejadghanbar, N., Arbabi, M. and Vafaeishoushtari, R. 2010. Study on geographical distribution and abundance of plant feeding mites on green parts and soil surface of tea plants gardens in eastern parts of Gilan Province of Iran. *Journal of Entomological Research*, 2 (4): 329-338.
- Noorbakhsh, S.H. 1993. Faunistic study of cereal mites in eastern ChaharMahal & Bakhtiari and biology of brown wheat mite *Petrobia latens* (Muller). M. Sc. thesis shahid Chamran University, Ahwaz, Iran, 120 pp.
- Noorbakhsh, S. H. and Kamali, K. 1995. Phytophagous and predaceous mites (Acari) associated with gramineous plants in ChaharMahal & Bakhtiari. Proceedings of 12th Iranian Plant Protection Congress, Tehran University, Karaj, P: 310.
- Olivier, P. A. S. 1998. The genus *Eupodes* Koch (Acari: Prostigmata: Eupodidae) from southern Africa with a redescription of *E. parafusifer* Meyer & Ryke and descriptions of two new species. *African Entomology*, 6 (2): 281-284.
- Olivier, P. A. S. 2008. Dendrochaetidae, a new family of mites (Acari: Prostigmata), with descriptions of a new genus and species from South Africa. *African Zoology*, 43 (1): 16-24.
- Olivier P. A. S. and Theron, P. D. 2000. Pentapalpidae, a new family of eupodoid mites Prostigmata: Eupodoidea) from South Africa. *Acarologia*, 40(1999): 385- 392.
- Olivier, P. A. S. and Theron, P. D. 2003. Descriptions of two new species of the genus *Cocceupodes* Thor, 1934 (Eupodidae), and a

- checklist of southern African Eupodoidea. African Entomology 11 (2):163-171.
- Ostovan, H. 1993. Faunistic study of stored product mites in Kazerun and biology of important species. M.S.c thesis, Tarbiat Modares University, Tehran, 172 pp.
- Qin, T. K. 1996. A review and cladistic analysis of the Eupodoidea (Acari: Acariformes). Systematic & Applied Acarology, 1: 77-105.
- Qin, T. K. 1998. A checklist and key to species of Eupodoidea (Acari: Prostigmata) from Australia and New Zealand and their subantarctic islands. Journal of the Royal Society of New Zealand, 28 (2): 295-307.
- Qin, T. K. and Halliday, B. 1997. Eriorhynchidae, a new family of Prostigmata (Acarina), with a cladistic analysis of eupodoid species of Australia and New Zealand. Systematic Entomology, 22: 151-171.
- Rahimi, H. 1991. Faunistic study of arthropods (Arthropoda) associated with saffron in Gonabad and Ghayen, and biology of important species. M.Sc. thesis, Faculty of Agriculture, Shahid Chamran University, Ahwaz, 141 pp.
- Russell, D. J., Hohberg, K. and Elmer, M. 2010. Primary colonisation of newly formed soils by actinedid mites. Soil Organisms, 82 (2): 237-251.
- Sadeghi Nameghi, H. 1990. Mites (Acari) associated with sugar cane and cereals with emphasis on biology of injurious species in Khuzestan, Iran. M. Sc. thesis, Shahid Chamran University, Ahwaz, 167 pp.
- Sadeghi Nameghi, and Kamali, K. 1993. Mites (Acari) associated with sugar cane and cereals in Khuzestan, Iran. The Scientific Journal of Agriculture, Shahid Chamran University, Ahwaz, 16 (1, 2): 3-13.
- Shiba, M. 1976. Taxonomic investigation on free-living Prostigmata from the Malay Peninsula. Nature and Life in Southeast Asia, 7: 83-229.
- Strandtmann, R. W. and Davies, L. 1972. Eupodiform mites from possession island, crozet islands, with a key to the species of *Eupodes* (Acarina: Prostigmata). Pacific Insects, 14: 39-56.
- Strandtmann, R. W. and Goff, M. L. 1978. The Eupodoidea of Hawaii (Acarina: Prostigmata). Pacific Insects, 3-4: 121-143.
- Umina, P. A., Hoffmann, A. A. and Weeks, A. R. 2004. Biology, ecology and control of the *Penthaleus* species complex (Acari: Penthaleidae). Experimental and Applied Acarology, 34 (3-4): 211-237.
- Zacharda, M., 1979. Strandtmanniidae – a new family of Eupodoidea (Acarina: Prostigmata). Vestník Československe Spolecnosti Zoologicke, 43 (1): 76-80.
- Zacharda, M. 1980. Soil mites of the family Rhagidiidae (Actinedida: Eupodoidea). Morphology, systematics, ecology. Acta Universitatis Carolina-Biologica, 1978 (5/6): 489-785.
- Zacharda, M. 1983. The Rhagidiidae (Acarina: Prostigmata) from the Central Caucasus, Siberia and some other parts of the USSR. Věstník Československé společnosti zoologické, 47 (4): 304-319.
- Zacharda, M. 1993. Glacial relict Rhagidiidae (Acari: Prostigmata) from superficial underground enclosures in the Krkonoše Mountains, Czechoslovakia, Journal of Natural History, 27: 1, 47-61.
- Zacharda M. 1997. New taxa of Rhagidiidae (Acari: Prostigmata) from North America. Part V. Genus *Robustocheles* Zacharda, with a key to world species of the genus. Journal of Natural History, 31 (7): 1075-1103.
- Zacharda, M., Ueckermann, E. A., Rostami, E. Abbasipour, H. and Khanjani, M. 2012. A survey of Rhagidiidae (Acari: Prostigmata) in fruit orchards in Iran, with description of a new species of *Robustocheles*. International Journal of Acarology, 38 (1): 30-34.
- Zhang, Z. Q., Fan, Q. H., Pesic, V., Smit, H., Bochkov, A. V., Khaustov, A. A., Baker, A. Wohltmann, A., Wen, T., Amrine, J. W., Beron, P., Lin, J., Gabrys, G. and Husband, R. 2011. "Order Trombidiformes Reuter, 1909. In: Zhang, Z. Q. (Ed.) Animal Biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148, Magnolia Press, Auckland, New Zealand, 237 p.

فهرست کنه‌های بالاخانواده (Acari: Prostigmata) Eupodoidea در ایران

مریم درب امامیه^۱، حمیدرضا حاجی قنبر^{۱*} و محمد خانجانی^۲

۱- گروه حشره‌شناسی، دانشکده کشاورزی، دانشگاه تربیت مدرس، صندوق پستی ۳۳۶-۱۴۱۱۵، تهران، ایران.

۲- گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه بوعلی سینا همدان، همدان، ایران.

* پست الکترونیکی نویسنده مسئول مکاتبه: hajiqanbar@modares.ac.ir

دریافت: ۲۸ اردیبهشت ۱۳۹۲؛ پذیرش: ۳ شهریور ۱۳۹۲

چکیده: فهرست حاضر دربرگیرنده همه گزارش‌های انجام شده از بالاخانواده Eupodoidea در ایران شامل ۱۹ گونه از ۱۳ جنس و ۵ خانواده است که از بین آنها خانواده Cocceupodidae Jesionowska, 2010 (به دلیل جابه‌جایی جنس‌ها به خانواده جدید) و دو گونه *Foveacheles (Foveacheles)* *Linopodes antennaepes* Banks, 1894 و *cegetensis* Zacharda, 1983 گزارش‌های جدید برای فون کنه‌های ایران هستند. برخی تصحیحات در اسامی گونه‌های گزارش شده قبلی به همراه پراکنش همه گونه‌ها در ایران، زیستگاه آنها و نیز پراکنش این گونه‌ها در نقاط مختلف جهان به همراه این لیست آورده شده است.

واژگان کلیدی: کنه‌های یوپودوئیدا، چک لیست، رکورد جدید، *Foveacheles (Foveacheles)* *Linopodes antennaepes* *cegetensis* ایران