

Research Article

Molecular characterization of aster yellows phytoplasma associated with citrus varieties, using Multiplex PCR

Cobra Moslemkhani¹, Raheleh Shahbazi¹, Sareh Baghaee Ravari² and Fatemeh Khelghatibana¹

1. Seed and Plant Certification and Registration Institute. P. O. Box: 31535-1516, Karaj, Iran.

2. Department of Crop Protection, Faculty of Agriculture, Ferdowsi University of Mashhad, P. O. Box: 91775-1163, Mashhad, Iran.

Abstract: A survey was carried out on citrus nurseries in Mazandaran and Kerman provinces as the major citrus growing regions of Iran. Different varieties of orange and tangerine with abnormal symptoms were examined for the occurrence of *Candidatus Liberobacter asiaticus*, *Spiroplasma citri* and phytoplasma as the most important vascular limited pathogens in citrus. By visual inspection and molecular examination of 360 samples for the presence of fastidious prokaryotes only an occasional occurrence of phytoplasma and *Spiroplasma citri* were detected, using phytoplasma universal primer pairs P1/P7 and fU5/rU3 (in single tube nested PCR assay) and *Spiroplasma* specific primers P89-r/f and P58-6f/4r. PCR products of phytoplasmas were digested with *MseI*, *AluI*, *BamHI* and *TaqI* restriction enzymes and the digestion results showed phytoplasma related to aster yellows phytoplasma group. No amplicon was obtained for *Candidatus Liberobacter asiaticus*. The findings suggest when pathogen's titre is low; symptoms of vascular limited disease are almost confused with disorder and deficiencies signs. So for sanitary selection and certification program, using sensitive detection method is recommended. Multiplex PCR assay used in this study successfully detected *Candidatus Liberobacter asiaticus*, *Spiroplasma citri* and phytoplasma in one reaction and suggests for health certification programs.

Keywords: vascular limited, citrus, Phytoplasma, Multiplex PCR

Introduction

Iranian Seed and Plant Certification and Registration Institute (SPCRI) founded in order to get an improvement of seed and plant material production. The main goal of SPCRI is the genetic and phytosanitary certification of plant propagative material. Citrus is one of the most important fruit crops in Iran. That is susceptible to a large number of pathogens. The

laboratories of SPCRI are planning to expand certification program for citrus nuclear stock and mother trees to enhance plant health monitoring.

Lime witches broom caused by *Candidatus phytoplasma aurantifolia* (Bove, *et al.*, 2000), Stubborn caused by *Spiroplasma citri* (Bove, 2006) and huanglongbing caused by *Candidatus Liberobacter asiaticus* (Mohkami *et al.*, 2011) are most serious citrus diseases facing us today. These diseases have been experimentally transmitted from infected trees to several citrus cultivars by grafting (Salehi *et al.*, 2007, Raju *et al.*, 1981, Bove, 2006, Frison and Taher, 1991, da Graça, 1991 OEPP/EPPO,

Handling Editor: Dr. Naser Safaie

* **Corresponding author**, e-mail: moslemkhany@yahoo.com
Received: 15 October 2013, Accepted: 14 January 2014
Published online: 3 September 2014

2006). These phloem-limited agents spread through vegetative propagation and phloem feeding vector (Seemuller 2002).

The symptoms of mentioned diseases are not always reliable for diagnosis because they can vary according to environmental conditions and pathogen concentrations.

The early detection of greening, Stubborn and witches broom are vital for revealing their real distribution and for phytosanitary certification. PCR as a sensitive molecular tool has been developed for detection of these diseases (Hung *et al.*, 1999; Heinrich *et al.*, 2001; Minsavage *et al.*, 1994 and Yokomi *et al.*, 2008). In view of the increasing interest in plant pathology for the detection of more than one targets, such as mixed infection pathogens in single reaction (Bertolini *et al.*, 2003), multiplex PCR protocols have been developed. We describe for the first time, a multiplex PCR for the detection of prokaryotic agents frequently infecting citrus trees. Using healthy mother trees and clean bud wood could prevent spreading of diseases Thus; this study was conducted to detect the presence of mentioned diseases in Iranian citrus orchards and nurseries.

Materials and Methods

Plant samples

A total of 360 samples from different varieties of orange and tangerine (such as Navelate, Okitsu, Fukumoto, Valencia late, Washington navel foyus, Hoshimoto, Washington navel, Clementine clime new less1, Clematis manitol, Lime Bears, Delta seedless and Navelina) were collected from commercial nurseries in Mazandaran and Kerman provinces of Iran. Samples were placed in labeled plastic bags and stored on ice until return to the laboratory. Leaf's midribs were grinded in liquid nitrogen in a pre-chilled mortar and pestle followed by extraction of nucleic acid with DNA extraction kit according to manufacturer (Fermentas, Vilnius, Lithuania).

Infected citrus samples with *Spiroplasma citri*, *Candidatus* phytoplasma aurantifolia and *Candidatus* Liberobacter asiaticus were obtained from plant protection organization of Iran and used as Positive controls.

Detection

All of plants were visually inspected for the presence of phytoplasma related disease in the field. Samples were analyzed with Dienes' staining method. Hand cut of tissues transferred to Dienes' stain for 10 min. The stain was prepared by dissolving 2.5 g methylene blue, 1.25 g azure II, 10 g maltose and 0.25 g sodium carbonate (Na₂CO₃) in 100 mL distilled water filter through filter paper and dilute to 0.2% (v/v) in distilled water (Deeley *et al.*, 1979; Musetti, 2013) The stained sections were later washed in distilled water and mounted in distilled water on a glass slide and examined microscopy.

Molecular detection and identification of phytoplasma was done by direct PCR, using primer pair P1/P7 (Duduk *et al.*, 2013) or by single tube nested PCR in a 50 µl reaction volume containing 50 ng DNA extract, 0.4 mM of each dNTPs, 1x PCR buffer, 3 mM MgCl₂, 3U Taq DNA polymerase (Fermentas, Vilnius, Lithuania) and 0.1 µM of each primers P1/P7 (Deng and Hiruki, 1991) and 0.3 µM of each primers fU5/rU3 (Lorenz *et al.*, 1995).

The PCR temperature profile was 94 °C (pre-denaturation) for 4 min and consisted of 40 cycles at 94 °C (denaturation) for 35s, 53 °C (annealing) for 35s, 72 °C (extension) for 45s, and final extension at 72 °C for 5 min.

Detection of *Candidatus* Liberobacter asiaticus and *Spiroplasma citri* were performed by specific primers respectively OI1/OI2 and A2/J5 (Hocquellet *et al.*, 1999), P89-r/f and P58-6f/4r (Yokomi *et al.* 2008).

Multiplex PCR was used for detection of phytoplasma, *Candidatus* Liberobacter asiaticus and *Spiroplasma citri* in one reaction. Primers P1/P7, A2/J5 and p58-6f/4r were used in one reaction for amplifying

DNA template of mixed positive controls. Multiplex PCR was performed in a volume of 150 μ l containing 150 ng template DNA, 1 mM of each dNTPs, 1x PCR buffer, 9 mM MgCl₂, 9 U Taq DNA polymerase (Fermentas, Vilnius, Lithuania) and 0.4 μ M of each primers P1/P7 and 0.4 μ M of each primers P58-6f/4r and 0.4 μ M of each primers A2/J5.

The PCR temperature profile was 94 °C (pre-denaturation) for 5 min and consisted of 40 cycles at 94 °C (denaturation) for 45s, 52 °C (annealing) for 45s, 72 °C (extension) for 1 min, and final extension at 72 °C for 10 min.

Approximately 20 μ l of each reaction mixture was electrophoresed in a 1.5% agarose gel and the products were visualized using UV transillumination after staining by ethidium bromide.

RFLP analysis

Ten microliters (approximately 200 ng of DNA) of direct PCR products of P1/P7 from phytoplasma infected samples were separately digested using 2 units of restriction endonucleases *TaqI*, *AluI*, *BamHI* and *MseI* in buffers supplied by the manufacturer (Fermentas, Vilnius, Lithuania). The restriction patterns were electrophoresed in a 2% agarose gel and visualized by staining with ethidium bromide and UV illumination.

Results

Dienes' staining as preliminary method verified the presence of phytoplasma in plant with deficiency symptoms. The cells of Phytoplasma can be observed as purple spots scattered in sieve tubes (Fig 1).

Primers OI1/OI2, A2/J5, P89-r/f, P58-6f/4r, P1/P7 and fU5/rU3 that used in this study respectively amplify 1160, 703, 707, 450, 1800 and 876 bp products from genome of *Ca. l. asiaticus*, *S. citri* and *Ca. Ph. aurantifolia* as positive controls (Fig 2).

The PCR results indicated that only phytoplasma and *Spiroplasma citri* infection is widespread in some of the citrus varieties. (Fig3). No amplicon was obtained for *Ca. L. asiaticus* in 360 samples collected from citrus nurseries.

Multiplex PCR assay successfully detected phytoplasma, *S. citri* and *Ca. L. asiaticus* from mixed positive control samples in one reaction (fig 4).

In young trees that exhibited yellowing and deficiencies symptoms, the presence of phytoplasma was indicated in 37/360 samples when using single tube nested PCR (Fig. 3). Lime witches broom as positive control and infected samples showed the 1800 and 876 bp fragment using P1/P7 and fU5/rU3 in one reaction, respectively.




Figure 1 Micrograph of hand cut stems of citrus stained with Dienes' stain. Section of healthy plant with unstained phloem (A), Phloem cells are stained in section obtained from infected plant (B).




Figure 2 PCR amplification products of *Ca. Liberobacter asiaticus*, *Spiroplasma citri* and *Ca. Phytoplasma aurantifolia* as positive controls with Primers OI1/OI2, A2/J5, P89-r/f, P58-6f/4r, P1/P7 and fU5/rU3, NC: health citrus sample, L: 1kb DNA ladder.




Figure 3 Detection of phytoplasma infection, using primer pairs P1/P7 and fU5/rU3 (1800 and 876 bp) in single tube reaction (A). Detection of *Spiroplasma citri*, using P58-6f/4r (450 bp), from extracted DNA of citrus varieties (B). (1: Okitsu, 2: Hoshimoto, 3: lime bear, 4: Navelate, 5: Delta seedless, 6: Fukumoto, 7: Washington navel). PC: *Candidatus Phytoplasma aurantifolia* and *Spiroplasma citri* infected samples used as positive controls, NC: health citrus sample, L: 1kb DNA ladder.

Phytoplasma infection observed in five of Navelate, six of Okitsu, four of Fukumoto, four of Valencia late, three of Washington navel foyus, four of Hoshimoto, four of Washington navel, one of Clementine clime new less1, three of Clemantis manitol, one of Lime Bears, five of Delta seedless and two of Navelina samples.

PCR products of P1/P7 were used in RFLP analysis (Fig 5). Following *MseI*, *AluI*, *BamHI*

and *TaqI* digestion, restriction pattern of phytoplasma isolate showed the same pattern with aster yellows phytoplasma as one of the members of 16Sr I group (Lee *et al.* 1998).

Stubborn disease have been identified in 17/360 citrus samples using PCR method by P89-r/f and P58-6f/4r primer pairs. Infection of *Spiroplasma citri* were observed in eight of Fukumoto, one of Navelina, five of Navelate and three of Valencia late samples.

Discussion

The multiplex PCR assay that used in present study, successfully detected *Spiroplasma citri* and *Candidatus Liberobacter asiaticus* and phytoplasma infections in one reaction. The multiplex PCR assay was used as a faster and cheaper detector of the fastidious prokaryotes in citrus as compared with standard PCR. These results represent an important step towards routine diagnosis of some citrus pathogens in laboratory, and an additional aid for certification with respect to these agents.

PCR-RFLP detected phytoplasma positive sample belong to Aster yellows group. Typical symptoms of phytoplasma infection with little leaf and proliferation were not observed in infected samples so may be, detected phytoplasma is mild strains or citrus varieties were not very susceptible host for these phytoplasma strains. On other hand there is evidence that phytoplasma concentration can vary greatly in infected plants (Berges *et al.* 2000). Sieve tubes in susceptible hosts allows to pathogen to multiply and spread along tube. In

contrast, in some citrus genotypes or varieties phytoplasma titre is extremely low and only non-specific yellowing symptoms can be developed. (Kartte and Seemuller 1991). Probably in these citrus varieties due to inability of phytoplasma to increase concentration, typical symptoms were not observed because phytoplasma concentration is usually regarded as important pathogenicity factor (Marcone 2010).

The citrus cultivars showing the yellowing and mottling symptoms caused by "phytoplasma" have been observed earlier in citrus orchards of China (Chen *et al.*, 2009), Brazil (Teixeira, 2008), Pakistan (Mannan, *et al.*, 2010), Saudi Arabia (Alhudaib *et al.*, 2009).

Confused of the symptoms be prompted by nutrient deficiencies or vascular limited pathogen such as phytoplasma, *Candidatus Liberobacter sp.* and *Spiroplasma citri* in young trees was reported (Bove, 2006; Weintraub and Jones 2010; Silva-Stenico *et al.* 2009). So for sanitary selection and certification program, using sensitive detection method is recommended.




Figure 4 Ethidium bromide-stained gel of Multiplex PCR and standard PCR. Detection of phytoplasma. (P1/P7 primers), *Spiroplasma citri* (P58-6f/4r primers) and *Candidatus Liberobacter asiaticus* (A2/J5 primers) by multiplex PCR (lane 1) and in separate reactions (lane 3, 4 and 5). L: 1kb DNA ladder.




Figure 5 RFLP analysis of phytoplasma isolated from orange. P1/P7 product was digested with *MseI*, *BamHI*, *TaqI* and *AluI*. Lane M, 1kb DNA ladder.

Our investigation determined co-infection of Spiroplasmas and Phytoplasma in different citrus varieties. Infection of many varieties of sweet orange with different range of susceptibility to *S. citri* was reported (Nejat *et al.*, 2007). Since citrus is not only host plant of *S. citri* and transmission of *S. citri* occurs by vectors and propagative material (Nejat *et al.*, 2011), stubborn disease widely distributed in citrus culture area in Iran.

References

- Alhudaib, K., Arocha, Y., Wilson, M. and Jones, P. 2009. Molecular identification, potential vectors and alternative hosts of the phytoplasma associated with a lime decline disease in Saudi Arabia. *Crop Protection*, 28: 13-18.
- Berges, R., Rott, M. and Seemüller, E. 2000. Range of phytoplasma concentrations in various plant hosts as determined by competitive polymerase chain reaction. *Phytopathology*, 90: 1145-1152.
- Bertolini, E., Olmos, A., Lopez, M. M. and Cambra, M. 2003. Multiplex nested reverse transcription-polymerase chain reaction in a single tube for sensitive and simultaneous detection for RNA viruses and *Pseudomonas savastanoi* pv. *savastanoi* in Olive trees, *Phytopathology*, 93: 286-292.
- Bové, J. M., Danet, J. L., Bananej, K., Hassanzadeh, N., Taghizadeh, M., Salehi, M. and Garnier, M. 2000. Witches' broom disease of lime in Iran. In: Proceedings of the fourteenth Conference of IOCV. IOCV, Riverside (US), pp. 207-212.
- Bové, J. M. 2006. Huanglongbing: A destructive, newly-emerging, century-old disease of citrus. *Journal of Plant Pathology*, 88: 7-37.
- Chen, J., Pu, X., Deng, X., Liu, S., Li, H. and Civerolo, E. 2009. A phytoplasma related to *Candidatus Phytoplasma asteris* detected in citrus showing Huanglongbing (yellow shoot disease) symptoms in Guangdong, P. R. China. *Phytopathology*, 99: 236-242.
- Da Graça, J. V. 1991. Citrus greening disease. *Annual Review of Phytopathology*, 29: 109-136.
- Deeley, J., Stevens, W. A. and Fox, R. T. V. 1979. Use of Diens Stain to detect plant diseases induced by mycoplasma-like organisms. *Phytopathology*, 69: 1169-1171.
- Deng, S. and Hiruki, C. 1991. Amplification of 16S rRNA genes from culturable and nonculturable mollicutes. *Journal Microbiology methods*, 14: 53-61.
- Duduk, B., Paltrinieri, S., Lee, I-M., Bertaccini, A., 2013. Nested PCR and RFLP analysis based on the 16S rRNA gene. *Methods in Molecular Biology*, 938: 159-171.

- Fahmeed, F., Arocha Rosete, Y., Acosta Perez, K., Boa, E. and Lucas J. 2009. First report of 'Candidatus Phytoplasma asteris' (Group 16SrI) infecting fruits and vegetables in Islamabad, Pakistan. *Journal of Phytopathology*, 157: 639-341.
- Frison, E. A. and Taher, M. N. 1991. FAO/IBPGR Technical Guidelines for the safe movement of citrus germplasm. Food and Agriculture Organization of the United Nations, Rome /International Board for Plant Genetic Resources, Rome.
- Gundersen, D. E. and Lee, L. M. 1996. Ultra-sensitive detection of phytoplasmas by nested PCR assays using two universal primers pair. *Phytopathologia Mediterranea* 35: 144-151.
- Heinrich, M., Botti, S., Caprara, L., Arthofer, W., Strommer, S., Hanzer, V., Katinger, H., Bertaccini, A. and Machalo, M. L. C. 2001. Improved detection methods for fruit tree phytoplasmas. *Plant Molecular Biology Reporter*, 19: 169-179.
- Hocquellet, A., Toorawa, P., Bove, J. M. and Garnier, M. 1999. Detection and identification of the two Candidatus Liberibacter sp. Associated with citrus huanglongbing by PCR amplification of ribosomal protein genes of beta operon. *Molecular and Cellular Probes*. 13: 373-379.
- Hung, T. H., Wu, M. L. and Su, H. J. 1999. Development of a rapid method for the diagnosis of citrus greening disease using the polymerase chain reaction. *Journal of Phytopathology*, 147: 599-604.
- Karte, S. and Seemüller, E. 1991. Histopathology of apple proliferation in *Malus taxa* and hybrids of different susceptibility. *Journal of Phytopathology*, 131: 149-160.
- Lee, I. M., Gundersen, R. D. E., Davis, R. E. and Bartozyk, I. M. 1998 Revised classification scheme of phytoplasmas based on RFLP analysis of 16S rRNA and ribosomal protein gene sequences. *International journal of systematic bacteriology*, 48: 1153-1169.
- Lorenz, K. H., Schneider, B., Ahrens, U., Seemüller, E. 1995. Detection of the apple proliferation and pear decline phytoplasmas by PCR amplification of ribosomal and nonribosomal DNA. *Phytopathology*, 85: 771-776
- Mannan, S., Ibrahim, M., Nadeem Chohan, S. and Holford, P. 2010. Detection of phytoplasma in citrus orchards of Pakistan *Journal of pathology*, 2: 49-54.
- Marcone, C. 2010. Movement of Phytoplasmas and the Development of Disease in the Plant in: Weintraub P, Jones P, eds, 2010. *Phytoplasmas: genomes, plant hosts and vectors*. allingford, UK, CABI Publishing.
- McCoy, R. E., Caudwell, A., Chang, C. J., Chen, T. A., Chiykowski, L. N., Cousin, M. T., Dale, J. L., de Leeuw, G. T. N., Golino, D. A., Hackett, K. J., Kirkpatrick, B. C., Marwitz, R., Petzold, H., Sinha, R. C., Sugiura, M., Whitcomb, R. F., Yang, I. L., Zhu, B. M. and Seemüller, E. 1989. Plant diseases associated with mycoplasma-like organisms. In: Whitcomb RF, Tully JG. (Eds.), *The Mycoplasmas*. Vol. V. Academic Press/ San Diego, California, pp 545-640.
- Mohkami, A., Satari, R., Lori, Z., Ehsani, A. and Nazemi, A. 2011. First report of citrus huanglongbing in the Orzooiyeh region in Kerman province. *Iranian Journal of Plant Pathology*. 47: 29 (In Fasi With English Summary).
- Musetti, R. 2013. *Phytoplasma*, methods in molecular biology. 938: 109-113.
- Nejat, N., Salehi, M., Fayyazi, M. and Izadpanah, K. 2007. Survey of sweet orange cultivars for stubborn disease resistance in Iran. *Bulletin of Insectology*, 60: 305-306.
- Nejat, N., Vadamalai, G., Sijam, K. and Dickinson, M. 2011. First report of *Spiroplasma citri* associated with periwinkle lethal yellows in southeast Asia. *Plant Disease*, 95: 1312-1320.
- OEPP/EPPO, 2006. *Candidatus Phytoplasma aurantifoliae*. OEPP/EPPO Bulletin. 36: 117-119.

- Raju, B. C., Nyland, G., Backus, E. A. and McLean, D. L. 1981. Association of Spiroplasma with brittle root of horseradish. *Phytopathology*, 71: 1067-1072.
- Salehi, M., Izadpanah, K., Siampour, M., Bagheri, A. and Faghihi, S. M. 2007. Transmission of *Candidatus Phytoplasma aurantifolia* to Bakraee (Citrus reticulata hybrid) by feral leafhopper *Hishimonus phycitis* in Iran. *Plant Disease*, 91: 466.
- Seemuller, E., Garnier, M. and Schneider, B. 2002. Mycoplasmas of plants and insects. In: Razin S, Herrmann R. (Eds.), *Molecular Biology and Pathology of Mycoplasmas*. Academic/Plenum Kluwer, London, pp 91-116.
- Silva-Stenico, M. E., Pacheco, F. T. H., Pereira-Filho, E. R., Rodrigues, J. L. M., Souza, A. N., Etchegaray, A. D., Gomes, J. E. and Tsai, S. M. 2009. Nutritional deficiency in citrus with symptoms of citrus variegated chlorosis disease. *Brazilian Journal of Biology*, 69 (3): 859-864.
- Teixeira, D. C., Wulff, N. A., Martins, E. C., Kitajima, E. W, Ayres, A. J., Eveillard, S., Saillard, C. and Bove, J. M. 2008. A phytoplasma closely related to the pigeon pea witches' broom phytoplasma (16SrIX) is associated with citrus huanglongbing symptoms in the state of São Paulo, Brazil. *Phytopathology*, 98: 977-984.
- Weintraub, P. and Jones. P, eds, 2010. *Phytoplasmas: genomes, plant hosts and vectors*. allingford, UK, CABI Publishing.
- Yokomi, R. K., Mello, A. F. S., Saponari, M. and Fletcher, J. 2008. Polymerase chain reactionbased detection of *Spiroplasma citri* associated with citrus stubborn disease. *Plant Disease*, 92: 253-260.

تشخیص مولکولی فیتوپلاسمای گروه زردی مینا در ارقام مختلف مرکبات با استفاده از PCR چندگانه

کبری مسلم خانی^۱، راحله شهبازی^۱، ساره بقایی راوری^۲ و فاطمه خلقتی بنا^۱

۱- مؤسسه تحقیقات ثبت و گواهی بذر و نهال، کرج، ایران.

۲- دانشگاه فردوسی مشهد، مشهد، ایران.

* پست الکترونیکی نویسنده مسئول مکاتبه: Moslemkhani@yahoo.com

دریافت: ۲۳ مهر ۱۳۹۲؛ پذیرش: ۲۴ دی ۱۳۹۲

چکیده: سلامت برخی از نهالستان‌های مرکبات واقع در استان‌های کرمان و مازندران به عنوان مناطق اصلی پرورش مرکبات در ایران ارزیابی گردید. ارقام مختلف با علائم غیرطبیعی از نظر آلودگی به بیماری‌های آوندی در مرکبات بررسی شدند. براساس مشاهدات چشمی و آزمایشات مولکولی ۳۶۰ نمونه، تنها آلودگی به عوامل فیتوپلاسمایی و *S. citri* محرز گردید. با استفاده از پرایمرهای P1/P7 و FU5/RU3 (در PCR آشیانه‌ای یک مرحله‌ای) آلودگی فیتوپلاسمایی ردیابی شد. همچنین آلودگی به *S. citri* با استفاده از پرایمرهای P89-R/F و P58-6F/4R مورد آزمون قرار گرفت. نتایج هضم محصول PCR به وسیله آنزیم‌های *TaqI*، *BamHI*، *AluI* و *MseI* نشان داد فیتوپلاسمای جدا شده متعلق به گروه زردی مینا است. همچنین با ارزیابی‌های صورت گرفته هیچ نوع آلودگی *Candidatus Liberobacter asiaticus* مشخص نگردید. تمایز علائم بیماری‌های محدود به آوندها خصوصاً زمانی که غلظت پاتوژن اندک است با علائم کمبود بسیار دشوار است لذا در خصوص برنامه‌های گواهی سلامت استفاده از روش‌های تشخیصی دقیق ضروری است. روش PCR چندگانه که در این مطالعه موفق به ردیابی هم‌زمان *Candidatus Liberobacter asiaticus*، *Spiroplasma citri* و *phytoplasma* در یک واکنش گردید به منظور استفاده در برنامه‌های گواهی سلامت توصیه می‌گردد.

واژگان کلیدی: بیماری، مرکبات، فیتوپلاسم و PCR چندگانه