J. Crop Prot. 2015, 4 (3): 267-276_





Laboratory survey on biological and demographic parameters of *Cryptolaemus montrouzieri* (Mulsant) (Coleoptera: Coccinellidae) fed on two mealybug species

Gholam Ali Abdollahi Ahi¹, Ali Afshari^{1*}, Valiollah Baniameri², Hemmat Dadpour³, Mohsen Yazdanian¹ and Ali Golizadeh⁴

1. Department of Plant Protection, Faculty of Plant Production, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran.

2. Department of Agricultural Entomology, Plant Protection Research Institute, Tehran, Iran.

3. Biological Control Research Laboratory, Plant Protection Research Institute, Amol, Iran.

4. Department of Plant Protection, Faculty of Agriculture, University of Mohaghegh Ardabili, Ardabil, Iran.

Abstract: Biological and demographic parameters of the mealybug ladybird, Cryptolaemus montrouzieri (Mulsant) were evaluated on citrus mealybug, Planococcus citri (Risso) and obscure mealybug, Pseudococcus viburni (Signoret) under 24 ± 2 °C, $80 \pm 5\%$ relative humidity and a photoperiod of 16:8 (L: D) h. The mealybugs had a significant effect on developmental time and reproductive and demographic parameters of the ladybird. Life span (egg to adult death) was obtained 220.85 ± 5.78 and 119.44 ± 2.1 days when fed on Pl. citri and Ps. viburni, respectively. However, mean number of eggs per female (fecundity) and mean percent of egg hatching were significantly higher on Ps. viburni than on Pl. citri. In addition, values of intrinsic rate of increase (r_m) , finite rate of increase (λ) and net reproductive rate (R_0) were significantly higher on Ps. viburni than on Pl. citri. The values of intrinsic rate of increase were estimated 0.081 and 0.094 day⁻¹ on *Pl. citri* and *Ps. viburni*, respectively. Results of this study suggested that the obscure mealybug is a more suitable prey than the citrus mealybug as the ladybird displayed shorter developmental time, and higher fecundity and growth rate when fed with obscure mealybug.

Keywords: Life table, Mealybug ladybird, Pseudococcidae

Introduction

Mealybugs (Hemiptera: Pseudococcidae) are an important group of plant pests worldwide (Williams, 1985; Ben-Dov, 1994; Miller *et al.*, 2002; Miller *et al.*, 2005) whose feeding on host tissues may cause leaf yellowing, defoliation, reduced plant growth, and in some cases, plant death (Hill, 1983). The citrus mealybug,

Planococcus citri (Risso), is a polyphagous pest that attacks a wide range of crops and ornamental plants in tropical and subtropical regions (Hill, 1983; Ben-Dov, 1994; Miller, 2005). The obscure or tea mealybug, *Pseudococcus viburni* (Signoret) has been reported as a vineyard and pome fruit orchard pest s in New Zealand, United States and South Africa (Charles *et al.*, 2010; Varela *et al.*, 2006; Mudavanhu, 2009).

In Iran, mealybugs are prevalent greenhouse and orchard pests which cause significant damage and yield loss each year (Behdad, 1997). Mafi Pashakolaei (1997) reported the

Handling Editor: Dr. Ali Asghar Talebi

^{*}**Corresponding author**, e-mail: Afshari@gau.ac.ir Received: 1 November 2014, Accepted: 13 March 2015 Published online: 8 April 2015

activity of five mealybug species on citrus and other host plants in northern Iran, among which *Pl. citri* was determined to be the dominant species. Abbasipour and Taghavi (2007) confirmed the occurrence and distribution of *Ps. viburni* in tea orchards of northern Iran, while in southern provinces of Iran, the spherical mealybug, *Nipaecoccus viridis* (Newstead), has been reported as the dominant mealybug species on host plants including citrus (Ghanbari *et al.*, 2012). Despite increased concerns about the prevalence, damage, seasonal outbreak, and economic impact of mealybugs in Iran our information in this regard is very limited.

Recommended practices for controlling mealybugs using insecticides are complicated by the fact that they hide in bark crevices and other inaccessible places and secrete thick layers of protective wax (Hill, 1983; Gill et al., 2012). As an alternative, biological control by predators could act as an efficient method of control (Simmonds et al., 2000, Mustu et al., 2008). The mealybug ladybird, Cryptolaemus montrouzieri (Mulsant) is one of the most important biological control agents that preys upon a wide range of mealybug species in many parts of the world (Babu and Azam, 1987; Heidari and Copland, 1992; Simmonds et al., 2000). This predator, which has also been reported from northern Iran, has been shown to play an important role in the population suppression of mealybugs in citrus and tea orchards (Mafi Pashakolaei, 1997; Mafi Pashakolaei et al., 2011).

Life table study of natural enemies on various species provides useful prey information for understanding their ability to control pests and provides information for improving their mass rearing feasibility (Hassell, 1978; Carey and Vargas, 1985; Carey, 1993). There are various studies on life history, life table, and behavior of C. montrouzieri, feeding on citrus mealybug and pink mealybug, Maconellicoccus hirsutus Green as prey (e.g. Persad and Khan, 2002; Özgökçe et al., 2006; Elsherif et al., 2010; Ghorbanian et al., 2011; Villegas-Mendoza et al., 2012), but our knowledge about the biology and demography of *C. montrouzieri* feeding on obscure mealybug is scant. The main objective of this study is to construct the age specific life table and demographic parameters of *C. montrouzieri* and compare their biology on citrus and obscure mealybugs, under laboratory conditions.

Materials and Methods

Laboratory rearing and maintenance of mealybugs and ladybird

Pl. citri and *Ps. viburni* were originally collected from a citrus orchard in Amol (52[°] 22[′] 50[″] E and 36[°] 26[′] 27[″] N) and a tea orchard in Nashtarood (51[°] 0[′] 37[″] E and 36[°] 45[′] 15[″] N), Mazandaran, northern Iran, respectively and then reared on squash (*Cucurbita maxima* Duchesne) fruits and potato (*Solanum tuberosum* L.) tuber buds in a climate-controlled room, 26 ± 2 °C, $80 \pm 5\%$ RH, and a photoperiod of 16L: 8D h.

The *C. montrouzieri* colony was originally provided by Tea Research Centre, Nashtarood, Mazandaran, Iran, and reared on infested squash fruits and potato tuber buds under the above-mentioned conditions. Prior to initiating feeding experiments on each prey (*Pl. citri* or *Ps. viburni*), the ladybird was reared for three generations on the corresponding prey to experience the same nutrient and physiological conditions.

Biological and demographic parameters

All experiments were conducted under similar conditions in an incubator at $24 \pm 2^{\circ}$ C, $80 \pm 5\%$ RH, and a photoperiod of L16:D8 h. Life table experiments were conducted using Carey's method (Carey, 1993). In order to obtain an egg cohort, 8 pairs of ladybirds were placed into two separate Petri dishes (10cm diameter) containing different developmental stages of mealybugs. One hundred eggs of C. montrouzieri, laid within a 24h period, were randomly chosen and transferred individually to a new Petri dishand provided with an ad libitum mixed diet of different developmental stages of mealybugs. Developmental and survival data of

Abdollahi Ahi et al. _

the ladybirds were recorded daily. After adult ladybirds emerged, females were paired with males of the same age and each pair was individually transferred to a separate Petri dish containing different developmental stages of the prey (37 and 41 pairs of adult ladybird in case of *Pl. citri* and *Ps. viburni*, respectively). Survival and fecundity data of the paired adults were recorded daily until the death of female individuals.

Statistical Analyses

The data obtained for durations of developmental stages of ladybird on two prey species were analyzed using Student's t-test (Proc t-test, SAS Institute, 2007). Demographic parameters were compared using the Jackknife method (Maia et al., 2000). The age-specific survival (l_r) , fecundity (m_r) , and life expectancy (e_x) and demographic parameters including intrinsic rate of increase (r_m) , finite rate of increase (λ), net reproduction rate (R_0), doubling time (DT) and the mean generation time (T) were calculated according to Carey (1993, 2001). The intrinsic rate of increase was estimated by solving iteratively Euler's equation: $\int_{\sum e^{-rx} l_x m_x = 1}^{\theta}$ (Carey, 1993). The mean generation time was defined as the length of

time required for a given population to complete one generation and calculated using $T = \ln R_0 / r_m$ equation (Carey, 1993).

Results

Mealybug species had a significant effect on developmental time of ladybird stages and instars including egg, larvae I, II and III, total larval, prepupa and pupae as well as on preoviposition, oviposition, and postoviposition periods (Table 1). In contrast, the duration of larval instar IV of the predator was not significantly different while feeding on the two mealybug species (t = 1.46, DF = 1,154, P > 0.05).

Mean developmental time for different stages or periods of ladybird fed on two mealybug species are illustrated in Table 2. Developmental times of most stages as well as periods of pre-oviposition, oviposition, and post-oviposition were significantly longer on *Pl. citri* than on *Ps. viburni*. Total life span (egg to adult death) was obtained 220.85 \pm 5.78 and 119.44 \pm 2.1 days on *Pl. citri* and *Ps. viburni*, respectively. Longevity of female ladybirds fed with *Pl. citri* (183.53 \pm 5.78 days) was significantly higher than those that were fed with *Ps. viburni* (78.50 \pm 2.1 days). Ovipositional period on *Pl. citri* (73.73 \pm 5.72 days) was also longer than that on *Ps. viburni* (41.32 \pm 0.23 days).

Table 1 The results of *t*-student analysis for durationsof developmental stages/periods of *Cryptolaemusmontrouzieri* fed on two mealybug species.

Developmental	t	df_{te}^{1}	Pvalue
stage/period		ι, υ	vanae
Egg incubation	159.96	1, 162	< 0.0001
Larva I	175.60	1, 156	< 0.0001
Larva II	56.36	1, 156	< 0.0001
Larva III	32.97	1, 155	< 0.0001
Larva IV	1.46	1, 154	< 0.2295
Total larval period	90.06	1, 156	< 0.0001
Prepupa	56.31	1, 154	< 0.0001
Pupa	703.75	1, 153	< 0.0001
Pre-ovipositional period	46.51	1, 152	< 0.0001
Ovipositional period	35.61	1,76	< 0.0001
Post- ovipositional period	141.70	1,73	< 0.0001

 1 df_{t,e} indicates values of degree of freedom for treatment and error, respectively.

Means of reproductive parameters of *C.* montrouzieri fed on two mealybug species are presented in Table 3. Despite longer oviposition period, all reproductive parameters including mean number of eggs per female, mean number of hatched eggs, mean percent of egg hatching, mean number of male offspring, and mean number of female offspring were higher on *Ps. viburni* than on *Pl. citri*. Mean number of eggs laid during whole life time of a female ladybird was estimated to be 278.62 ± 0.21 and 385.33 ± 0.84 when fed with *Pl. citri* and *Ps. viburni*, respectively. The oviposition rate of *C. montrouzieri* fed on the two mealybug species are shown in Fig. 1. The majority of eggs on both prey species were laid on early days of oviposition period; the highest number of eggs was laid 3 and 7 days after oviposition began when ladybirds were fed with *Pl. citri* and *Ps. viburni*, respectively.

All demographic parameters of *C. montrouzieri* including intrinsic rate of increase (r_m) , finite rate of increase (λ) , net reproduction rate (R_0) , mean generation time (T), and doubling time (DT) were significantly affected by mealybug species (Table 4). Values of r_m , λ and R_0 were statistically higher on *Ps. viburni* than on *Pl. citri*. Values of intrinsic rate of increase as the most important demographic parameter were estimated 0.081 and 0.094 day⁻¹ on *Pl. citri* and *Ps. viburni*, respectively. In contrast, values of *T* and *DT* were significantly higher on *Pl. citri* than on *Ps. viburni* and were estimated 57.11, 8.52 and 54.57, 7.37 days, respectively.

Age-specific survival (l_x) and life expectancy (e_x) rates of *C. montrouzieri* fed on two mealybug species are displayed in Fig. 2. According to the age specific life table constructed based on a cohort of 100 newly laid eggs, survival rate of egg, larval, and pupal stages of ladybird on *Pl. citri* and *Ps. viburni* were estimated 76.0, 96.05, 98.6 percent and 87.0, 95.4, 100 percent, respectively. In other words, 29.35 and 17.60 percent of the ladybird mortality (fed on *Pl. citri* than *Ps. viburni*, respectively) occurred during immature stages.

The maximum life expectancy (*ex*) of ladybirds, fed on both *Pl. citri* and *Ps. viburni*, was obtained by first instar larvae (196.13 and 100.52 days, respectively). Life expectancy of newly emerged fourth instar larvae fed on *Pl. citri* and *Ps. viburni* were estimated 187.13 and 93.63 days, respectively, while the values of life expectancy for newly emerged female ladybirds fed on these mealybugs were calculated 170.48 and 68.7 days, respectively.

 Table 2 Developmental time of different stages of Cryptolaemus montrouzieri fed on two mealybug species, Planococcus citri and Pseudococcus viburni.

Stage/Period	Developmental time (days) (Mean \pm SE) ¹	
	Planococcus citri	Pseudococcus viburni
Egg	6.66 ± 0.05 a (<i>n</i> = 79)	5.58 ± 0.07 b (<i>n</i> = 85)
Larva I	5.15 ± 0.08 a (<i>n</i> = 73)	3.35 ± 0.10 b (<i>n</i> = 85)
Larva II	3.33 ± 0.07 a (<i>n</i> = 73)	2.62 ± 0.06 b (<i>n</i> = 85)
Larva III	2.74 ± 0.06 b (n = 73)	3.25 ± 0.06 a (<i>n</i> = 84)
Larva IV	4.27 ± 0.07 a (<i>n</i> = 73)	4.43 ± 0.10 a (<i>n</i> = 83)
Total larval period	15.50 ± 0.06 a (73)	13.64 ± 0.10 b (82)
Prepupa	3.80 ± 0.04 a (<i>n</i> = 73)	3.01 ± 0.09 b (<i>n</i> = 83)
Pupa	11.34 ± 0.10 b (<i>n</i> = 73)	18.70 ± 0.20 a (<i>n</i> = 82)
Pre-ovipositional period	7.66 ± 0.10 a (<i>n</i> = 72)	6.33 ± 0.16 b (<i>n</i> = 82)
Ovipositional period	73.73 ± 5.72 a (<i>n</i> = 37)	41.32 ± 0.23 b (<i>n</i> = 41)
Post-ovipositional period	102.14 ± 5.78 a (<i>n</i> = 36)	30.85 ± 2.14 b (<i>n</i> = 39)
Adult longevity	183.53 ± 5.78 a (<i>n</i> = 37)	78.50 ± 2.14 b (<i>n</i> = 41)
Life cycle period	220.85 ± 5.78 a (<i>n</i> = 37)	119.44 ± 2.14 b (<i>n</i> = 41)

¹ Means followed by different letters in each row are significantly different (*t*-test, p < 0.05).

Abdollahi A	Ahi et	al.
-------------	--------	-----

Table 3 Reproductive parameters of Cryptolaemus montrouzieri fed on two mealybug species, Planococcus citri and Pseudococcus viburni.

Parameter	Reproductive parameters $(Mean \pm SE)^1$		
	Planococcus citri	Pseudococcus viburni	
Number of eggs per female	278.62 ± 0.21 b (<i>n</i> = 37)	385.33 ± 0.84 a (<i>n</i> = 41)	
Number of hatched eggs	220.11 ± 0.16 b (<i>n</i> = 37)	326.14 ± 0.71 a (<i>n</i> = 41)	
Percent of egg hatching	79.00 ± 0.02 b (<i>n</i> = 37)	85.00 ± 0.01 a (<i>n</i> = 41)	
Number of male offspring	102.48 ± 0.08 b (<i>n</i> = 37)	147.15 ± 0.32 a (<i>n</i> = 41)	
Number of female offspring	117.53 ± 0.09 b (<i>n</i> = 37)	178.98 ± 0.40 a (<i>n</i> = 41)	

¹ Means followed by different letters in each row are significantly different (*t*-test, p < 0.05).



Figure 1 Age-specific oviposition rate of *Cryptolaemus montrouzieri* fed on two mealybug species, *Planococcus citri* and *Pseudococcus viburni*.



Figure 2 Age-specific survival (l_x) , and life expectancy (e_x) rates of *Cryptolaemus montrouzieri* fed on two mealybug species, *Planococcus citri* and *Pseudococcus viburni*.

Biological and demographic parameters of C. montrouzieri ____

Table 4 Population growth parameters of Cryptolaemus montrouzieri fed on two mealybug species, Planococcus
citri and Pseudococcus viburni.

Parameter	Population growth parameters $(Mean \pm SE)^1$		
	Planococcus citri	Pseudococcus viburni	
Intrinsic rate of increase (r_m) (day ⁻¹)	0.081 ± 0.001 b (n = 37)	0.094 ± 0.001 a (n = 41)	
Finite rate of increase (λ)	1.085 ± 0.001 b (n = 37)	1.099 ± 0.001 a (n = 41)	
Net reproduction rate (R_0) (female / female/generation)	103.86 ± 5.73 b (n = 37)	169.27 ± 5.98 a (n = 41)	
Mean generation time (T) (days)	57.11 ± 0.75 a (n = 37)	54.57 ± 0.14 b (n = 41)	
Doubling time (DT) (days)	8.52 ± 0.12 a (n = 37)	7.37 ± 0.05 b (n = 41)	

¹ Means followed by different letters in each row are significantly different (*t*-test, p < 0.05).

Discussion

Because of high risk of using conventional insecticides, mass rearing and releasing of C. montrouzieri has been recently considered as an effective and safe biological method for controlling mealybugs populations in tea and citrus orchards of northern and southern Iran (Mafi Pashakolaei, 1997; Mafi Pashakolaei et al., 2011; Mossadegh et al., 2008). Our findings showed that C. montrouzieri developed and reproduced well when fed on both citrus and obscure mealybug, even though a shorter life cycle and higher fecundity and intrinsic rate of increase (r_m) was observed when they fed on Ps. viburni compared to Pl. citri, Therefore obscure mealybug is suggested as a more suitable prey than citrus mealybug.

Effect of prey quality on biological and parameters demographic of ladybirds particularly aphidophagous species has been reported in many studies (e.g. Omkar and James, 2004; Tsaganou et al., 2004; Cabral et al., 2006). However, little information is available on the development and demography of C. montrouzieri, especially on obscure or tea mealybug, Ps. viburni. C. montrouzieri has been often known as a specialist predator on mealybugs and a lower developmental or reproductive rate has been reported when this predator fed on other non-specialist prevs (Hodek and Honek, 2009).

Developmental and oviposition periods of C. montrouzieri obtained in our study are close to those values that reported for this predator feeding on Pl. citri (Baskaran et al., 2002; Özgökçe et al., 2006) and other mealybug species (Mani and Krishnamoorthy, 1997; Harmeet et al., 2010), but notably higher than the values that reported for other non-specialist preys. Larval developmental time observed in our study (13.64 days on Ps. viburni) was very close to 13.92 days that was reported for this ladybird when fed on *M. hirsutus* (Green) (Hemiptera: Pseudococcidae) (Parabal and Balasubramanian, 2000), but was notably lower than values that were reported for non-specialist preys such as *Aleurodicus dispersus* R. (Hemiptera: Aleurodidae) (Mani and Krishnamoorthy, 1999), Aphis gossypii (Glover) (Hemiptera: Aphididae) and Dactylopius tomentasus (Lamarck) (Hemiptera: Dactylopiidae) (Parabal and Balasubramanian, 2000; Baskaran et al., 2002). Ghorbanian et al., (2011) reported lower values for incubation, larval, prepupal and pupal periods of C. montrouzieri fed on Pl. citri compared to our study. We attribute these discrepancies to the higher temperature used in their study (27 °C vs. °C), temperature-dependent 24 such development and demography have been reported to many of other ladybird species (e.g. Kontodimas et al., 2004; Atlihan and Chi, 2008; Eliopoulos et al., 2010; Castro et al., 2011).

Abdollahi Ahi et al. _

There are a few studies reporting the demographic parameters of *C. montrouzieri* feeding on different mealybugs species (Persad and Khan, 2002; Özgökçe *et al.*, 2006; Elsherif *et al.*, 2010; Ghorbanian *et al.*, 2011), but there are apparently no reports published regarding the demography and life table of this predator feeding on *Ps. viburni*. Values of intrinsic rate of increase (r_m) in our study (0.081and 0.094 day⁻¹ on *Pl. citri* and *Ps. viburni*, respectively) were in the range of 0.077 day⁻¹ (Elsherif *et al.*, 2010) to 0.135 day⁻¹ (Persad and Khan, 2002) that have been reported on *Phenacoccus medeirensis* Green and *M. hirsutus*, respectively.

In conclusion, our findings can be helpful from both a biological control standpoint and a mass-rearing feasibility standpoint. We showed the high potential of C. montrouzieri to feed, develop, and reproduce successfully on both Pl. citri and Ps. viburni that could eventually suppress populations of these mealybugs in citrus and tea orchards of southern and northern Iran. We also showed that the ladybirds tend to feed more on Ps. viburni than on Pl. citri, which may have consequences on the efforts to rear this predator, mass especially in commercial insectaries.

Acknowledgments

We thank Plant Protection Research Institute, Tehran, Iran and Biological Control Laboratory, Amol, Mazandaran Province, Iran, for technical support. We are also thankful to Mahmood Hasanzadeh for help with collecting and rearing of insects and Majid Ghaninia for critical review of the manuscript.

References

- Abbasipour, H., and Taghavi, A. 2007.
 Description and seasonal abundance of the tea mealybug, *Pseudococcus viburni* (affinis) (Signoret) (Hom.; Pseudococcidae) found on tea in Iran. Journal of Entomology, 4: 474-478.
- Atlihan, R., and Chi, H. 2008. Temperaturedependent development and demography of

J. Crop Prot. (2015) Vol. 4 (3)

Scymnus subvillosus (Coleoptera: Coccinellidae) reared on *Hyalopterus pruni* (Homoptera: Aphididae). Journal of Economic Entomology, 101 (2): 325-333.

- Babu, T. R., and Azam, K. M. 1987. Biology of *Cryptolaemus montrouzieri* Mulsant. (Col., Coccinellidae) in relation with temperature. Entomophaga, 32 (4): 381-386.
- Baskaran, R. K. M., Srinivasan, T. R. and Mahadeven, N. R. 2002. Life table of Australian ladybird beetle, *Cryptolaemus* montrouzieri Mulsant feeding on mealybugs, Maconellicoccus hirsutus and Dactylopius tomentosus. Indian Journal of Agricultural Sciences, 72: 54-56.
- Behdad, E. 1997. Pests of Fruit Crops in Iran. Yadbood Pub., Isfahan, Iran, 843 pp.
- Ben-Dov, Y. 1994. A Systematic Catalogue of the Mealybugs of the World (Insecta, Hom., Coccoidea, Pseudococcidae and Putoidae) with Data on Geographical Distributions, Host plants, Biology and Economic Importance. Intercept Limited, Andover, UK, 686 pp.
- Cabral, S, Soares, A. O., Moura, R. and Garcia,
 P. 2006. Suitability of *Aphis fabae*, *Myzus persicae* (Homoptera: Aphididae) and *Aleyrodes proletella* (Homoptera: leyrodidae) as prey for *Coccinella undecimpunctata* (Coleoptera: Coccinellidae) Biological Control, 39: 434-440.
- Carey, J. R. 1993. Applied Demography for Biologists with Special Emphasis on Insects. Oxford University Press, New York, 206 pp.
- Carey, J. R. 2001. Insect biodemography. Annual Review of Entomology, 46: 79-110.
- Carey, J. R. and Vargas, R. I. 1985. Demographic analysis of insect mass rearing: a case study of three tephritids. Journal of Economic Entomology, 78: 523-527.
- Castro, C. F., Almeida, L. M. and Penteado, S. R. C. 2011. The impact of temperature on biological aspects and life table of *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae). Florida Entomologist, 94(4): 923-932.
- Charles, J. G., Bell, V. A. Lo, P. L., Cole, L. M. and Chhagan, A. 2010. Mealybugs

(Hemiptera: Pseudococcidae) and their natural enemies in NewZealand vineyards from 1993-2009. New Zealand Entomologist, 33: 84-91.

- Eliopoulos, P. A., Kontodimas, D. C. and Stathas, G. J. 2010. Temperature-dependent development of *Chilocorus bipustulatus* (Coleoptera: Coccinellidae). Environmental Entomology, 39 (4):1352-1358.
- Elsherif, M. E., Al-Humari, A. A., Naser, K. S. A. and Atif, J. Y. M. 2010. Fecundity and life table of female predator *Cryptolaemus montrouzieri* on *Phenacoccus madeirensis*. Annals of Agricultural Sciences, 55 (2): 321-326.
- Ghanbari, G., Ghajariyeh, H., Alichi, M. and Kheradmand, K. 2012. A study of the population dynamics of *Nipaecoccus viridis* Newstead in Shiraz region: effective factors on population decrease. Plant Protection (Scientific Journal of Agriculture), 34 (2): 47-58.
- Ghorbanian, S., Ranjbar Aghdam, H.,
 Ghajarieh, H. and Malkeshi, S. H. 2011. Life cycle and population growth parameters of *Cryptolaemus montrouzieri* Mulsant (Col.: Coccinellidae) reared on *Planococcus citri* (Risso) (Hom.: Pseudococcidae) on coleus. Journal of the Entomological Research Society, 13 (2): 53-59.
- Gill, H. K., Goyal, G., Gillet-Kaufman, J. 2012.
 Citrus Mealybug *Planococcus citri* (Risso) (Insecta: Hemiptera: Pseudococcidae).
 Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Available on: edis.ifas.ufl.edu/in947 (Revised April 2013).
- Harmeet, K., Virk, J. S. and Rabinder, K. 2010. Biology of Australian ladybird beetle, *Cryptolaemus montrouzieri* Mulsant on *Phenacoccus solenopsis* Tinsley. Journal of Biological Control, 24: 123-125.
- Hassell, M. P. 1978. The Dynamics of Arthropod Predator- Prey System. Princeton University Press, Princeton, NJ. 237 pp.
- Heidari, M. and Copland, M. J. W. 1992. Host finding by *Cryptolaemus montrouzieri* (Col.: Coccinellidae) a predator of mealybugs

(Hom.: Pseudococcidae). Entomophaga, 37: 621-625.

- Hill, D. S. 1983. *Planococcus citri* (Rossi). Agricultural Insect Pests of the Tropics and Their Control, 2nd Edition. Cambridge University Press, 746 pp.
- Hodek, I. and Honek, A. 2009. Scale insects, mealybugs, whiteflies and psyllids (Hemiptera, Sternorrhyncha) as prey of ladybirds. Biological Control, 51: 232-243.
- Kontodimas, D. C., Eliopoulos, P. A., Stathas, G.J. and Economou, L.P. 2004. Comparative temperature-dependent development of Nephus includens (Kirsch) and Nephus bisignatus (Boheman) (Coleoptera: Coccinellidae) preying on Planococcus citri (Risso) (Homoptera: Pseudococcidae): evaluation of a linear and various nonlinear models using specific criteria. Environmental Entomology, 33 (1):1-11.
- Mafi Pashakolaei, Sh. A. 1997. Identification of mealybugs (Hom.; Pseudococcidae) in Mazandaran province and study of their dominant species and natural enemies. M. Sc. Thesis, Tarbiat Modarres University, Tehran, Iran, 112 pp.
- Mafi Pashakolaei, Sh. A., Barari, Н., R., Babaei, 2011. Valiolahpor, M. Investigation on the impact of pesticide applications in citrus orchards on the developmental stages of *Cryptolaemus* montrozieri (Mulsant), in Mazandaran Province. Proceedings of the Second Iranian Pest Management Conference (IPMC), Kerman, Iran, pp. 197-206.
- Maia, A. H. N., Luiz, A. J. B. and Campanhola, C. 2000. Statistical inference on associated fertility life table parameters, using Jackknife technique: computational aspects. Journal of Economic Entomology, 93: 511-518.
- Mani, M., Krishnamoorthy, A. 1999. Predatory potential and development of the Australian ladybird beetle, *Cryptolaemus montrouzieri* Muls. (Col.: Coccinellidae) on the spiraling whitefly, *Aleurodicus dispersus* Russel. Entomon, 24 (2): 197-198.
- Mani, M. and Krishnamoorthy, A. 1997. Australian ladybird beetle *Cryptolaemus*

Abdollahi Ahi et al. _

montrouzieri. Madras Agricultural Journal, 84: 237-249.

- Miller, D. R. 2005. Selected scale insects groups (Hemiptera: Coccoidea) in the southern of the United States. Florida Entomologists, 88 (4): 482-501.
- Miller, D. R., Miller, G. L. and Watson, G. W. 2002. Invasive species of mealybugs (Hemiptera: Pseudococcidae) of the United States and their threat to U.S. Agriculture. Proceedings of the Entomological Society of Washington, 104: 825-836.
- Miller, G. L., Miller, D. R., Hodges, G. S. and Davidson, J.A. 2005. Introduced scale insects (Hemiptera: Coccoidea) of the United States and their impact on U. S. Agriculture. Proceedings of the Entomological Society of Washington, 107: 123-158.
- Mossadegh, M. S., Eslamizadeh, R. and Esfandiari, M. 2008. Biological study of mealybug *Nipaecoccus viridis* (New.) and possibility of its biological control by *Cryptolaemus montrouzieri* Mulsant in citrus orchards of North Khuzestan. Proceeding of 18th Iranian Plant Protection Congress, Hamedan, Iran, p. 35.
- Mudavanhu, P. 2009. An investigation into the integrated pest management of the obscure mealybug, *Pseudococcus viburni* (Signoret) (Hemiptera: Pseudococcidae), in pome fruit orchards in the Western Cape Province, South Africa. M.Sc. Tthesis, University of Stellenbosch, South Africa, 96 pp.
- Muştu, M., Kilinçer, N., Ülgentürk, S. and Kaydan, M. B. 2008. Feeding behavior of *C. montrouzieri* on mealybugs parasitized by *Anagyrus pseudococci*. Phytoparasitica, 36: 360-367.
- Omkar and James, B. E. 2004, Influence of prey species on immature survival, development, predation and reproduction of *Coccinella transversalis* Fabricius (Col., Coccinellidae). Journal of Applied Entomology, 128: 150-157.
- Özgökçe, M. S., Atlihan, R. and Karaca, I. 2006. The life table of *Cryptolaemus montrouzieri*

Mulsant (Col., Coccinellidae) after different storage periods. Journal of Food, Agriculture and Environment, 4: 282-287.

- Parabal, S. and Balasubramanian, A. 2000. Feeding potential and larval development of *Cryptolaemus montrouzieri* Mulsant on aphids and mealybug. Journal of the Agricultural Science Society of North East India, 13 (1): 8-11.
- Persad, A. and Khan, A. 2002. Comparison of life table parameters for *Maconellicoccus hirsutus*, *Anagyrus kamali*, *Cryptolaemus montrouzieri* and *Scymnus coccivora*. BioControl, 47: 137-149.
- SAS Institute. 2007. PROC user's manual, version 8th ed. SAS Institute, Cary, NC.
 Simmonds, M. S. J., Manlove, J. D.,
- Simmonds, M. S. J., Manlove, J. D., Blaney,W.M. and Khambay, B.P.S. 2000. Efect of botanical insecticides on the foraging and feeding behavior of the coccinellid predator *Cryptolaemus montrouzieri*. Phytoparasitica, 28: 99-107.
- Tsaganou, F. C., Hodgson, C. J., Athanassiou, C.
 G., Kavallieratos, N. G., Tomanovie', Z.,
 2004. Effect of *Aphis gossypii* Glover, *Brevicoryne brassicae* (L.), and *Megoura viciae* Buckton (Hemiptera: Aphidoidea) on the development of the predator *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae). Biological Control, 31: 138-144.
- Varela, L. G., Rhonda, J. S., Battany, M. and Bentley, W. 2006. Grape, obscure or vine which mealybug is it; why should you care? Practical winery and vineyard, XXVII (6): 37-46.
- Villegas-Mendoza, J. M., Rivera, G. and Rosas-García, Y. N. M. 2012. Behavioral analysis of *Cryptolaemus montrouzieri* Mulsant while preying on the pink hibiscus mealybug under field conditions. Southwestern Entomologist, 37 (2):177-185.
- Williams, D. J. 1985. Australian Mealybugs. British Museums (Natural History), London, 431 pp.

J. Crop Prot.

ارزیابی پارامترهای زیستی و جمعیتی کفشدوزک Cryptolaemus montrouzieri Mulsant (Coleoptera: Coccinellidae) با تغذیه از دو گونه شپشک آردآلود در شرایط آزمایشگاهی

غلامعلى عبداللهي آهي'، على افشارى"*، ولىالله بنىعامرى'، همت دادپور''، محسن يزدانيان' و على گلىزاده ً

۱ – گروه گیاهپزشکی، دانشکده تولید گیاهی، دانشگاه علوم کشاورزی و منابع طبیعی گرگان، گرگان، ایران.
 ۲ – بخش تحقیقات حشرهشناسی کشاورزی، مؤسسهی تحقیقات گیاهپزشکی کشور، تهران، ایران.
 ۳ – آزمایشگاه تحقیقات کنترل بیولوژیک، مؤسسهی تحقیقات گیاهپزشکی کشور، آمل، ایران.
 ۴ – گروه گیاهپزشکی، دانشکده کشاورزی، دانشگاه محقق اردبیلی، اردبیل، ایران.
 * پست الکترونیکی نویسنده مسئول مکاتبه: ۲۱ سفند (معنیه)
 دریافت: ۱۰ آیان ۱۳۹۳، پذیرش: ۲۲ اسفند ۱۳۹۳

واژگان کلیدی: جدول زندگی، کفشدوزک کریپتولموس، Pseudococcidae