

Short Paper

Comparison of some natural broth media for production and virulence of *Beauveria bassiana* blastospores against the browntail moth, *Euproctis chrysorrhoea* (Lep.: Lymantriidae)

Parisa Bena-Molaei¹, Reza Talaei-Hassanloui^{1*} and Hassan Askary²

1. Department of Plant Protection, College of Agriculture and Natural Resources, University of Tehran, Karaj, Iran.

2. Plant Protection Research Institute, Tehran, Iran.

Abstract: Effects of three nutritional levels of beet root molasses, cheese permeate, wheat bran extract, rice bran extract and Sabouraud's Dextrose Broth (SDB) were evaluated for blastospore production by two isolates of *Beauveria bassiana sensu lato*. at an interval of 24 h for seven days. Depending on the isolate, maximum blastospore production was obtained in 12% rice bran extract and 20% cheese permeates on the 7th day. Both isolates produced the fewest blastospores in 4% cheese permeate. Virulence of blastospores, produced in liquid media containing beet root molasses, permeate, wheat bran extract and SDB (as control), on third instar larvae of brown tail moth *Euproctis chrysorrhoea* indicated that there were no significant differences among these nutritional media for either one of the isolates. Considering blastospore quantity and quality in terms of virulence and local accessibility, cheese permeate was found to be the best medium for mass production of *B. bassiana* blastospores.

Keywords: *Beauveria bassiana*, *Euproctis chrysorrhoea*, blastospore, virulence, natural medium

Introduction

Entomopathogenic fungi play an important role in decreasing pest insect populations. Use of these fungi as microbial control agents for pest management in large scale depends on the capability of high spore production and at reasonable costs. The ability of pathogenic fungi to grow and produce spores on artificial media is one of the main advantages in the commercial development of these fungi (Cisneros and Vera, 2000; Dalla Santa *et al.*, 2005). Approaches directed to medium optimization must consider not only spore yield but also spore qualities such

as desiccation tolerance, stability as a dry preparation and biocontrol efficacy. In fact the important part of the success of microbial control programs often depends on an adequate mass-production method (Khachatourians, 1986; Kamp and Bidochka, 2002; Jenkins *et al.*, 1998; Jenkins and Goettel, 1997).

Blastospores of mitosporic fungi produced in liquid media have potential to infect hosts (Kleespies and Zimmermann, 1992; Jenkins and Goettel, 1997). The type of growth medium and the nutritional and physical conditions of the mass production system greatly affect the number, type, stability, durability and virulence of fungal propagules (Ibrahim *et al.*, 1993; Feng *et al.*, 1994; Fargues *et al.*, 2002). There is not much information on the effects of culture media on virulence of entomopathogenic fungi (Ibrahim *et al.*, 2002; Safavi *et al.*, 2007).

Handling Editor: Saeid Moharrampour

* Corresponding author, e-mail: rtalaei@ut.ac.ir

Received: 20 December 2014, Accepted: 13 March 2015

Published online: 24 April 2015

Bena-Molaei *et al.* (2011) studied the effect of culture substrates on virulence of *Beauveria bassiana* conidia against the browntail moth *Euproctis chrysorrhoea* (Lep.: Lymantriidae). Their results showed that there were significant differences among nutritional substrates for their effects on the virulence of produced conidia. Browntail moth, is one of the most important pests of oak and broad leaf trees (Nikdel *et al.*, 2003). The objective of this research was the evaluation of some natural broth media for blastospore production of two isolates of *B. bassiana sensu lato*. We considered browntail moth larvae as a target host to test our hypothesis that growth media could affect the virulence of *B. bassiana* blastospores.

Materials and Methods

Media preparation for spore production assay

Cheap nutritional substrates were selected and evaluated for blastospore production. Substrates were beet root molasses, permeate, wheat bran extract and rice bran extract. In addition, Sabouraud's Dextrose Broth (SDB, Merck™) was considered as the control medium. Two isolates of *B. bassiana s.l.* (EUT105, of soil origin and EUT116 from a lepidopteran origin) were used. Both isolates are deposited at the fungal collection of the College of Agriculture and Natural Resources, University of Tehran.

Beet root molasses (obtained from the Naghadeh Sugar Factory, West Azerbaijan) and cheese permeate (obtained from Tabriz P-Azar Company) were dissolved in sterile distilled water and filtered through cheese cloth. Two hundred grams of each of wheat and rice bran were boiled separately in distilled water and filtered through cheese cloth. Sterile distilled water was then added to these 4 nutritional substrates to prepare final concentrations of 4, 12 and 20%. Experimental units were 250 ml Erlenmeyer flasks each containing 150 ml of one of the media. The flasks were autoclaved and then inoculated with one plug (0.5 cm) taken from actively growing fungal colonies on Sabouraud's Dextrose Agar + Yeast extract (SDAY) and then incubated at 24 ± 3 °C on a rotary shaker (GFL, Germany) at 120 rpm.

Blastospore production

Daily until 7 days post-inoculation, 1 ml of each suspension was added to 9 ml of sterile distilled water and blastospores were counted using a hemocytometer (Neubauer, Germany). This experiment was done in a factorial arrangement in a completely randomized design, each treatment had three replicates and the whole experiment was repeated once.

Bioassays with browntail moth larvae

Egg masses and first instar larvae of the browntail moth, *Euproctis chrysorrhoea* were collected from Arasbaran forests, East Azarbaijan, Iran and maintained at 25 ± 2 °C, $50 \pm 10\%$ RH and 16:8 h (L: D) photoperiod. Fresh *Ulmus* (elm) leaves were used as food. Third instar larvae were used for bioassays. Blastospores were harvested from each medium on the 7th day, filtered through two layers of cheese cloth and any possible contamination was checked. Three concentrations (10^5 , 10^6 and 10^7 blastospore/ml) of each isolate produced on each medium were prepared. Third instar larvae were randomly selected from the stock colony and inoculated with 250 µL of prepared concentrations using a sprayer, depositing approximately 2.1×10^3 bl/cm² with 10^7 blastospore/ml. Inoculated larvae were transferred with a hair brush to octagonal dishes ($4.5 \times 11 \times 11$ cm) and control larvae were treated with sterile distilled water. Treated larvae were maintained at 25 ± 1 °C, $90 \pm 10\%$ RH and a photoperiod of 16h L: 8h D, fed daily and their mortalities were recorded every other day for 10 days. Dead larvae were placed in moist Petri dishes for observation of mycosis. The viability of blastospores was always more than 95% as determined according to Goettel and Inglis (1997). This experiment was done in a factorial arrangement in a completely randomized design and each treatment had four replicates, each replicate consisted of 11 larvae and 1536 larvae were used totally. The whole experiment was conducted twice.

Data Analysis

Observed total mortality percentages were corrected for control mortality that was always less than 5% using Schneider_Orelli formula

(Püntener, 1981) and then pooled data of two-time repeats were analyzed by SYSTAT12. The General Linear Model (GLM) procedure was used to perform an ANOVA with the Tukey HSD test for means comparison. In the case of determining difference among different concentrations, regression analysis was used.

Results

Analysis of variance with the GLM procedure showed that there were significant differences in production of blastospores among nutritional media for both isolates on the 4th day (EUT105: $F_{4,30} = 5.4$, $P < 0.005$ and EUT116: $F_{4,30} = 6.8$, $P < 0.005$) and on the 7th day (EUT105: $F_{4,30} = 3.1$, $P < 0.05$ and EUT116: $F_{4,30} = 9.3$, $P < 0.001$).

On the 4th day, the highest and lowest numbers of blastospore production for EUT105

(2.2×10^8 and 7×10^7 bl/ml) were recorded for 12% wheat bran extract and 4% molasses, respectively. In *B. bassiana* EUT116, 20% permeate (2.3×10^8 bl/ml) had the highest blastospore production with no significant difference with 12% rice bran extract and 12% wheat bran extract. The lowest blastospore production (3.4×10^7 bl/ml) occurred in 4% molasses (Table 1).

Mean comparisons on the 7th day showed that with isolate EUT105, 12% rice bran extract and 4% molasses had the highest (3.5×10^8 bl/ml) and lowest (1.3×10^8 blastospore/ml) blastospore yields, respectively. But there was no significant difference among the three concentrations of each medium with this isolate using Regression analysis. With EUT116, 20% permeate gave the highest (3.3×10^8 /ml) blastospore production (Table 2).

Table 1 Quantities of blastospores produced in five different media for two isolates of *Beauveria bassiana* on 4th day post-inoculation.

Nutritional substrates	Yield (blastospores / ml) ²					
	<i>B. bassiana</i> EUT105			<i>B. bassiana</i> EUT116		
	4%	12%	20%	4%	12%	20%
Permeate	9.6×10^7 ab	1.4×10^8 a	1.8×10^8 a	1.5×10^8 a	1.8×10^8 a	2.3×10^8 a
Beetroot molasses	7.0×10^7 b	7.6×10^7 a	1.3×10^8 a	3.4×10^7 c	7.4×10^7 a	8.5×10^7 ab
Rice bran	9.6×10^7 ab	1.9×10^8 a	1.0×10^8 a	1.2×10^8 ab	2.2×10^8 a	9.0×10^7 ab
Wheat bran	1.9×10^8 a	2.2×10^8 a	1.1×10^8 a	1.2×10^8 ab	1.9×10^8 a	9.5×10^7 ab
SDB ¹	9.0×10^7 b	9.0×10^7 a	9.0×10^7 b	6.5×10^7 bc	6.5×10^7 a	6.5×10^7 b

1. Sabouraud's Dextrose Broth, only one concentration was considered for this medium.
2. Means followed by the same letters in each column are not significantly different (Tukey's test at $P < 0.05$).

Table 2 Quantities of blastospores produced in five different media for two isolates of *Beauveria bassiana* on 7th day post-inoculation.

Nutritional substrates	Yield (blastospores / ml) ²					
	<i>B. bassiana</i> EUT105			<i>B. bassiana</i> EUT116		
	4%	12%	20%	1.9×10^8 a	12%	20%
Permeate	1.3×10^8 a	2.6×10^8 a	3.2×10^8 a	6.8×10^7 b	3.2×10^8 a	3.3×10^8 a
Beetroot molasses	1.3×10^8 a	1.5×10^8 a	1.6×10^8 a	1.8×10^8 a	1.3×10^8 b	1.8×10^8 ab
Rice bran	2.8×10^8 a	3.5×10^8 a	3.0×10^8 a	2.2×10^8 a	2.9×10^8 ab	1.4×10^8 b
Wheat bran	2.7×10^8 a	3.0×10^8 a	1.7×10^8 a	1.7×10^8 a	3.3×10^8 a	2.1×10^8 ab
SDB ¹	2.4×10^8 a	2.4×10^8 a	2.4×10^8 a	1.9×10^8 a	1.7×10^8 ab	1.7×10^8 ab

1. Sabouraud's Dextrose Broth, only one concentration was considered for this medium.
2. Means followed by the same letters in each column are not significantly different (Tukey's test at $P < 0.05$).

Bioassay with third instars larvae of browntail moth

The nutritional substrates had no significant effects on blastospore virulence on browntail moth larvae for either isolate (EUT105: $F_{4,45} = 1.8$, $P > 0.05$ and EUT116: $F_{4,45} = 2.3$, $P > 0.05$). Applying 10^7 blastospore/ml with deposition rate of approximately 2.1×10^3 bl./cm², of EUT105 produced mortalities ranging from 73 to 82% and it was between 69 to 85% for EUT116 (Table 3).

Table 3 Percent mortality of the browntail moth, *Euproctis chryorrhoea* third-instar larvae caused by two isolates of *Beauveria bassiana* blastospores obtained from different culture substrates.

Nutritional substrates	% Mortality \pm SE ¹	
	<i>B. bassiana</i> EUT105	<i>B. bassiana</i> EUT116
Permeate		
Beetroot molasses	72.9 \pm 13.3	81.9 \pm 13.9
Wheat bran	77.2 \pm 10.2	79.7 \pm 9.5
Rice bran	81.8 \pm 9.2	86.2 \pm 11.0
SDB	75.5 \pm 9.7	69.3 \pm 10.7
Permeate	81.1 \pm 10.4	84.2 \pm 10.0

1. mortality at 10^7 blastospores / ml.

Discussion

For growth, *B. bassiana* needs carbon sources such as dextrose which can be replaced by polysaccharides such as starch or lipids. But the nutritional requirements of each species or isolate of entomopathogenic fungus must be considered separately (Smith and Grula, 1981; Taborsky, 1992). In our study, type of nutritional media significantly affected the production of *B. bassiana* blastospores. Increasing the concentration of permeates and molasses led to higher blastospore production, while wheat bran extract and rice bran extract didn't increase the quantity of blastospores produced. We also demonstrated that liquid culture composition influences spore production. Complementary nourishment of media with agricultural by-products enhanced chlamydospore production in liquid culture of

Fusarium oxysporum (Elzein and Kroschel, 2004). Puzari *et al.* (1997) reported that 3.9×10^8 conidia/ml of *B. bassiana* were produced using a medium of rice hulls and saw dust, whereas 1×10^9 conidia/ml were produced on molasses yeast broth (Shashi *et al.*, 1999). There is far less information for blastospore production using natural substrates.

Verhaar and Hijwegen (1993) reported production of 3×10^9 spore/ml of *Verticillium lecanii* in oat flour media. Farsi *et al.* (2003) obtained 1.02×10^8 spores/ml of *V. lecanii* in 5% molasses media. In our study, 1×10^8 - 2×10^8 and 1.5×10^8 - 3.5×10^8 blastospore/ml in our used media were produced on the 4th and 7th day, respectively. Thus, blastospore production was in optimal range. Rice bran extract, wheat bran extracts and permeates showed higher blastospore production than molasses and control. These media present good carbon and nitrogen sources and some essential macronutrients and microelements. Thus it seems that these media support better fungal growth providing nutritional requirements and good blastospore production. Media including rice bran and wheat bran are good sources for starch and permeate and molasses are very rich in sugar (Kamyab, 2009). However the amount of protein, phosphorous, magnesium, methionine and cysteine in molasses is less than in the other studied media and probably could be the reason for lower blastospore production. As in the case of *P. fumosoroseus* (Now *Isaria fumosorosea*), higher concentrations of nitrogen can increase blastospore production (Jackson, 1997; Cliquet and Jackson, 2005). Although molasses has more calcium and potassium than the other substrates used (Kamyab, 2009), there must be an equilibrium among nutritional requirements other than carbon and nitrogen; their ratios could be effective factors in production and quality of fungal spores (Jackson and Bothast, 1990; Schisler *et al.*, 1991).

pH of the medium is another important factor in spore production. Optimum pH for *B. bassiana* growth is 5.7-5.9 and for spore formation it is 7-8 (Taborsky, 1992). There is a close relationship between oxygen and moisture

content of media and decrease of available oxygen which can retard fungal growth. According to our results, increasing nutrient concentration from 12 to 20% in wheat bran extract and rice bran extract didn't result in higher blastospore production. We speculate that decreasing oxygen content could be a reason for this.

Considering factors such as availability, price and feasibility of production, extraction and filtration of wheat and rice bran, these media are more time-consuming and not cost-effective. The blastospores of *B. bassiana* isolates produced on the tested culture media demonstrated high virulence against the browntail moth larvae although there was no significant difference due to nutritional contents of the media. In contrast, Ibrahim *et al.* (2002) reported that culture media influenced fungal virulence. They showed that conidia of *M. anisopliae* grown on Sabouraud dextrose agar modified (SDAM) with KCl and minimal medium (MM) were more aggressive to *Myzus persicae* and *Meligethes aeneus* than conidia derived from Sabouraud dextrose agar (SDA) or yeast extract agar. This could be related to fungus reservoir during the growth on different nutrients. It is known that *B. bassiana* needs nutrition for penetrating insect integument and growth (St. Leger *et al.*, 1989). Thus low nutrition reservoir of blastospores or conidia can be a probable factor for low virulence.

This is the first study of *B. bassiana s.l.* blastospores against the browntail moth providing evidence for its control potential against this insect. On the basis of analysis for yield data, harvesting blastospores was dependent on the medium and isolate. For *B. bassiana* EUT105, using 12% wheat and rice bran extract after 48 hours, using 20% molasses after 72 hours and using 20% permeate after 96 hours is suggested. Whereas in the case of *B. bassiana* EUT116, the third day is the best time for blastospore harvesting from all media. In conclusion, cheese permeate medium is introduced as an appropriate medium for mass production of *B. bassiana* blastospores due to production of heavy blastospores of good

virulence to browntail moth larvae and more importantly, lower cost of spore production compared with SDB medium. As this medium is a waste product in local cheese production, it could be provided at high volume with very low cost. Introducing this production medium could be considered from the two aspects in future; researchers can attempt to optimize it and fungal-product factories can use it locally for the production of semi-massive-scale biopesticide on the basis of blastospores.

Acknowledgement

Financial supports by University of Tehran and Center of Excellence for Biological Control of Plant Pests on this project are acknowledged. We thank Research Institute of Forests and Rangelands for providing a part of research facilities.

References

- Taborsky, V. 1992. Small-Scale Processing Of Microbial Pesticides. Chapter 2. Standardization of Products. FAO Agricultural Services Bulletin No. 96. Available on: <http://www.fao.org/docrep/t0533e/t0533e04.htm>.
- Bena-Molaei, P., Talaei-Hassanloui, R. and Askary, H. 2011. Effect of culture substrates on virulence of *Beauveria bassiana* (Ascomycota: Cordycipitaceae) conidia against the browntail moth, *Euproctis chrysorrhoea* (Lepidoptera: Lymantriidae). *Biocontrol Science and Technology*, 21: 619-624.
- Cisneros, F. and Vera, A. 2000. Mass production *Beauveria brongniartii* inoculum, an economical, farm-level method. CIP Program Report, 155-160.
- Cliquet, S. and Jackson, M. A. 2005. Impact of carbon and nitrogen nutrition on the quality, yield and composition of blastospores of the bioinsecticidal fungus *Paecilomyces fumosoroseus*. *Journal of Industrial Microbiology & Biotechnology*, 32: 204-210.
- Dalla Santa, H. S., Dalla Santa, O. R., Brand, D., Vandenberghe, L. P. S. and Soccol, C. R.

2005. Spore production of *Beauveria bassiana* from agro-industrial residues. *Brazilian Archives of Biology and Technology*, 48: 51-60.
- Elzein, A. and Koroschel, J. 2004. Influence of agricultural by-products in liquid culture on chlamydospore production by the potential mycoherbicide *Fusarium oxysporum* Foxy 2. *Biocontrol Science and Technology*, 14: 823-836.
- Fargues, J., Smits, N., Vidal, C., Vey, A., Vega, F., Mercadier, G. and Quimby, P. 2002. Effect of liquid culture media on morphology, growth, propagule production, and pathogenic activity of the hyphomycete, *Metarhizium flavoviride*. *Mycopathologia*, 154: 127-138.
- Farsi, M. J., Askary, A., Talebi, Kh. and Kharazi-Pakdel, A. 2003. Pathogenicity of *Verticillium lecanii* (= *Lecanicillium muscarium*) on browntail moth, *Euproctis chrysorrhoea* (L.) larvae at laboratory condition. *Pajouhesh and Sazandegi*, 60: 52-58. (Abstract in English).
- Feng, M. G., Poprawski, T. J. and Khachatourians, G. G. 1994. Production, formulation and application of the entomopathogenic fungus *Beauveria bassiana* for insect control: current status. *Biocontrol Science and Technology*, 4: 3-34.
- Goettel, M. S. and Inglis, G. D. 1997. Fungi: Hyphomycetes. In: Lacey, L. A. (Ed.), *Manual of Techniques in Insect Pathology*. Academic Press, San Diego, pp. 213-249.
- Ibrahim, L., Butt, T. and Jenkinson, P. 2002. Effect of artificial culture media on germination, growth, virulence and surface properties of the entomopathogenic hyphomycete *Metarhizium anisopliae*. *Mycological Research*, 106: 705-715.
- Ibrahim, Y. B., Lim, T. K., Tang, M. K. and Teng, H. M. 1993. Influence of temperature, pH and selected growth media on germination, growth and sporulation of *Aschersonia placenta* and *Hypocrella raciborskii*. *Biocontrol Science and Technology* 3: 55-61.
- Jackson, M. A. 1997. Optimizing nutritional conditions for the liquid culture production of effective fungal biological control agents. *Journal of Industrial Microbiology and Biotechnology*, 19:180-187.
- Jackson, M. A. and Bothast, R. J. 1990. Carbon concentration and carbon-to-nitrogen ratio influence submerged-culture condition by the potential bioherbicide *Colletotrichum truncatum* NRRL 13737. *Applied and Environmental Microbiology*, 56: 3435-3438.
- Jenkins, N. E. and Goettel, M. S. 1997. Methods for mass-production of microbial control agents of grasshoppers and locusts. *Memories of the Entomological Society of Canada*, 171: 37-48.
- Jenkins, N. E., Heviefio, G., Langewald, J., Cherry, A. J. and Lomer, C. J. 1998. Development of mass production thechnology for aerial conidia for use as mycopesticides. *Biocontrol News and Information*, 19, 21N-31N.
- Kamp, A. M. and Bidochka, M. J. 2002. Conidium production by insect pathogenic fungi on commercially available agars. *Letters in Applied Microbiology*, 35: 74-77.
- Kamyab, A. 2009. A guide to the livestock and poultry food. Haghshenas Publications, 218 pp.
- Khachatourians, G. G. 1986. Production and use of biological pest control agents. *Trends in Biotechnology*, 4: 120-124.
- Kleespies, R. G. and Zimmermann, G. 1992. Production of blastospores by three strains of *Metarhizium anisopliae* (Metch.) Sorokin in submerged culture. *Biocontrol Science and Technology*, 2: 127-135.
- Nikdel, M., Sadaghian, B. and Dordaei, A. 2003. Evaluation of biology and natural enemies of brown tail moth *Euproctis chrysorroea* in Arasbaran forests. *Iranian Journal of Forest and Range Protection Research*, 1: 97-118.
- Püntener W., 1981 *Manual for field trials in plant protection*. Second edition. Agricultural Division, Ciba-Geigy Limited, Available on: Bakre, E., www.ehabsoft.com/ldpline/onlinecontrol.htm.
- Puzzari, K. C., Sarmahd, D. K. and Hazarika, L. k. 1997. Medium for mass production of *Beauveria bassiana* (Balsamo) Vuillemin. *Journal of Biological Control*, 11: 97-100.

- Safavi, S. A., Shah, F. A., Pakdel, A. K., Rasouljan, G. R., Bandan, A. R. and Butt, T. M. 2007. Effect of nutrition on growth and virulence of the entomopathogenic fungus *Beauveria bassiana*. FEMS Microbiology Letters, 270: 116-123.
- Schisler, D. A., Jackson, M. A. and Bothast, R. J. 1991. Influence of nutrition on conidiation of *Colletotrichum truncatum*, conidial germination and efficacy in inciting disease in *Sesbania exaltata*. Phytopathology, 81: 587-590.
- Shashi, S., Gupta, R. B. L. and Yadava, C. P. R. 1999. Mass multiplication and formulation of entomopathogenic fungi and their efficacy against white grubs. Journal of Mycology and Plant Pathology, 29: 299-305.
- Smith, R. J. and Gula, E. A. 1981. Nutritional requirements for conidial germination and hyphal growth of *Beauveria bassiana*. Journal of Invertebrate Pathology, 37: 222-230.
- St. Leger, R. J., Butt, T. M. and Robert D. J. 1989. Production in vitro of appressoria by the entomopathogenic fungus *Metarhizium anisopliae*. Experimental Mycology, 13: 247-288.
- Verhaar, M. A. and Hijwegen, T. 1993. Efficient production of phialoconidia of *Verticillium lecanii* for biocontrol of cucumber powdery mildew, *Sphaerotheca fuliginea*. European Journal of Plant Pathology, 99: 101-103.

مقایسه چند محیط غذایی طبیعی برای تولید بلاستوسپور قارچ *Beauveria bassiana* و زهر آگینی
 علیه پروانه دم‌قهوه‌ای بلوط، (*Euproctis chrysorrhoea* (Lep.: Lymantriidae)

پریسا بنامولایی^۱، رضا طلایی حسنلویی^۱ و حسن عسکری^۲

۱- گروه گیاهپزشکی، پردیس کشاورزی و منابع طبیعی دانشگاه تهران، کرج، ایران.

۲- مؤسسه تحقیقات گیاهپزشکی کشور، تهران، ایران.

* پست الکترونیکی نویسنده مسئول مکاتبه: rtalaei@ut.ac.ir

دریافت: ۲۹ آذر ۱۳۹۳؛ پذیرش: ۲۲ اسفند ۱۳۹۳

چکیده: اثر سه سطح از محیط‌های غذایی شامل ملاس چغندر قند، آب پنیر، عصاره سبوس گندم، عصاره سبوس برنج و محیط مایع سابورود دکستروز برای تولید بلاستوسپورهای دو جدایه از قارچ *Beauveria bassiana sensu lato* در فواصل زمانی ۲۴ ساعته به مدت ۷ روز مورد بررسی قرار گرفت. بسته به جدایه، بیشینه تولید بلاستوسپور در روز هفتم در عصاره سبوس برنج ۱۲ درصد و آب پنیر ۲۰ درصد بود. برای هر دو جدایه، کم‌ترین مقدار بلاستوسپور تولیدی از آب پنیر ۴ درصد به دست آمد. زهر آگینی بلاستوسپورهای تولید شده در محیط‌های مایع حاوی ملاس چغندر قند، آب پنیر، عصاره سبوس گندم و سابورود دکستروز (به عنوان شاهد) روی لارو سن سوم پروانه دم‌قهوه‌ای، *chrysorrhoea* نشان داد که در مورد هر دو جدایه بین این محیط‌های غذایی تفاوت معنی‌داری وجود ندارد. با توجه به کمیت و کیفیت بلاستوسپورها از نظر زهر آگینی و دسترسی محلی، آب پنیر به عنوان محیط مناسب برای تولید انبوه بلاستوسپورهای قارچ *B. bassiana* معرفی می‌شود.

واژگان کلیدی: *Beauveria bassiana*، پروانه دم‌قهوه‌ای بلوط، بلاستوسپور، زهر آگینی، محیط غذایی طبیعی