J. Crop Prot. 2015, 4 (4): 463-475_



Research Article

Antifeedant effect of gamma radiation and *Perovskia atriplicifolia* essential oil combination against *Tribolium castaneum* (Coleoptera: Tenebrionidae)

Mehrdad Ahmadi¹*, Saeid Moharramipour² and Adly Mohamed Abd alla³

1. Nuclear Agriculture Research School, Nuclear Science and Technology Research Institute, Karaj, Iran.

2. Department of Entomology, College of Agriculture, Tarbiat Modares University, Tehran, Iran

3. Department of Pests and Plant Protection, National Research Centre, Dokki, Giza, Egypt.

Abstract: Control of stored-product insect by gamma radiation could be enhanced by other possible techniques such as essential oils as effective alternatives to chemical insecticides. In this study, the efficiency of gamma radiation combined with Perovskia atriplicifolia (Benth) was verified to assess their enhanced antifeedant effect against the Tribolium castaneum (Herbst). Flour disc bioassay was employed to assess the nutritional indices, such as relative growth rate (RGR), relative consumption rate (RCR), efficiency of conversion ingested food (ECI) and feeding deterrence index (FDI). The results showed that irradiation, essential oil and combination of both of them significantly reduced RGR, RCR and ECI in treated larvae and adults. As irradiation at 100 Gy reduced RGR of the larvae and adults from 0.0884 and 0.0366 to 0.0596 and 0.0332 mg/mg/day respectively, when combined with doses of P. atriplicifolia essential oil, the reduction reached to 0.051 - 0.388 and 0.01- 0.224 mg/mg/day respectively. The reduction rates in larvae were even greater when combination of irradiation and EO treatments were used. The results showed a significant increase in the feeding deterrence due to irradiation, essential oil and specially their combination. Therefore, findings led to a conclusion that irradiation can enhance antifeedant activity of essential oils.

Keyword: *Perovskia atriplicifolia*, relative growth rate, relative consumption rate, efficiency of conversion of ingested food, feeding deterrence index

Introduction

The storage pest *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae) is a major economic pest in flour mills and storage facilities because of its cosmopolitan distribution and affinity for stored grain, especially in tropical and sub-tropical climates (Mills and White, 1994). Under optimum

conditions of 25-35°C and 60-80% R.H., the adults and larvae cause serious damage to infested grains. The insects can be found on the surface or at the depth of grain stores (Bagheri Zenouz, 1997). Members of genus *Tribolium* are known to produce toxic quinones which contaminate flour and flour products (Gorham, 1987). Damage is inflicted by both larvae and adults with a reduction in quality and quantity of stored products (Bagheri Zenouz, 1997). To control this pest, fumigants and other chemical insecticides such as methyl bromide and phosphine are widely used. The extensive use of these treatments leads to undesirable residues

Handling Editor: Fariba Mehrkhou

^{*} **Corresponding author**, e-mail: mahmadi@nrcam.org Received: 28 September 2014, Accepted: 05 May 2015 Published online: 28 June 2015

(Mensah et al., 1979) in addition to development of resistance in certain insect species (Rossi et al., 2010). However, methyl bromide was classified as an ozone depleter, and due to environmentally unfavorable effects, its use has recently been banned in many countries (Carpenter et al., 2000). Due to these restrictions new alternative strategies should be investigated. Irradiation and essential oils have become approved and feasible alternatives to the conventional method for controlling stored product pests; because of residue free chemical fumigation. advantages over Irradiation can also extend the shelf life of various products and maintain the quality of the product over a longer period of time (Pszczola, 1997) however depending on the dose of ionizing energy applied; high doses of radiation may affect the quality of foods (Urbain, 1986; Bothaina et al., 2002).

Insecticidal, repellency, anti-nutritional and genotoxic properties are very important in choosing the method to control stored product pests. Insecticidal effect of gamma radiation and also essential oil from *Perovskia atriplicifolia* (Benth) on the red flour beetle *T. castaneum* has been previously described (Brower and Tilton, 1973a, b; Brower *et al.*, 1973; Misra and Paravathy, 1998; Ayvaz *et al.*, 2002; Tuncbilek *et al.*, 2003; Ahmadi *et al.*, 2013). Moreover, genotoxic effect of gamma radiation on genital cells of *T. castaneum* has been reported by Ahmadi *et al.* (2011).

On the other hand, the increasing efforts to develop environmentally friendly pest control methods have attracted the attention of many researchers towards the use of antifeedants (Jermy, 1990). Insect gonads and midgut contain mitotically active tissues, and irradiated insects are often sterile and stop feeding soon after treatment (Follett and Griffin, 2006). Doses of 0.2 to 1.0 kGy of gamma radiation are not immediately lethal, and irradiated insects can survive for several weeks, but they feed less and are usually infertile (Katz and Weaver, 2003). According to Brower and Tilton (1973b), the amount of damage inflicted by live but sterile *Sitophilus oryzae* (L.) and *Rhyzopertha dominica* (F.) adults treated with gamma radiation in bulk grain would probably not be a serious problem because of the great reduction in feeding caused by radiation effects. A wide variety of plants may supply new sources of natural pesticides, antifungals, antifeedants, and repellents (Grainge and Ahmed, 1988; Arnason *et al.*, 1989; Ananthakrishnan, 1992; Cetinsoy *et al.*, 1998; Abbasipour *et al.*, 2011). In search for alternatives to conventional pesticides, the essential oil of *Perovskia* spp has been widely investigated, and showed toxic, repellent and antifeedant effects against stored product pest, red flour beetle (Arabi *et al.*, 2008).

In addition, the application of irradiation has been recommended not only as a possible alternative but also as a supplement for other control methods (Cornwell, 1964). The combination of gamma radiation with other treatments like microwave, infra-red radiation, insecticides and essential oils has also been evaluated by various researchers (Mehta et al., 2004; Tilton et al., 1972; Cogburn and Spiers, 1972; Cogburn et al., 1971; Ahmadi et al., 2013). Ahmadi and Moharramipour, (2011) have reported that combination of gamma radiation and Rosmarinus officinalis L. essential oil may result in synergistic interactions that reduce the feeding of T. castaneum larvae. Therefore, it seems likely that low doses of gamma radiation combined with essential oil treatment would have antiffedant effect in T. castaneum population. This work was undertaken to study the antifeedant action of gamma radiation and P. atriplicifolia essential oil combination as a safe control method against different life stages of T. castaneum.

Materials and Methods

Insect

Tribolium castaneum was reared on wheat flour mixed with yeast (10:1 w/w). The cultures were maintained in the dark in the growth chamber set at 27 ± 1 °C and $50 \pm 5\%$ R.H. All experiments were conducted under these conditions.

Ahmadi et al. _____

Plant materials

Aerial parts of *P. atriplicifolia* were collected in full bloom stage in Tehran in April 2011. The collected plant materials were dried on laboratory benches at room temperature (23-24 °C) for 5 days (or until crisp). The dried materials were stored at -24 °C until processed by hydrodistillation to extract its essential oil.

Extraction of essential oils

Essential oils were extracted from the plant samples according to the method previously described by Negahban and Moharramipour (2007). In brief, using a Clevenger-type apparatus, in which the plant materials were hydrodistilled. Portions of the material (40 g of an air-dried sample, 1: 10 plant / water ratio) were distilled for 4 h. Anhydrous sodium sulfate was used to remove water after the extraction. Extracted oil was stored in refrigerator at 4 °C.

Irradiation

Irradiation of the tested insects was administered using 60 cobalt gamma sources at the Nuclear Science and Technology Research Institute, Nuclear Agriculture Research School, Karaj, at a dose rate of 0.4 Gy/sec. The required dose rate was obtained by exposing treated insects for variable times. To determine lethal doses of gamma radiation, healthy and active adult (1-3 days old) and larvae (5-10 days old) were irradiated by doses of 0, 100, 200, 300, 400 and 500 Gy with five replicates for each dose. All stages were then maintained in the dark in growth chamber.

Combination of gamma radiation and essential oil treatments

Experiments were designed to combine gamma irradiation and essential oil treatments to investigate their combined effects on adults and larvae of *T. castaneum* feeding. Insects were exposed to five doses (100, 200, 300, 400 and 500 Gy) of radiation and flour discs treated with three concentrations of the diluted *P. atriplicifolia* oils with acetone (5.27, 7.56 and

13.11 μ l/l air). Each experiment was replicated five times with twenty insects per replicate.

Antifeedant effect of combined irradiation and essential oil treatment

Flour discs were prepared according to the method of Xie et al. (1996), in brief 10 g of flour was mixed with 50 ml of distilled water. Using a micropipette, 200 ml of the prepared suspension was poured on a nylon sheet to convert the suspension to spherical discs. The discs were placed in normal room conditions for 4 h. and then transferred to sterile Petri dishes with the help of fine forceps. The flour discs were stored for 12 h. inside the hood to dry completely. In each container 2 discs covered by different doses of oil were placed (acetone used as control), and 20 irradiated insect (larvae or adults) starved for 48 h before exposure. Five replicates of each treatment were used. The weight of the flour discs and insects were accurately measured and recorded at the beginning and at 3 days post start of the experiment.

Nutritional indices were calculated according to the formula of Manuwoto and Scriber, (1992); Farrar *et al.* (1989) and Isman *et al.* (1990), (Cited by Huang *et al.*, 1997) with modifications as follows:

Relative Growth Rate (RGR) = $(A - B) / (B \times DAY)$ Where A weight of live insects on the third day (mg) / number of insects on the third day

B = initial weight of insects (mg)/initial number of insects

Relative Consumption Rate (RCR) = $D/(B \times day)$, Where D = biomass ingested (mg)/number of live insects on the third day

Efficiency of Conversion of Ingested Food (ECI%) = RGR/RCR \times 100

Feeding Deterrence Index (FDI%) = $(C-T)/C \times 100$, Where C is consumption of control discs, and T is consumption of treated discs.

Statistical analysis

The dosage mortality response was determined by probit analysis (Finney, 1971). Before the statistical analysis, ECI and FDI were normalized. ANOVA and Tukey's HSD test were employed using the SPSS software to compare the means.

Results

Irradiation

The RGR, RCR, ECI and FDI values of irradiated adults and larvae of *T. castaneum* fed on flour discs for 72 h are given in Fig. 1. The results of the antifeedant test indicate that the rates of RGR and RCR indices in irradiated larvae and adults were significantly lower than those of non-irradiated ones (P < 0.05).

At a dose of 100 Gy, RGR of adults was reduced from 0.0366 to 0.0332 mg for each mg of insect weight comparing to the control (9.2% reduction rate), whereas administering the same dose for the larvae caused, 32.57% reduction rate (reduction from 0.0884 to 0.0596 mg mg-1, daily). There were 13.11, 12.02, 18.30 and 21.03% reduction rates in the RGR in adults and 32.35, 37.44, 40.72 and 44.34% reduction in larvae exposed to 200, 300, 400 and 500 Gy of gamma ray respectively. There was no significant difference in the RGR reduction rate in adults between the exposure to 200 and 300 Gy doses. Also the exposure of larvae to 100 and 200 Gy doses did not show significant difference in the RGR reduction rate (Fig. 1). The results show that 2.32, 2.26, 0.97, 3.46 and 5.01% reductions in the RCR of the adults were caused by the exposure to 100, 200, 300, 400 and 500 Gy of gamma ray respectively while these reduction rates in larvae were 38.62, 38.97, 46.90, 48.03 and 53.68% respectively. In addition, it was shown that comparative effects of 100 and 200 Gy doses on the RCR of the adults and larvae were not significant.

Moreover, with the exception of 500 Gy dose of gamma ray treatment which caused 1.85% reduction in the ECI of the exposed larvae, the other doses increased the ECI rate. On the other hand in adults, the ECI significantly (P<0.05) decreased by

increasing irradiation doses (7.14, 5.79, 18.73, 15.34 and 17.92% reduction rates by using of 100, 200, 300, 400 and 500 Gy doses, respectively). The data shows that gamma radiation has significant effect on the FDI of treated larvae and adults. The FDI of the adults and larvae increased from 0 to 2.39 and 60.9% respectively after exposure to 100 Gy of gamma ray. Generally the FDI amount increased by exposure to gamma ray, and the effect value among the larvae was much greater than that for the adults (Fig. 1).

Essential oil

The RGR significantly (P < 0.05) declined at all *P. atriplicifolia* essential oil concentrations used in this study. The treatment with the essential oil at doses of 5.27, 7.65 and 13.11 µl/l caused 41, 45.7 and 54.2% in larvae and 33.9, 34.4 and 67.7% reduction in adults respectively. This result indicates that the reduction rate gradually increased with increasing oil concentration. The results also indicate that the RCR in treated larvae and adults decreased significantly (P < 0.05) with increasing oil concentration. The reduction rate of RCR was higher in larvae than adults. In contrast, the same doses, significantly reduced the ECI in adults (30.7, 32.5 and 53.9%) more than in larvae (16.3, 16.9 and 22.49%) respectively. Similar to the RGR trend, the reduction rate of the ECI increased with increasing the oil concentrations (Fig. 2). The FDI values of the treated larvae and adults with essential oil increased with increasing the concentration. At doses of 5.27, 7.65 and 13.11 µl/l the FDI increased from 0 to 35.05, 45.12 and 49.55 in larvae and to 21.25, 28.22 and 35.45 in adults respectively. The data indicate that the increasing rate of the FDI is higher in treated larvae than adults (Fig. 2).



Figure 1 Effects of gamma radiation on relative growth rate (RGR), relative consumption rate (RCR), efficiency of conversion ingested food (ECI) and feeding deterrence index (FDI) of *Tribolium castaneum* adults and larvae. Means followed by the same letters in each line are not significantly different (Tukey's HSD test at 5% level). Vertical bars indicate standard error of mean.



Figure 2 Effects of *Perovskia atriplicifolia* essential oil on relative growth rate (RGR), relative consumption rate (RCR), efficiency of conversion ingested food (ECI) and feeding deterrence index (FDI) of *Tribolium castaneum* adults and larvae. Means followed by the same letters in each line are not significantly different (Tukey's HSD test at 5% level). Vertical bars indicate standard error of mean.

Combination of radiation and essential oil

The antifeedant effect of gamma radiation in combination with P. atriplicifolia on T. castaneum increased markedly with an increase of the doses. As the exposure to 100 Gy reduced RGR of the larvae and adults from 0.0884 and 0.0366 to 0.0596 and 0.0332 respectively, while when combined with doses of P. atriplicifolia essential oil the reduction rate reached 0.051-0.388 and 0.224-0.01 respectively. These combinations especially at high doses were found to be appropriate for the purpose of the experiments (Tables 1 and 2). In all irradiation doses (100, 200, 300, 400 and 500 Gy), significant reduction in the RGR of the larvae and adults occurred when they combined with 13.11 μ l/l of oil (34.2, 45.6, 47.9, 57.1 and 48.1% in the larvae and 69.8, 66.9, 67.4, 71.57 and 74.1% in the adults respectively). The results pointed out those adults had more reduction rate in their RGR compared with larvae. In the RCR of the treated larvae, low and high reductions were observed in combination of 200 Gy-5.27 $\mu l/l$ and 200 Gy-13.11 µl/l respectively. Also in the treated adults, low and high reductions were observed in combination of 100 Gy-5.27 μ l/l and 500 Gy-13.11 μ l/l respectively. Combined doses of the gamma radiation-P. atriplicifolia essential oil which induced low and high reduction of the ECI were 100 Gy-5.27 μ l/l and 400 Gy-13.11 μ l/l for the larvae and 100 Gy-5.27 µl/l and 200 Gy-13.11 μ l/l for the adults, respectively (Tables 1 and 2). The results showed that the combined treatments had greater effect on the ECI of the adults than the larvae. On the other hand, combined dose of 400 Gy-13.11 µl/l of gamma radiation and essential oil increased the FDI of larvae from 57.3 to 93.22. Also combined doses of 500 Gy-13.11 µl/l increased the FDI of the adults 4.01to 69.5. These results from demonstrate the high increasing of the FDI among all of the combination treatments

(Tables 1 and 2). In all experiments, the antifeedant effects of gamma radiation or essential oil separately were lower than the antifeedant rate of combination of these two treatments.

Discussion

Irradiation

Based on the results presented in this study, it was determined that gamma radiation has significant effects on the nutritional indices of T. castaneum at different doses and it was more effective on the larvae compared with the adults. The RGR and RCR indices were significantly reduced with irradiation; furthermore, the reduction rate was greater among larvae than adults. These differences could be due to high feeding rate of the treated larvae and their high sensitivity to gamma radiation compared with the treated adults. No significant differences were observed between the effect of 100 and 200 Gy doses on the RGR in larvae or 200 and 300 Gy in adults which indicates that the growth response of larvae and adults against different doses of radiation are different. In addition, the effect of radiation on the adult RCR was not significant which could be explained by the low feeding of irradiated and non-irradiated adults in comparison with the larvae. The effect of gamma irradiation on the ECI in larvae and adults was different to some extent. Except 500 Gy, in other doses the ECI percentage in the larvae was increased, whereas in the adults the use of gamma radiation caused significant decrease in amount of the ECI. It can be inferred that irradiation causes high reduction in the larval feeding, then the larvae which have a small part of food attempt to increase the ECI.

Essential oil (µl / Disk)	Gamma (Gy)	Mean \pm SE ¹				
		RGR (mg/mg/day)	RCR (mg/mg/day)	ECI (%)	FDI (%)	
0	0	0.0884 ± 0.0029 a	0.4603 ± 0.0321 a	19.38 ± 1.15 bc	0 j	
5.27	0	$0.0521\pm0.005~cd$	$0.3224 \pm 0.0122 \ b$	$16.21 \pm 1.05 \text{ d}$	35.05 ± 3.25 i	
7.65	0	$0.0480 \pm 0.0041 \ def$	$0.2755 \pm 0.0196 \ bcd$	$16.1\pm1.08\ d$	$45.12\pm2.2\ h$	
13.11	0	0.0405 ± 0.0022 hi	$0.2105 \pm 0.0286 \; e$	$15.05\pm1.55~def$	$49.55 \pm 2.45 \ h$	
0	100	$0.0596 \pm 0.002 \; b$	$0.2825 \pm 0.0232 \ bc$	$21.22\pm1.15\ ab$	$60.90\pm2.42~g$	
5.27	100	$0.0510 \pm 0.0025 \ cd$	$0.2322\pm0.011\text{ bcde}$	$16.05 \pm 0.95 \ de$	$78.30 \pm 2.85 \; f$	
7.65	100	0.0420 ± 0.0013 ghi	0.1825 ± 0.0122 cde	$15.58\pm1.28~\text{de}$	$80.66 \pm 3.22 \text{ ef}$	
13.11	100	$0.0388 \pm 0.0028 \; i$	$0.1524 \pm 0.0145 \ cde$	$14.25\pm1.05~defg$	$88.23\pm3.25\ bcd$	
0	200	$0.0598 \pm 0.0036 \; b$	$0.2809 \pm 0.0266 \ bc$	21.83 ± 1.92 a	$62.01 \pm 3.71 \text{ g}$	
5.27	200	$0.0485 \pm 0.0025 \ def$	0.231 ± 0.0215 bcde	15.55 ± 0.75 de	$79.45 \pm 4.22 \ ef$	
7.65	200	$0.0455 \pm 0.0025 \ efg$	$0.1752 \pm 0.0125 \ cde$	$14.65\pm0.85~def$	$82.10 \pm 3.56 \text{ def}$	
13.11	200	$0.0325 \pm 0.0015 \; j$	$0.1422 \pm 0.0225 \ e$	$14.2\pm0.95~defg$	$85.15\pm2.85\ cde$	
0	300	$0.0553 \pm 0.0031 \ bc$	$0.2444 \pm 0.0236 \ bcde$	21.56 ± 1.68 a	$61.65 \pm 2.24 \text{ g}$	
5.27	300	$0.0440 \pm 0.0028 \ fgh$	$0.1850 \pm 0.0189 \; cde$	$14.55\pm0.85~def$	$85.12\pm3.21~\text{cde}$	
7.65	300	$0.0325 \pm 0.0026 \ j$	$0.1820 \pm 0.0188 \ cde$	$14.25\pm1.57~defg$	$89.36 \pm 4.21 \ abc$	
13.11	300	$0.0288 \pm 0.0024 \; jk$	$0.1385 \pm 0.0222 \ e$	$13.85\pm0.89~efg$	$92.25\pm3.05~ab$	
0	400	$0.0524 \pm 0.0023 \ cd$	0.2392 ± 0.0123 bcde	21.93 ± 0.43 a	$57.30\pm0.89~g$	
5.27	400	$0.0445\pm0.0025~efgh$	$0.1888 \pm 0.0135 \; cde$	$14.21\pm0.65~defg$	$89.63 \pm 3.55 \text{ abc}$	
7.65	400	$0.0332 \pm 0.0011 \; j$	$0.1756 \pm 0.0111 \ cde$	$14.05\pm0.85~defg$	$89.95 \pm 4.55 \ abc$	
13.11	400	$0.0225 \pm 0.0018 l$	$0.1355 \pm 0.0215 \ e$	$13.1\pm0.58~fg$	93.22 ± 4.33 ab	
0	500	$0.0492 \pm 0.0011 \text{ de}$	$0.2132\pm0.0114\ bcde$	$19.02\pm1.05\ c$	$60.42 \pm 3.09 \text{ g}$	
5.27	500	$0.0325 \pm 0.0022 \; j$	$0.1765 \pm 0.0205 \ cde$	$13.25\pm1.55~fg$	$88.50\pm4.33\ bc$	
7.65	500	$0.0333 \pm 0.0036 j$	0.1655 ± 0.0233 de	$12.25\pm0.78~g$	91.22 ± 5.24 abc	
13.11	500	$0.0255 \pm 0.0036 \; kl$	$0.1345 \pm 0.0195 \ e$	$12.10 \pm 1.25 \text{ g}$	95.25 ± 5.05 a	

Table 1 Combined antifeedant effect of gamma radiation and Perovskia atriplicifolia essential oil on Triboilum castaneum larvae.

¹ Means followed by the same letters within a column are not significantly different (Tukey's HSD test at 5% level).

Essential oil	Gamma	Mean \pm SE ¹				
$(\mu l / D_{1SK})$	(Gy)	RGR	RCR	ECI (%)	FDI (%)	
0	0	(mg/mg/day)	(mg/mg/day)	16.02 - 1.42	0	
0	0	0.0366 ± 0.0019 a	0.2253 ± 0.01 a	16.23 ± 1.43 a	0 m	
5.27	0	$0.0242 \pm 0.004 \; d$	$0.1755 \pm 0.0117 \; b$	$11.25\pm1.05\ cd$	$21.25\pm2.33\ k$	
7.65	0	$0.0240 \pm 0.0025 \ d$	$0.1699 \pm 0.0129 \text{ bc}$	$10.95 \pm 1.35 \text{ d}$	$28.22\pm1.05\ j$	
13.11	0	$0.0118 \pm 0.0011 \ g$	$0.1322\pm0.01~def$	$7.48\pm0.88~efg$	$35.45 \pm 2.25 \ i$	
0	100	$0.0332 \pm 0.0016 \; b$	$0.2199 \pm 0.027 \ a$	$15.07\pm1.27~ab$	$2.39\pm0.53\ lm$	
5.27	100	$0.0224 \pm 0.0015 \text{ de}$	$0.1725 \pm 0.01 \text{ bc}$	$9.25\pm1.55\;de$	$39.45\pm1.25\ h$	
7.65	100	$0.0219 \pm 0.0016 \; de$	$0.1555 \pm 0.019 \; bcd$	$8.25\pm1.58~ef$	$41.25\pm1.66\ h$	
13.11	100	$0.010\pm0.0014~gh$	$0.1185\pm0.012~efgh$	$5.55\pm0.63~\text{ghi}$	$46.35\pm2.33~g$	
0	200	$0.0318 \pm 0.0025 \ bc$	$0.2202 \pm 0.019 \text{ a}$	$15.29\pm2.46\ ab$	$2.11\pm0.13\ lm$	
5.27	200	$0.0205 \pm 0.0018 \; e$	$0.1666 \pm 0.018 \ bc$	$8.65\pm0.58\;e$	$41.25\pm1.59\ h$	
7.65	200	$0.0195 \pm 0.001 \; e$	$0.1325 \pm 0.018 \; def$	$5.33\pm0.65~ghi$	$45.85 \pm 2.25 \ g$	
13.11	200	$0.0105 \pm 0.0012 \ gh$	$0.1300 \pm 0.011 \; def$	4.26 ± 0.75 hi	$51.25 \pm 2.33 \; f$	
0	300	$0.0322 \pm 0.0021 \ bc$	$0.2231 \pm 0.015 \ a$	$13.19\pm1.25\ bc$	$3.03\pm0.89\ lm$	
5.27	300	$0.0215 \pm 0.0019 \; de$	$0.1450\pm0.018~\text{cde}$	$6.33\pm0.87~fgh$	$48.88\pm1.25~fg$	
7.65	300	$0.0156 \pm 0.0018 \; f$	$0.1255\pm0.016~efg$	5.23 ± 0.85 hi	$55.35\pm3.56~e$	
13.11	300	$0.0105 \pm 0.0015 \ gh$	$0.1125\pm0.015~fghi$	5.20 ± 0.65 hi	$59.25\pm2.45~d$	
0	400	$0.0299 \pm 0.0014 \ bc$	$0.2175 \pm 0.014 \; a$	$13.74\pm1.21\ b$	$3.32\pm0.61\ lm$	
5.27	400	$0.0115 \pm 0.0015 \ g$	$0.1133\pm0.015~fghi$	$6.25\pm0.94~fgh$	$52.23 \pm 1.22 \text{ ef}$	
7.65	400	$0.0090 \pm 0.0008 \ gh$	$0.11\pm0.014~fghi$	$4.29\pm0.85~\text{hi}$	$61.25\pm2.22~cd$	
13.11	400	$0.0085 \pm 0.0021 \ gh$	0.0955 ± 0.018 hi	$4.12\pm0.83\ hi$	$63.44\pm3.85\ bc$	
0	500	$0.0289 \pm 0.0025 \ c$	0.2140 ± 0.011 a	$13.32\pm2.51~b$	$4.01\pm0.56l$	
5.27	500	$0.0119 \pm 0.0013 \; g$	$0.1055\pm0.016~fghi$	$4.25\pm0.45~\text{hi}$	$61.25 \pm 2.33 \text{ cd}$	
7.65	500	$0.0095 \pm 0.0008 \; gh$	0.0985 ± 0.017 ghi	$3.82\pm0.55\ i$	$66.33 \pm 4.25 \ ab$	
13.11	500	$0.0075 \pm 0.0017 \; h$	$0.08555 \pm 0.018 \ i$	$3.25\pm0.25\ i$	69.5 ± 3.05 a	

 Table 2 Combined antifeedant effect of gamma radiation and Perovskia atriplicifolia essential oil on Triboilum castaneum adults.

¹ Means followed by the same letters within a column are not significantly different (Tukey's HSD test at 5% level).

The results also showed that gamma radiation had feeding deterrence effects on either adults or larvae of *T. castaneum* and the effect on FDI was higher in larvae than adults; this fact illustrates that with the lowest dose of gamma radiation, sharp reduction on feeding of the larvae, (the most active stage) will be

produced. Reduction in the feeding by storedproduct beetles has been observed after radiation treatments, with a corresponding decrease in feeding damage (Cornwell, 1964; Watters and MacQueen, 1967; Brower and Tilton, 1973a, b). Johnson and Vail (1988, 1989) reported that feeding and damage caused

Ahmadi et al. _____

by navel orange worm larvae could be reduced by as much as 78% after 300 Gy treatments; this treatment in our study reduced the feeding of larval stages of *T. castaneum* about 61.65%.

Antifeedant effects of gamma radiation can likely be due to insect's physiology and sensitivity of cells to irradiation. Sensitivity of any tissue to radiation is dependent on the degree of cellular development activity and differentiation (Bergonie and Tribondeau, 1959). Cells of the midgut epithelium of feeding insects are frequently replaced by undifferentiated regenerative cells. Radiations usually destroy these regenerative cells resulting in histolysis of the midgut epithelium (Ashraf *et al.*, 1971), which can affect the feeding amount of irradiated insects.

Essential oil

This study demonstrates that the essential oil of P. atriplicifolia had toxic and significant effect on the nutritional indices of the larvae and adults of T. castaneum at different doses. P. atriplicifolia essential oil can significantly reduce the RGR and RCR of the adults and larvae, also the treated larvae showed more sensitivity to the oil effect compared with the treated adults. There are several reports on antifeedent activity of essential oils. Huang and Ho (1998) reported the antifeedant action of the Cinnamomun aromaticum Nees, on the red flour beetle adults and larvae. Also Abbasipour et al. (2011) stated that Datura stramonium L. (Solanaceae) extract had significant effects on the nutritional indices and mortality of T. castaneum adults at different concentrations. They showed that this oil significantly reduced the growth rate and food consumption and utilization of T. castaneum adults at higher concentrations. Negahban and Moharramipour (2007) also verified that Artemisia scoparia Waldst. & Kit and Artemisia sieberi Besser extracts had antifeedant property on T. castaneum adults. These studies confirm the validity of our experiments however, P. atriplicifolia essential oil had a more antiffedant effect on T. castaneum than did C. aromaticum and D. stramonium but, it's inhibition of feeding was less than that of *A. sieberi* and *A. scoparia*. Although contact toxicity of *P. atriplicifolia* was verified (Ahmadi *et al.*, 2013) and its EO was shown to affect the feeding of the treated larvae and treated adult of *T. castaneum*, however the applied doses were high. Therefore it may not be feasible to apply high just doses of this essential oil for insect control purposes.

Combination of radiation and essential oil

Similar to other food processing methods, irradiation can make certain changes in the commodity that can modify the composition and nutritive values of food (Wiendl, 1984). The use of low irradiation doses combined with other treatments is a way to minimize the negative effects of irradiation on food quality (Dionisio et al., 2009). The study of Ahmadi et al. (2013) indicated that P. atriplicifolia essential oil could be combined with gamma radiation to enhance the mortality effect of irradiation. They showed that the effect of the essential oil and gamma radiation can be increased and the doses can be reduced. The results demonstrated when a combination of P. atriplicifolia and gamma radiation was employed, insect mortality increased not only by combined toxicity of both treatment but also by their adverse effects on T. castaneum feeding behavior.

The antifeedant effects of some combined treatments to control the insects have been previously reported. Huang et al. (2002) specified that the toxicity of the deltamethrin could be enhanced to kill insects effectively. When a mixture of deltamethrin and eugenol or one of its analogues is applied, insects die due to the combined toxic and antifeedant effects. Moreover, the study of Ahmadi and Moharramipour (2011) indicated that the combination of gamma radiation and R. officinalis essential oil could reduce the feeding of T. castaneum larvae. They showed that relative growth rate had significantly decreased (P < 0.05) by combining the gamma radiation and R. officinalis and that the severity of the reduction rate increased by increasing doses. Also relative food consumption rate decreased

when gamma radiation and R. officinalis were combined and its reduction severity had convert relative with increasing of doses. This study demonstrates the validity of our results. Our data suggest that a combination of treatments with essential oils and irradiation can play a major role in management of stored-product insects. However, the vapor pressure of essential oils is too low to enable them to penetrate into commodities to kill pests, and the treatment can be employed only on a small scale. In order to make practical use of essential oils, these compounds should be formulated. The present study shows that P. atriplicifolia essential oil when applied with gamma radiation can control T. castaneum by killing its two active and harmful life stages by means of fumigant and antifeedant actions.

Acknowledgments

We thank Mohammad Babaei and Hadi Fatolahi, from Nuclear Agriculture Research School, Nuclear Science and Technology Research Institute for their technical assistance.

References

- Abbasipour, H., Mahmoudvand, M., Rastegar, F. and Hosseinpour, M. H. 2011. Bioactivities of jimsonweed extract, *Datura stramonium* L. (Solanaceae), against *Tribolium castaneum* (Coleoptera: Tenebrionidae). Turkish Journal of Agriculture and Forestry, 35: 623-629.
- Ahmadi, M. and Moharramipour, S. 2011.
 Antifeedant effects of gamma radiation and *Rosmarinus officinalis* to larval stage of *Tribolium castaneum* (Col: Tenebrionidae).
 International Journal of Artificial Organs, 34: 667-667.
- Ahmadi, M., Abd-Alla, A. M. M. and Moharramipour, S. 2013. Combination of gamma radiation and essential oils from medicinal plants in managing *Tribolium castaneum* contamination of stored products. Applied Radiation and Isotopes, 78: 16-20.
- Ahmadi, M., Moharramipour, S., Mozdarani, H. and Babaii, M. 2011. Induction of

micronuclei in ovaries of *Tribolium castaneum* exposed to gamma radiation and Rosmarinus officinalis essential oil. IOBC Bulletin, 69: 196.

- Ananthakrishnan, T. N. 1992. Dimensions of Insect-Plant Interactions. Oxford and IBH Publishing, New Delhi.
- Arabi, F., Moharramipour, S. and Sefidkon, F. 2008. Chemical composition and insecticidal activity of essential oil from *Perovskia abrotanoides* (Lamiaceae) against *Sitophilus oryzae* (Coleoptera: Curculionidae) and *Tribolium castaneum* (Coleoptera: Tenebrionidae). International Journal of Tropical Insect Science, 28 (3): 144-150.
- Arnason, J. T., Philogene, B. J. R. and Morand, P. 1989. Insecticides of Plant Origin. American Chemical Society, Washington, D. C.
- Ashraf, M., Brower, J. H. and Tilton, E. W. 1971. Effects of gamma radiation on the larval midgut of the Indian-meal moth, *Plodia interpunctella* (Lepidoptera: Phycitidae). Radiation Research, 45: 349-354.
- Ayvaz, A., Ozturk, F., Yaray, K. and Karahacio, E. 2002. Effect of the gamma radiations and malathion on confused flour beetle, *Tribolium confusum* J. du Val. Pakistan Journal of Biological Sciences, 5: 560-562.
- Bagheri-zenouz, E. 1997. Storage Pests and Their Control, Vol. 1. Sepehr Press, 309 pp.
- Bergonie, J. and Tribondeau, L. 1959. Interpretation of some results of radiotherapy and an attempt at determining a logical technique of treatment. Radiation Research, 11: 587-588.
- Bothaina, M. Y., Asker, A. A., El-Samahy, S. K. and Swailam, H. M. 2002. Combined effect of steaming and gamma irradiation on the quality of mango pulp stored at refrigerated temperature. Food Research International, 35: 1-13.
- Brower, J. H. and Tilton, E. W. 1973a. Comparative gamma radiation sensitivity of *Tribolium madens* (Charpentier) and *T. castaneum* (Herbst). Journal of Stored Products Research, 9: 93-100.
- Brower, J. H. and Tilton, E. W. 1973b. Weight loss of wheat infested with gamma-radiated

Ahmadi et al. ___

Sitophilus oryzae (L.) and Rhyzopertha dominica (F.). Journal of Stored Products Research, 9: 37-41.

- Brower, J. H., Hossain, M. M. and Tilton, E. W. 1973. Radiation senility of successively irradiated generations of *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae), and *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae). Journal of Stored Products Research, 9: 43-49.
- Carpenter, J., Gianessi, L. and Lynch, L. 2000. The Economic Impact of the Scheduled U.S. Phaseout of Methyl Bromide. NCFAP, 463 pp.
- Cetinsoy, S., Tamer, A. and Aydemir, M. 1998. Investigations on repellent and insecticidal effects of *Xanthium strumarium* L. on Colorado potato beetle *Leptinotrasa decemlineata* Say (Col: Chrysomelidae). Turkish Journal of Agriculture and Forestry, 22: 543-552.
- Cogburn, R. R. and Spiers, R. D. 1972. Toxicity of malathion to gamma irradiated and nonirradiated adult red flour beetles. Journal of Economic Entomology, 65: 185-191.
- Cogburn, R. R., Brower, J. H. and Tilton, E. W. 1971. Combination of gamma and infrared radiation for control of Angoumois grain moth in wheat. Journal of Economic Entomology, 64: 923-925.
- Cornwell, P.B. 1964. Insect control, In: Jefferson S. (Ed.), Massive Radiation Techniques, Massachusetts Radiation Technology, George Newness, London. pp. 141-198.
- Dionisio, A. P., Gomes, R. T. and Oetterer, M. 2009. Ionizing Radiation Effects on Food Vitamins-A Review. Brazilian Archives of Biology and Technology, 52 (5): 1267-1278.
- Farrar, R. R., Barbour, J. D. and Kennedy, G. G. 1989. Quantifying food consumption and growth in insects. Annals of the Entomological Society of America, 82: 593-598.
- Finney, D. J. 1971. Probit Analysis, Third Edition, Cambridge University Press, London.
- Follett, P. A. and Griffin, R. L. 2006. Irradiation as a phytosanitary treatment for fresh horticultural commodities: Research and regulations. In: Sommers, C. H. and

Fan, X. (Eds), Food Irradiation Research and Technology, Oxford, Blackwell, 143-168.

- Gorham, J. R. 1987. Insect and Mite Pests in Food: An Illustrated Key. United States Department of Agriculture, Agricultural Handbook, No. 655.
- Grainge, M. and Ahmed, S. 1988. Handbook of Plants with Pest-Control Properties. Wiley-Interscience, New York.
- Huang, Y. and Ho, S. H. 1998. Toxicity and antifeedant activities of cinnamaldehyde against the grain storage insects, *Tribolium castaneum* (Herbst) and *Sitophilus* zeamais Motsch. Journal of Stored Products Research, 34: 11-17.
- Huang, Y., Ho, S. H., Lee, H. C. and Yap, Y. L. 2002. Insecticidal properties of eugenol, iseugenol and methyl engenol and their effects on nutrition of *Sitophilus zeamais* Mostsch. (Col: Curculionidae) and *Tribolium castaneum* (Col: Tenebrionidae). Journal of Stored Products Research, 38: 403-412.
- Huang, Y., Tan, J. M. W. L., Kini, R. M. and Ho, S. H. 1997. Toxic and antifeedent action of nutmeg oil against *Tribulium castaneum* (Herbst.) and *Sitophilus zeamais* Motsch. Journal of Stored Products Research, 33 (4): 289-298.
- Isman, M. B., Koul, O., Luczynski, A. and Kaminski, J. 1990. Insecticidal and antifeedant bioactivities of neem oils and their relationship to azadiractin content. Journal of Agricultural and Food Chemistry, 38: 1406-1411.
- Jermy, T. 1990. Prospects of antifeedant approach to pest control-A critical review. Journal of Chemical Ecology, 16 (11): 3151-3166.
- Johnso, J. A. and Vail, P. V. 1988. Post treatment survival, development and feeding of irradiated Indian meal moth and navel orange worm larvae (Lepidoptera: Pyralidae). Journal of Economic Entomology, 81: 376-380.
- Johnso, J. A. and Vail, P. V. 1989. Damage to raisins, almonds and walnuts by irradiated Indian meal moth and navel orange worm larvae (Lepidoptera: Pyralidae). Journal of Economic Entomology, 82: 1391-1394.
- Katz, S. H. and Weaver, W. W. 2003. Gale encyclopedia of Food and Culture 3, p. 712.

- Manuwoto, S. and Scriber, J. M. 1992. Consumption and utilization of three maize genotypes by the southern armyworm. Journal of Economic Entomology, 75: 163-167.
- Mehta, V. K., Sethi, G. R., Garg, A. K. and Seth, R. K. 2004. Use of ionizing radiation in interaction with fumigants towards management of *Tribolium castaneum* (Herbst). In Proceedings of the International Conference of Controlled Atmosphere and Fumigation in Stored Products., Gold-Coast Australia, pp. 467-474.
- Mensah, G. W. K., Watters, F. L. and Webster, G. R. B. 1979. Insecticide residues in milled fractions of dry or tough wheat treated with malathion, bromophos, iodofenphos and pirimiphos-methyl. Journal of Economic Entomology, 72: 728-731.
- Mills, J. and White, N. 1994. Seasonal occurrence of insects and mites in a Manitoba feed mill. Proceedings of the Entomological Society of Manitoba, 49: 1-15.
- Misra, H. P. and Paravathy, B. 1998. Gamma radiation susceptibility of strains of *Tribolium castaneum* (Herbst) resistant and susceptible to fenvalerate. International Journal of Pest Management, 44: 145-147.
- Negahban, M. and Moharramipour, S. 2007. Efficiency of *Artemisia sieberi* and *Artemisia scoparia* essential oils on nutritional indices of *Tribolium castaneum* (col: Tenebrionidae). Iranian Journal of Medicinal and Aromatic Plants Research, 23 (1): 13-22.

- Pszczola, D. E. 1997. 20 ways to market the concept of food irradiation. Food Technology, 51: 46-48.
- Rossi, E., Cosimi, S. and Loni, A. 2010. Insecticide resistance in Italian populations of *Tribolium* flour beetles. Bulletin of Insectology, 63 (2): 251-258.
- Tilton, E. W., Brower, J. H., Brown, G. A. and Kirkpatrick, R. L. 1972. Combination of gamma and microwave radiation for control of the Angoumois grain moth in wheat. Journal of Economic Entomology, 65: 531-533.
- Tuncbilek, A. S., Ayvaz, A., Ozturk, F. and Kaplan, B. 2003. Gamma radiation sensitivity of larvae and adults of the red flour beetle, *Tribolium castaneum* Herbst. Anzeiger furSchdlingskunde, 76 (5): 129-132.
- Urbain, W. M. 1986. Food Irradiation. Food Science and Technology. A Series of Monographs. Academic Press, London.
- Watters, F. L. and MacQueen, K. F. 1967. Effectiveness of gamma irradiation for control of five species of stored-product insects. Journal of Stored Products Research, 3: 223-234.
- Wiendl, F. M. 1984. A salubridade dos alimentos irradiados. Sociedade Brasileira de Ciência e Tecnologia de Alimentos, 18: 48-56.
- Xie, Y. S., Bodnaryk, R. P. and Fields, P. G. 1996. A rapid and simple flour disk bioassay for testing natural substances active against stored-product insects. Canadian Entomologist, 128: 865-875.

J. Crop Prot. (2015) Vol. 4 (4)

Ahmadi et al. _____

تأثیر ضد تغذیهای تلفیق پر تو گاما با اسانس برازمبل Perovskia atriplicifolia روی شپشه آرد (Coleoptera: Tenebrionidae)

مهرداد احمدی'*، سعید محرمی پور ٔ و آدلی محمد عبدالا ّ

۱- پژوهشکده کشاورزی هستهای، پژوهشگاه علوم و فنون هستهای، کرچ، ایران. ۲- گروه حشرهشناسی کشاورزی، دانشکده کشاورزی، دانشگاه تربیت مدرس، تهران، ایران. ۳- گروه آفات و گیاهپزشکی، مرکز مطالعات ملی، دوکی، گیزا، مصر. * پست الکترونیکی نویسنده مسئول مکاتبه: mahmadi@nrcam.org دریافت: ۶ مهر ۱۳۹۳؛ یذیرش: ۱۵ اردیبهشت ۱۳۹۴

چکیده: کنترل آفات انباری با استفاده از پرتو گاما میتواند با کمک سایر روشها مانند اسانسهای گیاهی که خود جایگزینی برای آفتکشهای شیمیایی میباشند، بهبود یابد. در این تحقیق، کارایی تلفیق پرتو گاما با اسانس برازمبل Perovskia atriplicifolia بهمنظور افزایش تأثیر ضدتغذیهای آنها روی شپشه آرد Tribolium castaneum مورد ارزیابی قرار گرفت. جهت ارزیابی شاخصهای تغذیهای مانند: نرخ رشد نسبی (RGR)، نرخ مصرف نسبی (RCR)، کارایی غذای خورده شده (ECI) و شاخص بازدارندگی تغذیهای (FDI)، از آزمایش زیستسنجی با کمک دیسکهای آردی استفاده گردید. نتایج نشان داد که یرتو گاما، اسانس برازمبل و نیز تلفیق این دو روش مقدار RCR ،RGR و ECI را در لارو و حشرات کامل بهطور معنیداری کاهش میدهند. بهطوریکه دز ۱۰۰ گری از پرتو گاما مقدار RGR لاروها و حشرات كامل را از ۰/۰۸۸۴ و ۰/۰۳۶۶ میلی گرم بر میلی گرم وزن حشره به ۰/۰۵۹۶ و ۰/۰۳۳۲ میلیگرم بر میلیگرم وزن حشره کاهش میدهد، این درحالی است که تلفیق این دز پرتو با اسانسهای گیاهی میزان کاهش را تا ۳۸۸۸-۰/۰۵۱ و ۲/۲۰۴-۰/۰۱ میلی گرم بر میلی گرم وزن حشره می ساند. همچنین مشخص گردید که شدت و میزان کاهش در حالت تلفیق در لاروها نسبت به حشرات کامل شپشه آرد شدیدتر است. نتایج نشان داد که مقدار FDI لاروها و حشرات کامل در اثر پرتو گاما، اسانس برازمبل و نیز تلفیق هر دو روش، افزایش معنیداری داشته است. همچنین شدت افزایش در FDI در حالت تلفیق بیشتر از زمانی است که هر دو روش به تنهایی مورد استفاده قرار گرفته بودند. بنابر این از یافتههای بهدست آمده می توان دریافت که پرتودهی می تواند اثرات ضدتغذیهای اسانس های گیاهی را افزایش دهد.

واژگان کلیدی: برازمبل، نرخ رشد نسبی، نرخ مصرف نسبی، کارایی غذای خورده شده، شاخص بازدارندگی تغذیه