

Research Article

Evaluation of six cucumber greenhouse cultivars for resistance to *Tetranychus turkestanii* (Acari: Tetranychidae)

Somaye Mohammadi^{1*}, Ali Asghar Seraj¹ and Ali Rajabpour²

1. Plant Protection Department, Faculty of Agriculture, Shahid Chamran University, Ahvaz, Iran.

2. Department of Plant Protection, Faculty of Agriculture, Ramin Agriculture and Natural Resources University of Khuzestan, Ahvaz, Iran.

Abstract: The strawberry spider mite (SSM), *Tetranychus turkestanii* Ugarov & Nikolski, is an important pest of greenhouse cucumber in tropical and temperate regions of the world. The use of the host plant resistance is a basic method to reduce pesticide application in greenhouses. In this study, the biological responses of SSM to six greenhouse cucumber cultivars (Puia, Hedieh, Milad Ghadim, Milad Jadid, Khasib and Negin) were investigated. Non-choice tests were performed to evaluate the interaction between the host plant-mite. All tests of this study were carried out under laboratory conditions at 28 ± 1 °C, $60 \pm 5\%$ RH and 16:8 h (L: D) in a walk-in growth chamber. The results indicated that immature development time and adult longevity of SSM was significantly influenced by cucumber cultivars. The highest r , λ , and R_0 were 0.275 (day^{-1}), 1.316 (day^{-1}) and 36.180 (offspring/generation) all of which were recorded for the mites reared on the Hedieh cultivar. No significant difference was observed for these parameters among the other cultivars. Mean generation times of *T. turkestanii* reared on Negin (14.020 day) and Milad Ghadim (13.57 day) cultivars were significantly longer than for the other cultivars. This mite had the shortest generation time on Puia cultivar (11.43 day). In conclusion, it seems that Hedieh and Milad Jadid cultivars are the more susceptible and resistant cultivars to the SSM than the other tested cultivars, respectively.

Keywords: host plant resistance, greenhouse cucumber, non-choice test, life table, strawberry spider mite

Introduction

The greenhouse cultivations are rising in recent years, at the same time the problems of pests are increasing as well. Cucumber (*Cucumis sativus* L.) is one of the most important and widely grown crops in greenhouse conditions (Rich *et al.*, 2013; Khaghani, 2009). The greenhouse crops are

affected by many pests that cause extensive yield losses (Zhang, 2003). Manufacturers use chemical pesticides to control the pests, and this process threatens the health of consumers, particularly because these crops are consumed as raw and fresh (Kosary and Kharazy Pakdel, 2006; Baniameri, 2003). Among the wide range of pests which attack to greenhouse crops, strawberry spider mite (SSM), *Tetranychus turkestanii* Ugarov & Nikolski, is an economically important pest (Martinez-Ferrer *et al.*, 2006). Mite population increases enormously in the constantly warm and moist conditions of

Handling Editor: Yaghoob Fathipour

* **Corresponding author**, e-mail: smentomology@gmail.com

Received: 26 May 2015, Accepted: 09 August 2015

Published online: 28 September 2015

greenhouses (Zhang, 2003). The pest further infests the underside of leaves, where it produces plenty of web and feeds in a piercing-sucking manner that damages plant cells and tissue. The feeding results in appearance of yellow chlorotic spots on leaves (Martinez-Ferrer *et al.*, 2006). One of the strategies for sustainable management of this pest would be the use of resistant host plants.

Host plant resistance can be a fundamental factor of a successful integrated pest management (IPM) system. Integration of resistant plants with other pest control methods results in conservation of beneficial natural enemies and decreased usage of pesticides (Lorenzen *et al.*, 2001). In the first step, recognition of resistance sources in plants is consequential (Jyoti *et al.*, 2001). Also, information about the type of resistance is essential for breeding programs (Stoner and Shelton, 1988). Plant resistance to a pest can be explained in three main assortments: antixenosis, antibiosis and tolerance, or some combinations of these mechanisms (Kogan and Ortman, 1978). Antibiosis is the most important because of its direct effect on the biology of a pest. Therefore, comparison of biological parameters of pest on different host plant cultivars can be used to select resistant cultivars to the pest (Ofomata *et al.*, 2000; Li *et al.*, 2004). Different host plant cultivars affect the life table parameters of SSM differently. Host plant resistance to *T. turkestanii* was reported on some host plants such as melon (Mansour *et al.*, 1987), raspberry (Wilde *et al.* 1991), strawberry (Gimenez-Ferrer *et al.* 1994), watermelon (Fadel *et al.*, 1994), tomato (Saeidi, 2006), bean (Mohammadi *et al.*, 2008), soybean (Sedaratian *et al.*, 2011), cotton (Kabiri *et al.*, 2012) and eggplant (Khanamani *et al.*, 2012, 2013). The establishment of spider mites on different cultivars is affected by diverse factors, including secondary metabolites, morphological structures, nutritive value of leaf and leaf age.

Defensive chemical compounds or poor-quality plants can make changes in life history and reproductive parameters of spider mites (Krips *et al.*, 1998; Agrawal, 2000; Gotoh and Gomi, 2003).

A life-table study provides two types of information including basic data and derived parameters (Jha *et al.*, 2012; Huang and Chi, 2012). The age-stage specific growth rate, age-stage specific developmental rate, age-stage specific survival rate and age-stage specific fecundity are the basic data that are calculated directly from the life history data of a cohort. These basic data are used to estimate derived parameters, such as the intrinsic rate of increase (r), finite rate of increase (λ), net reproductive rate (R_0) and mean generation time (T). The derived parameters are applicable only to a population with a stable age distribution, which is unlikely to occur under changing environmental conditions. According to basic data, an age-stage, two-sex life table offers a way to simulate population growth for a month without assuming a stable age distribution (Chi and Liu, 1985). A simulation based on an age-stage, two-sex life table represents the stage structures of both sexes in a pest population at any time (Chi, 1988, 1990; Tsai and Chi, 2007; Kavousi *et al.*, 2009; Huang and Chi, 2012).

The aim of the present study is to determine the effects of six greenhouse cucumber cultivars on life table parameters of strawberry spider mite. The data obtained from these studies are used to understand the mechanism of population build-up of pest on different cucumber cultivars and to develop a comprehensive pest management program for cucumber in greenhouses.

Materials and Methods

Cultures

In this study, six greenhouse cucumber (*Cucumis sativus*) cultivars, namely Puia, Hedieh, Milad Ghadim, Milad Jadid, Khasib and Negin were used as host plants. The

cultivars were selected by advisement of plant protection experts in Khuzestan province, Iran. Seeds were drenched 24 h. before planting. Then, seedlings were planted in pots (10 liters in size) that were filled with coco peat and perlite (1:1) in a hydroponic system. The pots were kept in a greenhouse at 28 ± 2 °C, with a light intensity of 13000 Lux and were fed by the nutrient solution (Rash) three times daily.

The initial populations of mites (*T. turkestanii*) were collected from infested European bindweed, *Convolvulus arvensis* L., in the agricultural College campus of Shahid Chamran University, Ahvaz, Iran. Collected mites were separately reared on cucumber leaves of each experimental cultivar for three generations.

The mite rearing and experiments were done in a walk-in growth chamber at 28 ± 1 °C, $60 \pm 5\%$ RH and a photoperiod of 16:8 (L: D) h.

Experimental unit

The effect of cucumber cultivars on life parameters of the spider mite and non-choice test was assessed on leaf discs (2 cm diameter) placed on soaked cotton in a Petri-dish (3 cm diameter), in which a 1.5 cm diameter hole was drilled and covered with ruching for ventilation. To keep the leaves fresh, distilled water was daily added to cotton.

Non-choice test

This experiment was carried out in a completely randomized design with six treatments (cucumber cultivars), each with six replications. Five female adult mites were released on the leaf disc per unit. After 72 h., the damage score of the leaf discs, the number of eggs and the percentage of dead mites were recorded. The damage was assessed according to a damage score method as explained by Nihoul *et al.* (1992), Gimenes-Ferrer and Scheerens (1993), Smith (2005) and Seaidi (2006).

Life table

Female and male spider-mite cohorts were formed by placing a pair of SSM on a leaf disc. The mites were eliminated after 3 h. and only one egg was maintained on each leaf disc. Each greenhouse cucumber cultivar was tested with 40 replicates. Units were checked every 12 h. (at 7 am and 7 pm). After adult emergence, females were coupled with males taken from the colony of the mite on the same cultivar. Then the duration of egg, larva, protonymph and deutonymph, adult longevity, fecundity and mortality of the mite were recorded for each cultivar until the death of the last mite.

Life table parameters

The age-stage survival rate (s_{xj} ; where x and j are age and stage, respectively), the age-stage fecundity (f_{xj}), the age-specific survival rate (l_x), and the age-specific fecundity (m_x), as well as the population parameters, the intrinsic rate of increase (r), the finite rate of increase (λ), the net reproductive rate (R_0), and the mean generation time (T), were estimated, according to Chi (2015). The intrinsic rate of increase (r) was calculated by using the iterative bisection method from the Euler-Lotka formula with the age indexed from zero (Goodman, 1982).

$$\sum_{x=0}^{\infty} e^{-r(x+1)} l_x m_x = 1 \quad (1)$$

Where l_x and m_x were computed according to Chi and Liu (1985) as (k is the number of stages):

$$l_x = \sum_{j=1}^k s_{xj} \quad (2)$$

The mean generation time is the time length that a population needs to increase to R_0 -times of its size as the stable age distribution and the stable increase rate are reached. Therefore, the mean generation time is calculated by $T = \ln R_0 / r$. The R_0 is estimated at:

$$R_0 = \sum_{x=0}^{\omega} \sum_{j=1}^{\kappa} s_{xj} f_{xj} \quad (3)$$

Statistical analysis

Mean compression of the data was subjected to a one-way Analysis of Variance (ANOVA) using the Proc GLM procedure of SAS 9.2 using the LSD tests at $P < 0.05$ (SAS Institute Inc, 2006). The data analysis and population parameters (r , λ , R_0 and T) were calculated by using the TWOSEX-MS Chart program and charts were drawn with Sigma Plot 12.0 (Chi, 2015). The standard errors were estimated by using the bootstrap technique and the differences of life table bootstrap-values among the treatments were compared with paired bootstrap test by TWOSEX-MS Chart program (Efron, 1979; Huang and Chi, 2012, 2013).

Result

Non-choice test

The result of the non-choice tests showed a significant difference between the mean mortality percentages ($F_{5, 24} = 9.20$; $P = 0.0001$), mean egg numbers ($F_{5, 24} = 29.67$; $P = 0.0001$) and mean damage scores ($F_{5, 24} = 22.13$; $P = 0.0001$) of *T. turkestanii* fed on different cucumber cultivars leaf disc after 72 h (Table 1). The highest mortality percentage was observed on the Milad Jadid cultivar (36 percent/72 h) and those of other cultivars had no significant difference. The SSM females laid the highest eggs on the Milad Ghadim cultivar during 72 h (119.60 eggs/72 h) and the lowest on the Milad Jadid cultivar (38.80 eggs/72 h). The highest and lowest damage scores were observed on the Hedieh and Milad Jadid cultivars leaf discs (4.40 and 0.80 damage score/leaf disc), respectively (Table 1). The following aspects of life history of SSM as

affected by mite-host plant interactions were studied in more details.

Developmental time and adult longevity

Effects of different greenhouse cucumber cultivars on the developmental time and adult longevity are presented in table 2. Significant differences among some biological traits such as egg ($F_{5, 54} = 18.32$; $P < 0.0001$), larvae ($F_{5, 54} = 17.30$; $P < 0.0001$), protonymphal ($F_{5, 54} = 12.45$; $P < 0.0001$) and deutonymphal ($F_{5, 54} = 12.10$; $P < 0.0001$) durations, total immature developmental time and adult longevity of SSM were observed on the different tested cultivars. The longest (3.52 days) and shortest (2.10 days) egg development were observed on Hedieh and Puia cultivars, respectively. The longer durations of larvae (2.28 days), protonymph (2.28 day) and deutonymph (2.00 days) of the SSM were observed on Puia cultivar.

Statistical comparison between the mean adult pre-oviposition periods (APOP) of *T. turkestanii* on six cucumber cultivars showed that this mite has the longest APOP on Milad Jadid cultivar (1.32 days) and the shortest on Hedieh cultivar (0.62 day) ($F_{5, 54} = 32.80$; $P < 0.0001$). The shortest total pre-oviposition period (TPOP) (from egg to first oviposition) however was observed on Milad Jadid (9.06 days) and Puia (9.14 days) and the longest on Negin (9.5 days) and Hedie (9.41 days) cultivars ($F_{5, 54} = 3.44$; $P < 0.009$). It seems that Negin and Hedieh cultivars lead to increase in duration of the SSM immature life stages. Also, the female adult longevity was shortest (12.77 days) on Milad Jadid and longest on Hedieh cultivar (16.26 days) ($F_{5, 54} = 63.67$; $P < 0.0001$). On the whole, it seems that developmental time and adult longevity of this pest on Hedieh cultivar are longer than those of the other cultivars.

Table 1 Mean (\pm SE) percent mortality, number of eggs and damage score of *Tetranychus turkestanii* fed on the six tested greenhouse cucumber cultivars in the non-choice tests under laboratory conditions.

Cultivars	Mortality (%) (in 72 h)	No. eggs (in 72 h)	Damage score on leaf disk (in 72 h)
Puia	0.00 ± 0.00b*	45.20 ± 2.94c	4.00 ± 0.45ab
Hedieh	8.00 ± 4.91b	74.80 ± 8.02b	4.40 ± 0.25a
Milad Jadid	36.00 ± 7.50a	38.80 ± 7.43c	0.80 ± 0.20d
Milad Ghadim	0.00 ± 0.00b	119.60 ± 6.04a	3.20 ± 0.20b
Khasib	2.00 ± 0.91b	75.20 ± 3.08b	3.20 ± 0.38b
Negin	4.00 ± 1.40b	104.20 ± 5.52a	2.00 ± 0.00c

* The means followed by the same letters in each column are not significantly different (LSD, $P < 0.05$).

Table 2 Mean (\pm SE) of *Tetranychus turkestanii* development time and adult longevity reared on the six tested greenhouse cucumber cultivars in laboratory condition.

Development time (day)	Puia	Hedieh	Milad Jadid	Milad Ghadim	Khasib	Negin
Egg	2.10 ± 0.04d*	3.52 ± 0.76a	2.55 ± 0.08c	2.95 ± 0.03b	2.92 ± 0.03b	2.93 ± 0.03b
Larval	2.28 ± 0.10a	1.77 ± 0.10c	2.05 ± 0.08b	2.10 ± 0.06b	2.05 ± 0.05b	2.08 ± 0.04b
Protonymph	2.16 ± 0.12a	1.60 ± 0.08b	1.78 ± 0.08b	1.63 ± 0.08b	1.18 ± 0.06c	1.77 ± 0.09b
Deutonymph	2.00 ± 0.12a	1.65 ± 0.09b	1.48 ± 0.55b	1.60 ± 0.11b	1.59 ± 0.09a	1.29 ± 0.10a
Total immature	8.58 ± 0.16ab	8.55 ± 0.73ab	7.82 ± 0.09d	8.30 ± 0.12b	8.14 ± 0.07d	8.71 ± 0.60a
APOP**	0.91 ± 0.18c	0.62 ± 0.13d	1.32 ± 0.25a	1.00 ± 0.14bc	1.12 ± 0.10b	1.15 ± 0.18b
TPOP***	9.04 ± 0.23c	9.41 ± 1.03ab	9.05 ± 0.25c	9.30 ± 0.17abc	9.24 ± 0.13bc	9.50 ± 0.17a
Adult longevity (Male)	10.50 ± 0.5b	11.80 ± 0.58a	10.00 ± 0.58c	12.00 ± 1.00a	12.00 ± 0.44a	11.83 ± 0.7a
Adult longevity (Female)	13.17 ± 0.40cd	16.26 ± 1.13a	12.77 ± 0.57d	13.82 ± 0.67b	13.57 ± 0.48bc	13.91 ± 0.63b
Total longevity	13.08 ± 0.47cd	15.60 ± 1.01a	12.49 ± 0.52d	13.25 ± 0.63bc	13.14 ± 0.46bcd	13.76 ± 0.56b

Notes: * The means followed by the same letters in each row are not significantly different (LSD, $P < 0.05$). **APOP, adult pre-ovipositional period; *** TPOP, total pre-ovipositional period (from egg to first oviposition).

Survival and mortality rate

The result showed that there was a difference among mortality and survival rates of *T. turkestanii* reared on the six cucumber cultivars, comparatively (Fig. 1). Mortality of this pest on the tested cucumber cultivars nearly occurred between 5-6 days, except that it started on the 7 day on Hedieh cultivar (Fig. 1). The mortality rate of this pest on cultivar Hedieh ascended with a quick slope between 7-22 days and was fixed from 22nd day till closing date (47th day). On the other cultivars, mortality started on 5th or 6th day and increased till closing date (47th day) (Fig. 1). These results suggest that survival rate of *T. turkestanii* fed on Hedieh cultivar was comparatively slowed down.

Population projection

The population growth rates of the SSM reared on the six greenhouse cucumber cultivars were projected. The resulting figures revealed the change in the age-stage structure of *T. turkestanii* reared on the mentioned host plants. After the second peak, the trend of population growth of SSM on the Hedieh cultivar was equable in the each stage and then the population increased. However, population growth of this pest on the other cultivars was descending after the second peak (Fig. 2).

Life table parameters

The result showed that there was a significant difference between the intrinsic rate of increase (r) ($F_{4, 22} = 4.22$; $P = 0.0012$), the finite rate of increase (λ) ($F_{5, 54} = 4.43$; $P =$

0.0008) and the net reproductive rate (R_0) ($F_{5, 54} = 6.77$; $P = 0.0001$) of the SSM fed on the different tested cultivars (Table 3). The highest r , λ , and R_0 were 0.275 (day^{-1}), 1.316 (day^{-1}) and 36.18 (offspring/generation) recorded for the mites that were reared on the Hedieh cultivar. No significant difference was observed for these parameters among the other

cultivars. Mean generation time of *T. turkestanii* reared on Negin (14.020 day) and Milad Ghadim (13.570 day) cultivars was significantly longer than that of other cultivars ($F_{5, 54} = 4.86$; $P = 0.0003$). This parameter was least for mites that reared on Puia cultivar (11.430 day) (Table 3).




Figure 1 Age-specific survival rate (solid white circle, left Y-axis) and mortality rate (solid black circle, right Y-axis) of *Tetranychus turkestanii* reared on six greenhouse cucumber cultivars in laboratory conditions.




Figure 2 Population projection of age-stage structure of *Tetranychus turkestani* reared on the six greenhouse cucumber cultivars during population growth.

Table 3 Mean (\pm SE) population parameters of *Tetranychus turkestani* reared on six cucumber greenhouse cultivars.

Cucumber cultivars	Population parameter (means \pm SE) ¹			
	Intrinsic rate of increase (day ⁻¹)	Finite rate of increase (day ⁻¹)	Net reproductive rate (offspring/generation)	Mean generation time (day)
Puia	0.183 \pm 0.026b*	1.201 \pm 0.03b	8.15 \pm 1.85b	11.43 \pm 0.41c
Hedieh	0.275 \pm 0.26a	1.316 \pm 0.02a	36.18 \pm 5.19a	13.06 \pm 0.27ab
Milad Jadid	0.207 \pm 0.02b	1.233 \pm 0.03b	15.050 \pm 4.05b	13.04 \pm 0.43ab
Milad Ghadim	0.199 \pm 0.02b	1.221 \pm 0.02b	15.05 \pm 4.05b	13.57 \pm 0.42a
Khasib	0.209 \pm 0.02b	1.232 \pm 0.02b	13.433 \pm 2.37b	12.40 \pm 0.23b
Negin	0.184 \pm 0.02b	1.200 \pm 0.03b	13.333 \pm 3.09b	14.02 \pm 0.29a

¹The means followed by the same letters in each column are not significantly different (Paired bootstrap test, P < 0.05).

Discussion

The main purpose of this study was to evaluate resistance of six greenhouse cucumber cultivars to the strawberry spider mite. The pests have different survival, development and reproductive rates on the plant species and cultivars because plants differ greatly in suitability as hosts. Shorter development time and high reproduction of pests on host plants indicate greater suitability of those plants for colonization of pests (van Lenteren and Noldus 1990). Previous studies demonstrated that not only the quantity, but also the quality of food sources is a consequential factor for growth and development of all organisms (Eischen and Dietz, 1987; Hagley and Barber, 1992, Awmack and Leather, 2002; Blanco *et al.*, 2006; Lee, 2007; Hwang *et al.*, 2008). The results of this study confirmed the previous studies regarding effect of quality and quantity of host plant on the pest biology.

Egg development time of the SSM showed significant difference in the various tested cucumber cultivars. This means that embryonic development of eggs laid by mite females on the different cucumber cultivars was affected by the feeding of females before oviposition. However, Sedaratian *et al.* (2009) pointed out that egg incubation period of *T. urticae* was not significantly different on soybean genotypes. These noticeable differences might be related to differences in, for example the host plant and mite species.

Mean development time of *T. turkestanii* varied significantly on the tested cucumber cultivars. According to previous studies, these differences could be due to variation in plant quality, particularly host plant nutrient levels and secondary compounds (Awmack and Leather, 2002; Blanco *et al.*, 2006; Lee, 2007; Hwang *et al.*, 2008).

Khanamani *et al.* (2013) and Kabiri *et al.* (2012) reported that developmental time of *T. urticae* fed on the resistant eggplant and cotton cultivars was longer than on susceptible cultivars. Uckan and Ergin (2002) suggested that increasing the life stage duration of pests

may be a major fitness for the pest for survival of a generation when food is inadequate, because only a limited number of females have the potentiality to survive. In this study *T. turkestanii* had a longer development time on Hedieh cultivar and hence it seems this cultivar may be susceptible to the SSM while it was shorter on Milad Jadid cultivar which appears to be resistant. Different responses of this mite to cucumber, eggplant, cotton and or other host plants may be due to the different mite species or variant host plant.

The range of the value of r for the mites that fed on the six greenhouse cucumber cultivars varied from 0.183 to 0.275 (day^{-1}), which is nearly consistent with reported results by Helle and Sabelis (1985). Khanamani *et al.* (2013) reported that the intrinsic rate of increase of two spotted spider mite varied from 0.022 to 0.157 (day^{-1}) on different eggplant cultivars. *T. urticae* reared on soybean genotypes had arranged between 0.392 and 0.236 (day^{-1}) (Sedaratian *et al.*, 2011). The declared values of R_0 on the different eggplant cultivars by Khanamani *et al.*, (2013) varied from 1.425 to 11.585 (offspring/generation), but the range of parameter for the SSM in our study was wider (8.150 to 36.180 offspring/generation) and on soybean genotypes was 45.521 at 12.149 (offspring/generation) (Sedaratian *et al.*, 2011). These results showed that the cucumber cultivars may be a more suitable host plant for the larva and nymph of the spider mite than eggplant cultivars but not better than soybean genotypes.

The population parameters (r , R_0 , λ and T) are the derived parameters. They are calculated based on the presumption that the environmental factors are constant and the population structure reaches a stable age-stage distribution as time approaches infinity (Huang and Chi, 2012). The intrinsic rate is a good parameter to reveal and compare the potential of insect populations under different treatments. Its practical application in pest management, however, is limited. The basic data, including survival rate, developmental rate, and fecundity, describe the life history and stage

differentiation. These data can be used in population projection a stop predicts growth trends, as well as the stage structure of a population in the short-term or long-term future.

These laboratory results may be related to antibiotic or antixenotic effects (or their combinations) of host plants on both life history and fecundity of the SSM. This study did not assay the effect of chemical compounds or morphological structure of these cultivars on biological parameters of *T. turkestanii*. Therefore further work is imperative to explain whether the differences observed in mentioned parameters in our study are related to the chemical compounds, the morphological structure of the plants or to both. As well as obtaining more applied results of field experiments should be considered in future.

In conclusion, the use of resistant and moderately resistant cultivars with biological and chemical control methods can help to success an IPM program. Different levels of resistance to *T. turkestanii* were observed in Milad Jadid, Puia, Negin, Khasib and Milad Ghadim cultivars. Hedieh and Milad Jadid cultivars were the more susceptible and resistant cultivars to the SSM than other tested cultivars, respectively.

Acknowledgements

This research has been supported by Department of Plant Protection, Shahid Chamran University of Ahvaz, Iran, which is greatly appreciated. We also express our appreciation to Prof. Chi providing us his software.

Reference

- Agrawal, A. A. 2000. Host-range evolution: adaptation and trade-offs in fitness of mites on alternative hosts. *Ecology*, 81: 500-508.
- Awmack, C. S., Leather, S. R. 2002. Host plant quality and fecundity in herbivorous insects. *Annual Review of Entomology*, 47: 817-844.
- Baniameri, V. 2003. Study on life characteristics and predatory behavior of predator bug *Orius niger* Wolff on onion thrips *Thrips tabaci* Lind under lab conditions. Ph. D. Dissertation. Shahid Chamran University. 120 pp.
- Blanco, C. A., Teran-Vargas, A. P., Abel, Portilla, C. A. M., Rojas, M. G., Morales-Ramos, J. A. and Snodgrass, G. L. 2006. Plant host effect on the development of *Heliothis virescens* F. (Lepidoptera: Noctuidae). *Environmental Entomology*, 37: 1538-1547.
- Chi, H. 1988. Life-table analysis incorporating both sexes and variable development rate among individuals. *Environmental Entomology*, 17: 26-34.
- Chi, H. 1990. Timing of control based on the stage structure of pest populations: a simulation approach. *Journal of Economic Entomology*, 83: 1143-1150.
- Chi, H. 2015. TWO SEX-MS Chart: A computer program for the age-stage, two-sex life table analysis. Available on: <http://140.120.197.173/Ecology/Download/TwoSexMSChart.zip>
- Chi, H. and Liu, H. 1985. Two new methods for the study of insect population ecology. *Bulletin of the Institute of Zoology, Academia Sinica*, 24: 225-240.
- Efron, B. 1979. Bootstrap methods: another look at the jackknife. *Annual Review of Entomology*, 7: 1-26.
- Eischen, F. and Dietz, A. 1987. Growth and survival of *Galleria mellonella* (Lepidoptera: Pyralidae) larva fed diets containing honey bee-collected plant resins. *Annals of the Entomological Society of America*, 80: 74-77.
- Fadel, M., Shain, Z., Karchi, Z. and Grson, U. 1994. Resistance of selected melon lines to the carmine spider mite *Tetranychus cinnabarinus* (Acari: Tetranychidae) field and laboratory experiments. *Bulletin of Entomological Research*, 84: 265-267.
- Gimenes-Ferrer, R. M., Scheerens, J. C. and Erb, W. A. 1993. In vitro screening of 76 strawberry cultivars for two-spotted spider mite resistance. *Horticultural Science*, 28: 841-844.
- Gimenez-Ferrer, R. M., Erb, W. A., Bishop, B. L. and Scheerens, J. C. 1994. Host-pest

- relationship between the two spotted spider mite and strawberry cultivars with differing levels of resistance. *Journal of Economic Entomology*, 87: 168-175.
- Goodman, D. 1982. Optimal life histories, optimal notation, and the value of reproductive value. *American Naturalist*, 119: 803-823.
- Gotoh, T., and K. Gomi. 2003. Life-history traits of the Kanzawa spider mite *Tetranychus kanzawai* (Acari: Tetranychidae). *Applied Entomology and Zoology*, 38: 7-14.
- Hagley, E. A. C. and Barber, D. R. 1992. Effect of food sources on the longevity and fecundity of *Pholetes orornigis* (Weed) (Hymenoptera: Braconidae). *Canadian Entomologist*, 124: 341-346.
- Helle, W. and Sabelis, M. W. 1985. Spider mites their biology, natural enemies and control. Elsevier Science Publisher, 405 pp. <http://140.120.197.173/Ecology/Download/TwoSexMSChart.zip>. (Accessed 10, February, 2015).
- Huang, Y. B. and Chi, H. 2012. Life tables of *Bactrocera cucurbitae* (Diptera: Tephritidae): with an invalidation of the jackknife technique. *Journal of Applied Entomology*, 137: 327-339.
- Huang, Y. B. and Chi, H. 2013. Age-stage, two-sex life tables of *Bactrocera cucurbitae* (Coquillett) (Diptera: Tephritidae) with a discussion on the problem of applying female age-specific life tables to insect populations. *Journal of Insect Science*, 19: 263-273.
- Hwang, S. Y., Liu, C. H. and Shen, T. C. 2008. Effects of plant nutrient availability and host plant species on the performance of two Pieris butterflies (Lepidoptera: Pieridae). *Biochemical Systematics and Ecology*, 36: 505-513.
- Jha, R. K., Chi, H. and Tang, L. 2012. Stage-structured population growth of *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae): a simulation approach. *Journal of Formosan Entomology*, 32: 139-153.
- Jyoti, J., Shelton, A. and Earle, E. 2001. Identifying sources and mechanisms of resistance in crucifers for control of cabbage maggot (Diptera: Anthomyiidae). *Journal of Economic Entomology*, 94: 942-949.
- Kabiri, H. Saboori, A. and Allahyari, H. 2012. Impact of different cotton (*Gossypium* spp.) cultivars, as host plant, on development and fertility life-table parameters of *Tetranychus urticae* (Acari: Tetranychidae). *International Journal of Acarology*, 38 (1): 46-50.
- Kavousi, A., Chi, H., Talebi, K., Bandani, A., Ashouri, A. and Naveh, V. H. 2009. Demographic traits of *Tetranychus urticae* (Acari: Tetranychidae) on leaf discs and whole leaves. *Journal of Formosan Entomology*, 102: 595-601.
- Khanamani, M., Fathipour, Y., Hajiqanbar, H. and Sedaratian, A. 2012. Reproductive performance and life expectancy of *Tetranychus urticae* (Acari: Tetranychidae) on seven eggplant cultivars. *Journal of Crop Protection*, 1 (1): 57-66.
- Khanamani, M., Fathipour, Y. and Hajiqanbar, H. 2013. Population growth response of *Tetranychus urticae* to eggplant quality: application of female age-specific and age-stage two sex life tables. *International Journal of Acarology*, 39 (8): 638-648.
- Kogan, M. and Ortman, E. F. 1978. Antixenosis a new term proposed to replace painter's 'non-preference' modality of resistance. *Bulletin of the Entomological Society of America*, 24: 175-176.
- Kosary, A. A. and KharaziPakdel, A. 2006. Preferences prey of *Orius albidipennis* on onion thrips and two spotted-spider mite under lab conditions. *Journal of entomological society of Iran*, 26(1): 73-91.
- Krips, O. E., A. P. Witul, E. L. Willems, and M. Dicke. 1998. Intrinsic rate of population increase of the spider mite *Tetranychus urticae* on the ornamental crop gerbera: intraspecific variation in host plant and herbivore. *Entomologia Experimentalis et Applicata*, 89: 159-168.
- Lee, K. P. 2007. The interactive effects of protein quality and macronutrient imbalance on nutrient balancing in an insect herbivore. *Journal of Experimental Biology*, 210: 3236-3244.

- Li, Y., Hill, C. B. and Hartman, G. L. 2004. Effect of three resistant soybean genotype on the fecundity, mortality, and maturation of soybean aphid (Homoptera: Aphididae). *Journal of Economic Entomology*, 97: 1106-1111.
- Lorenzen, J., Balbyshev, N., Lafta, A., Casper, H., Tian, X. and Sagredo, B. 2001. Resistant potato selections contain leptine and inhibit development of the Colorado potato beetle (Coleoptera: Chrysomelidae). *Journal of Economic Entomology*, 94: 1260-1267.
- Mansour, F., Karchi, Z. and Omari, N. 1987. Resistance of melon to the carmine spider mite, *Tetranychus cinnabarinus* (Boisduval) (Acari: Tetranychidae). *Bulletin of Entomological Research*, 77: 603-607.
- Martinez-Ferrer, M., Jacas, J., Piolles-Moles, J. and Aucejo-Romero, S. 2006. Approaches for sampling the two spotted spider mite (Acari: Tetranychidae) on clementines in Spain. *Journal of Economic Entomology*, 99: 1490-1499.
- Mohammadi, S. 2008. Resistance of different red and white bean genotypes to two-spotted spider mite, *Tetranychus urticae* Koch. (Acari: Tetranychidae), in Shahrekord. M. SC. Thesis, Shahid Chamran University of Ahvaz. 96 pp.
- Nihoul, P. T., Hance, B. M. and Van-Impe, G. 1992. Physiological aspect of damage caused by spider mite on tomato leaflets. *Journal of Applied Entomology*, 113 (5): 487-492.
- Ofomata, V. C., Overholt, W. A., Lux, S. A., Huis, A. V. and Egwuatu, R. I. 2000. Comparative studies on the fecundity, egg survival, larval feeding, and development of *Chilo orichalcociliellus* (Lepidoptera: Crambidae) on five grasses. *Annals of the Entomological Society of America*, 93:492-499.
- Rich, J. R., Webb, S. E., Paret, M. L. and Momol, M. T. 2013. Considerations for managing greenhouse pests-Vegetable production handbook. IFAS Extension, University of Florida, 3.
- Khaghani, B. 2009. Guide to Greenhouse Cultivation and General Information about Vegetable. Publications Hamim, 148.
- Saeidi, Z. 2006. Nature of resistance to two-spotted spider mite *Tetranychus urticae* Koch. (Acari: Tetranychidae) in *Lycopersicon* Species. Ph. D. Dissertation, University of Agricultural Sciences Bangalore, India. pp. 159.
- SAS Institute. 2006. GLM: A Guide to Statistical and Data Analysis, Version 9.2. SAS Institute, Cary.
- Sedaratian, A., Fathipour, Y. and Moharrampour, S. 2009. Evaluation of resistance in 14 soybean genotypes to *Tetranychus urticae* (Acari: Tetranychidae). *Journal of Pest Science*, 82: 163-170.
- Sedaratian, A., Fathipour, Y. and Moharrampour, S. 2011. Comparative life table analysis of *Tetranychus urticae* (Acari: Tetranychidae) on 14 soybean genotypes. *Insect Science*, 18: 541-553.
- Smith, M. C. 2005. Plant Resistance to Arthropods Molecular and Conventional Approaches. Springer, 421 pp.
- Stoner, K. A. and Shelton, A. M. 1988. Role of non-preference in the resistance of cabbage varieties to the onion thrips (Thysanoptera: Thripidae). *Journal of Economic Entomology*, 81: 1062-1067.
- Tsai, T. J. and Chi, H. 2007. Temperature-dependent demography of *Supella longipalpa* (Blattodea: Blattellidae). *Journal of Medical Entomology*, 44: 772-778.
- Uckan, F. and Ergin, E. 2002. Effect of host diet on the immature developmental time, fecundity, sex ratio, adult longevity, and body size of *Apanteles galleriae* (Hymenoptera: Braconidae). *Environmental Entomology*, 13: 168-171.
- Van Lenteren, J. C. and Noldus, L. P. J. J. 1990. Whitefly-plant relationship: behavioral and biological aspects. In: Gerling, D. (Ed), Whitefly: their Bionomics, Pest Status and Management. Intercept, Andover, pp 47-89.
- Wilde, G., Thomas, W. and Hall, H. 1991. Plant resistance to two-spotted spider mite (Acari: Tetranychidae) in raspberry cultivars. *Journal of Economic Entomology*, 84 (5): 251-255.
- Zhang, Z. Q. 2003. Mites of greenhouses: Identification, biology and control. CABI International. 256 p.

ارزیابی مقاومت شش رقم خیار گلخانه‌ای نسبت به *Tetranychus turkestanii* (Acari: Tetranychidae)

سمیه محمدی^{۱*}، علی اصغر سراج^۱ و علی رجب پور^۲

۱- گروه گیاهپزشکی، دانشکده کشاورزی، دانشگاه شهید چمران اهواز، اهواز، ایران.

۲- گروه گیاهپزشکی، دانشگاه کشاورزی و منابع طبیعی رامین خوزستان، ایران.

* پست الکترونیکی نویسنده مسئول مکاتبه: smentomology@gmail.com

دریافت: ۵ خرداد ۱۳۹۴؛ پذیرش: ۱۸ مرداد ۱۳۹۴

چکیده: کنه تارتن ترکستانی (*Tetranychus turkestanii* Ugarov & Nikolski (SSM)) آفت مهم خیار گلخانه‌ای در مناطق گرمسیری و معتدل جهان است. یکی از روش‌های مهم جهت کاهش مصرف آفت‌کش‌ها در گلخانه‌ها استفاده از میزبان گیاهی مقاوم می‌باشد. در این تحقیق به بررسی واکنش‌های زیستی کنه تارتن ترکستانی روی شش رقم خیار گلخانه‌ای (پویا، هدیه، میلاد قدیم، میلاد جدید، خسیب و نگین) پرداخته شد. برای ارزیابی برهمکنش میزبان گیاهی کنه، از تست غیرانتخابی استفاده گردید. تمامی آزمایش‌های این تحقیق در اتاقک رشد با شرایط دمایی 1 ± 28 درجه سانتی‌گراد، رطوبت نسبی 5 ± 60 درصد و نسبت روشنایی به تاریکی ۸:۱۶ ساعت انجام شد. نتایج نشان داد که مجموع مدت زمان رشد و نمو کنه در مرحله نابالغ و طول زندگی بالغین آن به‌طور معنی‌داری تحت تأثیر ارقام متفاوت خیار بود. بیش‌ترین R_0 و λ کنه تارتن ترکستانی روی رقم هدیه مشاهده شد (به ترتیب ۰/۲۷۵ بر روز، ۱/۳۱۶ بر روز و ۳۶/۱۸۰ نتاج/نسل). اما تفاوت معنی‌داری برای پارامترهای مذکور کنه روی ارقام دیگر خیار مشاهده نشد. مدت زمان تکمیل یک نسل کنه تارتن پرورش یافته روی رقم نگین (۱۴/۰۲۰ روز) و میلاد قدیم (۱۳/۵۷ روز) به‌طور معنی‌داری طولانی‌تر از ارقام دیگر بود. این کنه دارای کم‌ترین مدت زمان تکمیل یک نسل روی رقم پویا (۱۱/۴۳ روز) بود. به‌طور کلی به‌نظر می‌رسد ارقام هدیه و میلاد جدید به ترتیب حساسیت و مقاومت بیشتری به کنه تارتن ترکستانی نسبت به دیگر ارقام مورد آزمایش دارند.

واژگان کلیدی: مقاومت میزبان گیاهی، خیار گلخانه‌ای، آزمایش غیر انتخابی، جدول زندگی، کنه تارتن ترکستانی