

Expansion of Bessel and g -Bessel sequences to dual frames and dual g -frames

M. S. Asgari^a and G. Kavian^{a,*}

^aDepartment of Mathematics, Faculty of Science, Islamic Azad University,
Central Tehran Branch, PO. Code 14168-94351,
Tehran, Iran

Abstract. In this paper we study the duality of Bessel and g -Bessel sequences in Hilbert spaces. We show that a Bessel sequence is an inner summand of a frame and the sum of any Bessel sequence with Bessel bound less than one with a Parseval frame is a frame. Next we develop this results to the g -frame situation.

© 2013 IAUCTB. All rights reserved.

Keywords: G -frames, dual frames, dual g -frames.

2010 AMS Subject Classification: 46H20, 46H25, 46G05.

1. Introduction

Let \mathcal{H} denote a separable Hilbert space. A sequence $\{f_i\}_{i \in I}$ in \mathcal{H} is called a frame if there exist constants $0 < A \leq B < \infty$ such that

$$A\|f\|^2 \leq \sum_{i \in I} |\langle f, f_i \rangle|^2 \leq B\|f\|^2 \quad \forall f \in \mathcal{H}. \quad (1)$$

We call A and B the lower and upper frame bounds, respectively. The sequence $\{f_i\}_{i \in I}$ is a Bessel sequence if at least the upper bound in (1) is satisfied. For any frame $\{f_i\}_{i \in I}$ there exists at least one dual frame, i.e., a frame $\{g_i\}_{i \in I}$ for which $f = \sum_{i \in I} \langle f, g_i \rangle f_i$ for all $f \in \mathcal{H}$. If $\{f_i\}_{i \in I}$ is a Bessel sequence with bound $B < 1$, how can we find two

*Corresponding author.

E-mail addresses: msasgari@yahoo.com (M. S. Asgari), kaviangolsa@yahoo.com (G. Kavian).

sequences $\{g_i\}_{i \in I}$ and $\{p_i\}_{i \in I}$ such that $\{f_i + g_i\}_{i \in I}$ and $\{p_i\}_{i \in I}$ are dual frames, i.e., such that

$$f = \sum_{i \in I} \langle f, p_i \rangle (f_i + g_i) = \sum_{i \in I} \langle f, f_i + g_i \rangle p_i. \quad (2)$$

In this paper we aim at the more general results of the type in (2). For any Bessel sequence $\mathcal{F} = \{f_i\}_{i \in I}$ the synthesis operator is defined as follows:

$$T_{\mathcal{F}} : \ell^2(I) \rightarrow \mathcal{H}, \quad \text{with} \quad T_{\mathcal{F}}(\{c_i\}_{i \in I}) = \sum_{i \in I} c_i f_i.$$

The analysis operator for \mathcal{F} is $T_{\mathcal{F}}^*$ and is given by $T_{\mathcal{F}}^* f = \{\langle f, f_i \rangle\}_{i \in I}$. The frame operator is the positive self-adjoint invertible operator $S_{\mathcal{F}} = T_{\mathcal{F}} T_{\mathcal{F}}^*$ and satisfies $S_{\mathcal{F}} f = \sum_{i \in I} \langle f, f_i \rangle f_i$. The reconstruction formulas are as follows:

$$f = \sum_{i \in I} \langle f, f_i \rangle S_{\mathcal{F}}^{-1} f_i = \sum_{i \in I} \langle f, S_{\mathcal{F}}^{-1} f_i \rangle f_i \quad \forall f \in \mathcal{H}. \quad (3)$$

Frames were first introduced in 1952 by Duffin and Schaeffer [4] in the study of non-harmonic Fourier series, reintroduced in 1986 by Daubechies, Grossman and Meyer in [3]. G -frames for Hilbert spaces first formally were defined by Sun in [6].

Let \mathcal{H}, \mathcal{K} be two separable Hilbert spaces and $\{W_i\}_{i \in I}$ be a sequence of closed subspaces of \mathcal{K} , where I is a subset of \mathbb{Z} . Let $\mathcal{L}(\mathcal{H}, W_i)$ be the collection of all bounded linear operators from \mathcal{H} into W_i . Recall that a family of operators $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, W_i) : i \in I\}$ is said to be a generalized frame, or simply a g -frame for \mathcal{H} with respect to $\{W_i\}_{i \in I}$ if there exist constants $0 < C \leq D < \infty$ such that

$$C \|f\|^2 \leq \sum_{i \in I} \|\Lambda_i f\|^2 \leq D \|f\|^2 \quad \forall f \in \mathcal{H}. \quad (4)$$

The constants C and D are called g -frame bounds and $\sup_{i \in I} \Lambda_i$ is called the multiplicity of the g -frame. We call Λ a tight g -frame if $C = D$ and it is a Parseval g -frame if $C = D = 1$. If the right-hand side of (4) holds, then Λ is said a g -Bessel sequence for \mathcal{H} with respect to $\{W_i\}_{i \in I}$. The representation space associated with a g -Bessel sequence $\Lambda = \{\Lambda_i\}_{i \in I}$ is defined by

$$\left(\sum_{i \in I} \oplus W_i \right)_{\ell^2} = \left\{ \{g_i\}_{i \in I} \mid g_i \in W_i \text{ and } \sum_{i \in I} \|g_i\|^2 < \infty \right\}. \quad (5)$$

The synthesis operator of Λ given by

$$T_{\Lambda} : \left(\sum_{i \in I} \oplus W_i \right)_{\ell^2} \rightarrow \mathcal{H} \quad T_{\Lambda}(\{g_i\}_{i \in I}) = \sum_{i \in I} \Lambda_i^* g_i.$$

The adjoint operator of T_{Λ} , which is called the analysis operator also obtain as follows

$$T_{\Lambda}^* : \mathcal{H} \rightarrow \left(\sum_{i \in I} \oplus W_i \right)_{\ell^2} \quad T_{\Lambda}^* f = \{\Lambda_i f\}_{i \in I}.$$

By composing T_Λ with its adjoint T_Λ^* , we obtain the g -frame operator

$$S_\Lambda : \mathcal{H} \rightarrow \mathcal{H} \quad S_\Lambda f = T_\Lambda T_\Lambda^* f = \sum_{i \in I} \Lambda_i^* \Lambda_i f,$$

which is a bounded, self-adjoint, positive and invertible operator and $CI_{\mathcal{H}} \leq S_\Lambda \leq DI_{\mathcal{H}}$. The canonical dual g -frame for $\{\Lambda_i\}_{i \in I}$ is defined by $\{\tilde{\Lambda}_i\}_{i \in I}$ where $\tilde{\Lambda}_i = \Lambda_i S_\Lambda^{-1}$, which is also a g -frame for \mathcal{H} with g -frame bounds $\frac{1}{D}$ and $\frac{1}{C}$, respectively. The reconstruction formulas are also as follows:

$$f = \sum_{i \in I} \Lambda_i^* \tilde{\Lambda}_i f = \sum_{i \in I} \tilde{\Lambda}_i^* \Lambda_i f \quad \forall f \in \mathcal{H}.$$

For more details about the theory and applications of frames we refer the readers to [1, 5], about g -frames to [5, 6].

For using of the reconstruction formulas we need to invert the g -frame operator, which can be complicated. In the following similar to frame algorithm we use a g -frame algorithm to obtain approximations of linear operator $U \in B(\mathcal{H}, \mathcal{K})$.

Theorem 1.1 Let $\Lambda = \{\Lambda_i\}_{i \in I}$ be a g -frame with g -frame bounds C, D . For every $U \in B(\mathcal{H}, \mathcal{K})$, we define the sequence $\{U_n\}_{n \in \mathbb{N}}$ by

$$U_n = \begin{cases} 0 & n = 0 \\ U_{n-1} + \frac{2}{C+D}(U - U_{n-1})S_\Lambda & n \geq 1 \end{cases}$$

Then we have $U = \lim_{n \rightarrow \infty} U_n$ with the error estimate

$$\|U - U_n\| \leq \left(\frac{D-C}{D+C}\right)^n \|U\|.$$

Proof. By the g -frame condition for each $f \in \mathcal{H}$ we have

$$-\frac{D-C}{D+C} \|f\|^2 \leq \langle (I_{\mathcal{H}} - \frac{2}{C+D} S_\Lambda) f, f \rangle \leq \frac{D-C}{D+C} \|f\|^2.$$

Thus

$$\|Id_{\mathcal{H}} - \frac{2}{C+D} S_\Lambda\| \leq \frac{D-C}{D+C}.$$

Using the definition of $\{U_n\}_{n \in \mathbb{N}}$ we obtain

$$U - U_n = (U - U_0) \left(Id_{\mathcal{H}} - \frac{2}{C+D} S_\Lambda \right)^n.$$

Hence,

$$\|U - U_n\| \leq \left(\frac{D-C}{D+C}\right)^n \|U\|.$$

■

A family of operators $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, W_i) : i \in I\}$ is called an orthonormal g -basis for \mathcal{H} if it holds in the following conditions.

- (1) $f = \sum_{i \in I} \Lambda_i^* \Lambda_i f \quad \forall f \in \mathcal{H}.$
- (2) $\langle \Lambda_i^* g, \Lambda_j^* g' \rangle = \delta_{ij} \langle g, g' \rangle \quad \forall g, g' \in W_i, \forall i, j \in I.$

Lemma 1.2 Let $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, W_i) : i \in I\}$ be a collection of partial isometries. Then the sequence $\{\Lambda_i\}_{i \in I}$ is a Parseval g -frame for \mathcal{H} , if and only if the sequence $\{\Lambda_i^* \Lambda_i\}_{i \in I}$ be an orthonormal g -basis for \mathcal{H} .

Proof. First note that Λ_i is a partial isometry, if and only if $\Lambda_i^* \Lambda_i$ is an orthogonal projection on \mathcal{H} . Now the claim follows from

$$\|f\|^2 = \sum_{i \in I} \|\Lambda_i f\|^2 = \sum_{i \in I} \langle \Lambda_i^* \Lambda_i f, f \rangle = \sum_{i \in I} \|\Lambda_i^* \Lambda_i f\|^2 \quad \forall f \in \mathcal{H}.$$

■

2. Duality of Bessel and g -Bessel sequences

Li and Sun in [5] expanded every Bessel sequence to a tight frame by adding some elements. In this section we show that the sum of any Bessel sequence with Bessel bound less than one with a Parseval frame is a frame for \mathcal{H} and we prove that a Bessel sequence is an inner summand of a frame.

Let $\mathcal{F} = \{f_i\}_{i \in I}$ and $\mathcal{G} = \{g_i\}_{i \in I}$ be two Bessel sequences for \mathcal{H} with synthesis operators $T_{\mathcal{F}}$ and $T_{\mathcal{G}}$ respectively. Then we say that \mathcal{F} and \mathcal{G} are dual frames for \mathcal{H} if $T_{\mathcal{F}} T_{\mathcal{G}}^* = I_{\mathcal{H}}$ or $T_{\mathcal{G}} T_{\mathcal{F}}^* = I_{\mathcal{H}}$, i.e.,

$$f = \sum_{i \in I} \langle f, g_i \rangle f_i = \sum_{i \in I} \langle f, f_i \rangle g_i \quad \forall f \in \mathcal{H}.$$

Notation 2.1 For every countable (or finite) index set I , we define the space $\ell(I, \mathcal{H})$ by

$$\ell(I, \mathcal{H}) = \left\{ \{f_i\}_{i \in I} : f_i \in \mathcal{H}, \text{ and } \sup_{i \in I} \|f_i\| < \infty \right\}.$$

It is easy to check that $\ell(I, \mathcal{H})$ with the pointwise operations and norm defined by

$$\|\{f_i\}_{i \in I}\| = \sup_{i \in I} \|f_i\|,$$

is a Banach space. Let $\mathcal{B}(I, \mathcal{H})$ be the set of all Bessel sequences, $\mathcal{F}(I, \mathcal{H})$ be the collection of all frames and $\mathcal{P}(I, \mathcal{H})$ denote the set of all Parseval frames indexed by I for \mathcal{H} respectively. Then $\mathcal{B}(I, \mathcal{H})$ is a subspace of $\ell(I, \mathcal{H})$ and $\mathcal{P}(I, \mathcal{H}) \subset \mathcal{F}(I, \mathcal{H}) \subset \mathcal{B}(I, \mathcal{H}) \subset \ell(I, \mathcal{H})$.

The following theorem shows that the sum of any Bessel sequence with Bessel bound less than one with a Parseval frame is a frame for \mathcal{H} .

Theorem 2.2 Let $\mathcal{F} = \{f_i\}_{i \in I} \in \mathcal{B}(I, \mathcal{H})$ with Bessel bound $B < 1$. Then

$$\mathcal{F} + \mathcal{P}(I, \mathcal{H}) \subset \mathcal{F}(I, \mathcal{H}).$$

Proof. Let $\mathcal{G} = \{g_i\}_{i \in I} \in \mathcal{P}(I, \mathcal{H})$. Then for all $f \in \mathcal{H}$ we have

$$\begin{aligned} \|T_{\mathcal{F}}T_{\mathcal{G}}^*f\|^2 &= \sup_{\|g\|=1} |\langle T_{\mathcal{F}}T_{\mathcal{G}}^*f, g \rangle|^2 = \sup_{\|g\|=1} \left| \sum_{i \in I} \langle f, g_i \rangle \langle f_i, g \rangle \right|^2 \\ &\leq \sup_{\|g\|=1} \sum_{i \in I} |\langle f, g_i \rangle|^2 \sum_{i \in I} |\langle g, f_i \rangle|^2 \leq B\|f\|^2. \end{aligned}$$

Thus $\|T_{\mathcal{F}}T_{\mathcal{G}}^*\| \leq \sqrt{B} < 1$, and so $I_{\mathcal{H}} + T_{\mathcal{F}}T_{\mathcal{G}}^*$ an invertible operator in $\mathcal{L}(\mathcal{H})$. If we set $\Theta = (I_{\mathcal{H}} + T_{\mathcal{F}}T_{\mathcal{G}}^*)^{-1}$ and $\mathcal{U} = \mathcal{F} + \mathcal{G}$ with $\mathcal{U} = \{u_i\}_{i \in I}$. Then we compute

$$\begin{aligned} f &= (I_{\mathcal{H}} + T_{\mathcal{F}}T_{\mathcal{G}}^*)\Theta f \\ &= \sum_{i \in I} \langle \Theta f, g_i \rangle f_i + \sum_{i \in I} \langle \Theta f, g_i \rangle g_i \\ &= \sum_{i \in I} \langle f, \Theta^* g_i \rangle u_i, \end{aligned}$$

for all $f \in \mathcal{H}$. This shows that $\mathcal{U} \in \mathcal{F}(I, \mathcal{H})$ with frame bounds $\|\Theta\|^{-2}$ and $(1 + \sqrt{B})^2$. ■

The next result shows that a Bessel sequence is an inner summand of a frame.

Corollary 2.3 Let $\mathcal{F} \in \mathcal{B}(I, \mathcal{H})$ be a Bessel sequence. Then there exists a tight frame $\mathcal{G} \in \mathcal{F}(I, \mathcal{H})$ such that $\mathcal{F} + \mathcal{G} \in \mathcal{F}(I, \mathcal{H})$.

Proof. Let B be the Bessel bound for $\mathcal{F} = \{f_i\}_{i \in I}$, then $\{\frac{1}{\sqrt{2B}}f_i\}_{i \in I}$ is a Bessel sequence with the Bessel bound less than one. By Theorem 2.2 $\{\frac{1}{\sqrt{2B}}f_i + e_i\}_{i \in I}$ is a frame for \mathcal{H} , where $\{e_i\}_{i \in I}$ is an arbitrary Parseval frame. Define $g_i = \sqrt{2B}e_i$ for all $i \in I$, then $\mathcal{G} = \{g_i\}_{i \in I}$ is a tight frame and $\mathcal{F} + \mathcal{G} = \{\sqrt{2B}(\frac{1}{\sqrt{2B}}f_i + e_i)\}_{i \in I}$ is also a frame for \mathcal{H} . ■

The next theorem changes every Bessel sequence to a dual frame by summing it with any Parseval frame.

Theorem 2.4 Let $\mathcal{F} \in \mathcal{B}(I, \mathcal{H})$ with Bessel bound $B < 1$ and let $\mathcal{E} \in \mathcal{P}(I, \mathcal{H})$. Then there exists a $\mathcal{G} \in \mathcal{B}(I, \mathcal{H})$ such that $\mathcal{F} + \mathcal{E}$ and $\mathcal{G} + \mathcal{E}$ are dual frames.

Proof. Let $\mathcal{F} = \{f_i\}_{i \in I}$ and $\mathcal{E} = \{e_i\}_{i \in I}$. Since $B < 1$, hence $I_{\mathcal{H}} + T_{\mathcal{F}}T_{\mathcal{E}}^*$ is an invertible operator in $\mathcal{L}(\mathcal{H})$. If we define $\Theta = -(I_{\mathcal{H}} + T_{\mathcal{F}}T_{\mathcal{E}}^*)^{-1}T_{\mathcal{F}}T_{\mathcal{E}}^*$ and $g_i = \Theta^*e_i$ for all $i \in I$. Then $\mathcal{G} = \{g_i\}_{i \in I}$ is a Bessel sequence for \mathcal{H} and for all $f \in \mathcal{H}$ we have

$$\begin{aligned} f &= (I_{\mathcal{H}} + T_{\mathcal{F}}T_{\mathcal{E}}^*)\Theta f + T_{\mathcal{E}}T_{\mathcal{E}}^*f + T_{\mathcal{F}}T_{\mathcal{E}}^*f \\ &= T_{\mathcal{E}}T_{\mathcal{E}}^*\Theta f + T_{\mathcal{E}}T_{\mathcal{E}}^*f + T_{\mathcal{F}}T_{\mathcal{E}}^*\Theta f + T_{\mathcal{F}}T_{\mathcal{E}}^*f \\ &= \sum_{i \in I} \langle \Theta f, e_i \rangle e_i + \sum_{i \in I} \langle f, e_i \rangle e_i + \sum_{i \in I} \langle \Theta f, e_i \rangle f_i + \sum_{i \in I} \langle f, e_i \rangle f_i \\ &= \sum_{i \in I} \langle f, g_i + e_i \rangle (f_i + e_i), \end{aligned}$$

which this finishes the proof. ■

Corollary 2.5 Let $\mathcal{F} \in \mathcal{B}(I, \mathcal{H})$ with Bessel bound $B < 1$ and let $\mathcal{E} \in \mathcal{P}(I, \mathcal{H})$. Then there exists a $\mathcal{G} \in \mathcal{B}(I, \mathcal{H})$ such that $\mathcal{F} + \mathcal{E}$ and \mathcal{G} are dual frames.

Proof. Suppose that $\mathcal{F} = \{f_i\}_{i \in I}$ and $\mathcal{E} = \{e_i\}_{i \in I}$. Since $I_{\mathcal{H}} + T_{\mathcal{F}}T_{\mathcal{E}}^*$ is invertible on \mathcal{H} . Thus if we set $\Theta = (I_{\mathcal{H}} + T_{\mathcal{F}}T_{\mathcal{E}}^*)^{-1}$ and $g_i = \Theta^*e_i$ for all $i \in I$. Then for all $f \in \mathcal{H}$ we have

$$\begin{aligned} f &= (I_{\mathcal{H}} + T_{\mathcal{F}}T_{\mathcal{E}}^*)\Theta f = T_{\mathcal{E}}T_{\mathcal{E}}^*\Theta f + T_{\mathcal{F}}T_{\mathcal{E}}^*\Theta f \\ &= \sum_{i \in I} \langle \Theta f, e_i \rangle e_i + \sum_{i \in I} \langle \Theta f, e_i \rangle f_i = \sum_{i \in I} \langle f, g_i \rangle (f_i + e_i). \end{aligned}$$

From this completes the proof. \blacksquare

Corollary 2.6 For every $\mathcal{F} \in \mathcal{B}(I, \mathcal{H})$ there exist $\mathcal{G} \in \mathcal{B}(I, \mathcal{H})$ and a tight frame $\mathcal{U} \in \mathcal{F}(I, \mathcal{H})$ such that $\mathcal{F} + \mathcal{U}$ and \mathcal{G} are dual frames for \mathcal{H} .

Proof. Let B be the Bessel bound of $\mathcal{F} = \{f_i\}_{i \in I}$ and let $\{e_i\}_{i \in I}$ denote any Parseval frame for \mathcal{H} . By Theorem 2.4 there exists a Bessel sequence $\{v_i\}_{i \in I}$ for \mathcal{H} such that $\{\frac{1}{\sqrt{2B}}f_i + e_i\}_{i \in I}$ and $\{v_i + e_i\}_{i \in I}$ are dual frames for \mathcal{H} . Put $\mathcal{G} = \{g_i\}_{i \in I}$, $\mathcal{U} = \{u_i\}_{i \in I}$ with $g_i = \frac{1}{\sqrt{2B}}v_i + \frac{1}{\sqrt{2B}}e_i$ and $u_i = \sqrt{2B}e_i$ for all $i \in I$. Then for all $f \in \mathcal{H}$ we have

$$\begin{aligned} \sum_{i \in I} \langle f, g_i \rangle (f_i + u_i) &= \sum_{i \in I} \langle f, \frac{1}{\sqrt{2B}}v_i + \frac{1}{\sqrt{2B}}e_i \rangle (f_i + \sqrt{2B}e_i) \\ &= \sum_{i \in I} \langle f, v_i + e_i \rangle (\frac{1}{\sqrt{2B}}f_i + e_i) = f. \end{aligned}$$

From this the claim follows immediately. \blacksquare

In the following theorem we show that every Bessel sequence can be expanded to a dual frame by adding it to a Parseval frame. Another form of this result can be found in [5] Corollary 3.2.

Theorem 2.7 Let $\mathcal{F} = \{f_i\}_{i \in I}$ be a Bessel sequence with Bessel bound B and $\mathcal{E} = \{e_i\}_{i \in I}$ be a Parseval frame for \mathcal{H} . Then for all $\alpha > B$, there exists a Bessel sequence $\{g_i\}_{i \in I}$ for \mathcal{H} such that $\{f_i, e_i\}_{i \in I}$ and $\{\frac{1}{\alpha}f_i, g_i\}_{i \in I}$ are dual frames for \mathcal{H} .

Proof. Since $\alpha > B$, hence $\Theta = I_{\mathcal{H}} - \frac{1}{\alpha}T_{\mathcal{F}}T_{\mathcal{F}}^*$ is a linear bounded and positive operator on \mathcal{H} . Thus if we define $g_i = \Theta^*e_i$ for all $i \in I$. Then $\{g_i\}_{i \in I}$ is a Bessel sequence for \mathcal{H} and for all $f \in \mathcal{H}$ we have

$$\begin{aligned} \sum_{i \in I} \langle f, \frac{1}{\alpha}f_i \rangle f_i + \sum_{i \in I} \langle f, g_i \rangle e_i &= \sum_{i \in I} \langle f, \frac{1}{\alpha}f_i \rangle f_i + \sum_{i \in I} \langle \Theta f, e_i \rangle e_i \\ &= \frac{1}{\alpha}T_{\mathcal{F}}T_{\mathcal{F}}^*f + \Theta f = f \end{aligned}$$

which this finishes the proof. \blacksquare

Let $\Lambda = \{\Lambda_i\}_{i \in I}$ and $\Gamma = \{\Gamma_i\}_{i \in I}$ be g -Bessel sequences for \mathcal{H} with synthesis operators T_{Λ} and T_{Γ} respectively. Then we say that Λ and Γ are dual g -frames for \mathcal{H} if $T_{\Lambda}T_{\Gamma}^* = I_{\mathcal{H}}$ or $T_{\Gamma}T_{\Lambda}^* = I_{\mathcal{H}}$. In the following we show that any pair of g -Bessel sequences can be extended to pair of dual g -frames. This result, generalizes a result of Christensen, Oh Kim and Young Kim [2] to the situation of g -frames.

Theorem 2.8 Let $\Lambda = \{\Lambda_i\}_{i \in I}$ and $\Gamma = \{\Gamma_i\}_{i \in I}$ be two g -Bessel sequences for \mathcal{H} with respect to $\{W_i\}_{i \in I}$. Then there exist g -Bessel sequences $\{\Xi_j\}_{j \in J}$ and $\{\Omega_j\}_{j \in J}$ for \mathcal{H} with respect to $\{V_j\}_{j \in J}$, such that $\{\Lambda_i\}_{i \in I} \cup \{\Xi_j\}_{j \in J}$ and $\{\Gamma_i\}_{i \in I} \cup \{\Omega_j\}_{j \in J}$ form a pair of dual g -frames for \mathcal{H} with respect to $\{W_i\}_{i \in I} \cup \{V_j\}_{j \in J}$.

Proof. Assume that $\{\Phi_j\}_{j \in J}$ and $\{\Psi_j\}_{j \in J}$ are any pair of dual g -frames for \mathcal{H} respect to $\{V_j\}_{j \in J}$ and let $\Theta = I_{\mathcal{H}} - T_{\Gamma}T_{\Lambda}^*$. Then for any $f \in \mathcal{H}$ we have

$$f = \Theta f + T_{\Gamma}T_{\Lambda}^*f = \sum_{j \in J} \Psi_j^* \Phi_j \Theta f + \sum_{i \in I} \Gamma_i^* \Lambda_i f.$$

If we set $\Xi_j = \Phi_j \Theta$ and $\Omega_j = \Psi_j$ for all $j \in J$. Then $\{\Lambda_i\}_{i \in I} \cup \{\Xi_j\}_{j \in J}$ and $\{\Gamma_i\}_{i \in I} \cup \{\Omega_j\}_{j \in J}$ are dual g -frames for \mathcal{H} with respect to $\{W_i\}_{i \in I} \cup \{V_j\}_{j \in J}$. ■

The following corollaries are generalizations of the above results to the g -frames situation. We leave the proofs to interested readers.

Corollary 2.9 Let $\Lambda = \{\Lambda_i\}_{i \in I}$ be a g -Bessel sequence for \mathcal{H} with respect to $\{W_i\}_{i \in I}$ with g -Bessel bound $B < 1$. Then there exists g -Bessel sequence $\{\Gamma_i\}_{i \in I}$ for \mathcal{H} with respect to $\{W_i\}_{i \in I}$, such that $\{\Xi_i + \Lambda_i\}_{i \in I}$ and $\{\Xi_i + \Gamma_i\}_{i \in I}$ are dual g -frames for \mathcal{H} with respect to $\{W_i\}_{i \in I}$, where $\{\Xi_i\}_{i \in I}$ is a Parseval g -frame for \mathcal{H} with respect to $\{W_i\}_{i \in I}$.

Corollary 2.10 For every g -Bessel sequence $\Lambda = \{\Lambda_i\}_{i \in I}$ with Bessel bound $B < 1$ and each Parseval g -frame $\Xi = \{\Xi_i\}_{i \in I}$ for \mathcal{H} with respect to $\{W_i\}_{i \in I}$, there exists g -Bessel sequence $\{\Gamma_i\}_{i \in I}$ for \mathcal{H} with respect to $\{W_i\}_{i \in I}$ such that $\{\Lambda_i + \Xi_i\}_{i \in I}$ and $\{\Gamma_i\}_{i \in I}$ are dual g -frames for \mathcal{H} with respect to $\{W_i\}_{i \in I}$.

Corollary 2.11 For every g -Bessel sequence $\{\Lambda_i\}_{i \in I}$ for \mathcal{H} with respect to $\{W_i\}_{i \in I}$ there exist g -Bessel sequence $\{\Gamma_i\}_{i \in I}$ and a tight g -frame $\{\Xi_i\}_{i \in I}$ for \mathcal{H} with respect to $\{W_i\}_{i \in I}$ such that $\{\Lambda_i + \Xi_i\}_{i \in I}$ and $\{\Gamma_i\}_{i \in I}$ are dual g -frames for \mathcal{H} with respect to $\{W_i\}_{i \in I}$.

3. Conclusions

In this paper, using of g -frame algorithm we obtain an approximation of a bounded linear operator $U(\mathcal{H}, \mathcal{K})$. We also show that the sum of any Bessel sequence with Bessel bound less than one with a Parseval frame is a frame for \mathcal{H} and we prove that a Bessel sequence is an inner summand of a frame. The important result of this paper changing every Bessel sequence to a dual frame by summing it with any Parseval frame.

References

- [1] O. Christensen, An Introduction to Frames and Riesz Bases. *Birkhauser, Boston* (2003)
- [2] O. Christensen, H. O. Kim, R. Y. Kim, Extensions of Bessel sequences to dual pairs of frames. *Appl. Comput. Harmon. Anal.* **34**, (2013), 224-233.
- [3] I. Daubechies, A. Grossmann, Y. Meyer, Painless nonorthogonal expansions. *J. Math. Phys.* **27**, (1986), 1271-1283.
- [4] R. J. Duffin, A. C. Schaeffer, A class of nonharmonic Fourier series. *Trans. Amer. Math. Soc.* **72**, (2), (1952), 341-366.
- [5] D. F. Li, W. Sun, Expansion of frames to tight frames. *Acta Math. Sin. (Engl. Ser.)* **25** (2), (2009), 287292.
- [6] W. Sun, G -frames and G -Riesz bases. *J. Math. Anal. Appl.* **322**, (2006), 437-452.