

Probiotics in Goat Milk: A Promising Solution for Management of Drug-Resistant *Acinetobacter baumannii*

Leila Fozouni^{1*}, Maryam Yaghoobpour², Ania Ahani³

1. Department of Biology, Gorgan Branch, Islamic Azad University, Gorgan, Iran.
2. Department of Biology, Gorgan Branch, Islamic Azad University, Gorgan, Iran.
3. Department of Biology, Gorgan Branch, Islamic Azad University, Gorgan, Iran.

Article Type:

Original Article

Article History:

Received: 4 Aug 2019

Revised: 1 Sep 2019

Accepted: 30 Sep 2019

*Correspondence:

Leila Fozouni

Department of Biology,
Gorgan Branch, Islamic
Azad University, Gorgan,
Iran

lili_kia@yahoo.com



Abstract

Background and objectives: *Acinetobacter* is a genus of opportunistic pathogens that are commonly found in the environment. Given the unique ability of these bacteria to survive in the hospital, they are considered as one of the main causes of hospital-acquired infections. The emergence of multidrug-resistant *Acinetobacter* spp., particularly *Acinetobacter baumannii* has become a major health threat worldwide. In this study, we investigate antibacterial effects of probiotic isolates from goat milk on clinical isolates of *A. baumannii*.

Methods: In this study, 100 clinical specimens were taken from patients hospitalized in six hospitals in the Golestan Province, north of Iran. Following isolation and identification of *A. baumannii* strains, antibiotic resistance patterns of the isolates were investigated using the Kirby-Bauer method according to the Clinical and Laboratory Standards Institute (CLSI-2015) guidelines. Probiotic bacteria in goat milk were isolated and identified by culture in MRS and M17 media and carbohydrate fermentation tests. Antibacterial effects of the probiotic bacteria against resistant *A. baumannii* isolates were evaluated using the agar well diffusion method.

Results: Overall, 55% of the isolates were identified as *A. baumannii*. The highest resistance rates were observed against tobramycin (76.3%), mezlocillin (74.5%) and cefotaxime (74.5%). Resistance to levofloxacin, tetracycline, imipenem and minocycline was detected in 72.7%, 72.7%, 70.9% and 29.1% of the isolates, respectively. The most common probiotic isolates were *Lactobacillus plantarum* and *Lactococcus piscium* (30% each). The highest and lowest effects were exerted by *Lactococcus lactis* (34.54%) and *Lactobacillus bulgaricus* (3.63%), respectively.

Conclusion: Our results demonstrate that the prevalence of drug-resistant *A. baumannii* strains is high in the hospitals. Given the promising antimicrobial effects of the isolated probiotic bacteria, goat milk can be recommended as an adjuvant therapy or an alternative to common antibiotics for improving treatment outcome of infections caused by drug-resistant *A. baumannii*.

Keywords: *Acinetobacter*, Goat milk, Probiotic, Nosocomial infection

Introduction

Emergence of antibiotic-resistant microorganisms that cause nosocomial infections has become a major healthcare challenge (1). *Acinetobacter* is a genus of gram-negative coccobacilli from the Moraxellaceae family. *Acinetobacter* species, including *Acinetobacter baumannii*, are opportunistic pathogens that can cause both community-acquired and nosocomial infections, especially in intensive care unit and high-dependency unit patients (2,3). These bacteria have been isolated from various infections including ventilator-associated pneumonia, endocarditis, meningitis, skin and soft-tissue infections, urinary tract infection and prosthesis-related infections. *A. baumannii* has been isolated from diverse animal, human and environmental sources, but the control of its presence in healthcare settings is most importance (4,5). Until three decades ago, *A. baumannii* infections were treated effectively with conventional antibiotics (6,7), but, the recent emergence of multi-drug resistant *A. baumannii* strains has made them difficult to control (8). Therefore, much attention has been paid to discovery of novel, effective and non-toxic alternatives to conventional antibiotics. Probiotics are an example of such compounds that lack the side effects of antibiotics. Probiotics create holes in the phospholipid bilayer of bacteria by disrupting the cytoplasmic membrane and generating the proton motive force. Also, they prevent the probable growth and metastasis of bacteria through immunologic and non-immunologic mechanisms (9,10). The purpose of this study was to investigate in vitro antibacterial activity of goat milk as probiotics source against *A. baumannii*.

Materials and Methods

Sampling and bacterial isolation

This descriptive study was performed on 100 samples (blood, burn wound and trachea) from six hospitals in the Golestan Province, Iran. Before sampling, written informed consent was taken from all patients. The samples were cultured on blood agar and MacConkey agar. *A. baumannii* strains were identified by colony morphology, gram staining and biochemical tests including oxidase, catalase, citrate, motion, TSI, indole, methyl red, Voges-Proskauer and carbohydrate fermentation test.

Antimicrobial susceptibility testing

Antimicrobial susceptibility was evaluated using the disc diffusion method (Kirby-Bauer method). First, a suspension (equivalent to a 0.5 McFarland standard) was prepared from an overnight culture of *Acinetobacter* spp. isolates. After incubation at 37°C, the isolates were cultured on Mueller Hinton agar (Merck, Germany) using sterile swabs. Levofloxacin (5 µg), minocycline (30 µg), cefotaxime (30 µg), tobramycin (10 µg), mezlocillin (75 µg), imipenem (10 µg) and tetracycline (30 µg) disks were purchased from Padtan Teb Co. (Tehran, Iran). The antibiotics disks were placed on the culture medium. After 18-24 hours of incubation at 37 °C, results were interpreted as resistant, intermediate and susceptible by measuring the diameter of inhibition zone according to the Clinical and Laboratory Standards Institute guidelines (CLSI-M100-S25) (11).

Isolation of probiotic bacteria

Milk samples were collected from 1-3 years old Pakistani-Turkmen goats. The milk samples were cultured on MRS agar and M17 agar (Merck, Germany) and incubated at 37 °C

for 48 hours under anaerobic conditions. Bacterial species were identified based on colony morphology, gram staining, catalase test and ability to ferment cellobiose, fructose, galactose, glucose, lactose, maltose, mannitol, mannose, melezitose, ribose and trehalose.

In order to isolate antimicrobials, several colonies of lactic bacteria isolated from goat milk were inoculated into tubes containing M17 broth and MRS broth. After addition of paraffin to the media, the tubes were incubated at 37 °C for 4 days. Next, paraffin was removed and the tubes were centrifuged at 3500 rpm for 10 minutes. Sediment was removed under sterile conditions and supernatant containing bacterial metabolites was stored for future use.

Evaluation of antimicrobial activity of probiotic isolates

The agar well diffusion method was used to evaluate antimicrobial activity of the probiotic isolates. First, a suspension (equivalent to a 0.5 McFarland standard; 1.5×10^8 CFU/mL) from all antibiotic-resistant *A. baumannii* isolates was prepared in physiological saline solution and then cultured on Muller Hinton agar. Next, wells (7mm diameter) were created on the culture medium and 100 μ l of the bacterial suspension were added to each well. A well containing distilled water was considered as negative control. The plates were incubated at 37 °C for 24 hours. A growth inhibition zone diameter of ≥ 15 and ≤ 12 mm indicated susceptibility and resistance, respectively. Moreover, *A. baumannii* ATCC 19606 was used as a positive control.

Results

Overall, 55% of the isolates were identified as *A. baumannii*. The frequency of *A. baumannii* isolates was highest in men (76.4%) and patients aged 50 years and older (60%). The rate of antibiotic resistance among *A. baumannii* was highest against tobramycin (76.3%) and lowest against minocycline (29.1%) (Figure 1).





Figure 1. Relative frequency of antibiotic resistance among *A. baumannii* isolates

According to the results of the carbohydrate fermentation tests, *Lactobacillus plantarum* and *Lactococcus piscium* were identified as the probiotic strains in milk samples from three-year-old goats. *Lactobacillus bulgaricus* and *Lactococcus lactis* were identified as the probiotic strains in milk samples from two-year-old goats. *L. plantarum*, *Lactococcus raffinolactis* and *L. piscium* were identified as the probiotic strains in milk samples from one-year-old goats (Figure 2).



According to results of the agar well diffusion assay, *L. lactis* exerted the strongest antibacterial activity against *A. baumannii* isolates (Table 1).

Figure 2. Relative abundance of the probiotic strains isolated from goat milk.

Table 1. Antibacterial activity of the probiotic isolates from goat milk against drug-resistant *A. baumannii*

	<i>L. lactis</i>		<i>L. piscium</i>		<i>L. raffinolactis</i>		<i>L. plantarum</i>		<i>L. bulgaricus</i>	
	No	%	No	%	No	%	No	%	No	%
Grown	36	65.45	44	80	49	89.09	46	83.63	53	96.36
Inhibited	19	35.45	11	20	6	10.90	9	16.36	2	3.63
Total	55	100	55	100	55	100	55	100	55	100
P-Value	0.01		0.01		0.04		0.036		0.05	

Table 2. Mean diameter of growth inhibition zones caused by probiotic isolates around drug-Resistant *A. baumannii*^{a,b}

Probiotics	Mean ±SD
<i>L. bulgaricus</i>	15.5±4.01
<i>L. plantarum</i>	16.3±3.90
<i>L. lactis</i>	18.6±0.03
<i>L. piscium</i>	17.4±0.00
<i>L. raffinolactis</i>	15.8±0.00
^a Diameter of the Inhibition Zone, mm.	
^b P value < 0.01.	

Discussion

Management of nosocomial infections caused by *Acinetobacter* species has become challenging due to the recent emergence and spread of drug-resistant strains. *A. baumannii* is one of the most clinically important *Acinetobacter* species that can cause a wide range of infections (12). The purpose of our study was to evaluate the antagonistic activity of probiotic isolates from goat milk against drug-resistant *A. baumannii* isolates. Of 100 clinical specimens collected from patients in intensive care and high dependency units, 55 were identified as *A. baumannii*. In the antibiotic susceptibility test, we observed high resistance rates against tobramycin (76.3%), mezlocillin (74.5%), cefotaxime (74.5%), levofloxacin (72.7%), tetracycline (72.7%), imipenem (70.9%) and minocycline (29.1%).

In a previous study in Iran, the prevalence of imipenem- and ciprofloxacin-resistant *A. baumannii* isolates was 40.9% and 77.7%, respectively (13). A more recent study in Iran reported high rates of resistance against cefotaxime (93%), cefipime (91%), norfloxacin (87%), imipenem (86%) and tobramycin (67%) among *A. baumannii* clinical isolates (14). These results indicate the rising prevalence of resistant *A. baumannii* strains in recent years. According to a systematic review, the frequency of carbapenem-resistant *A. baumannii* strains increased in 2010-2013, while the rate of resistance to aminoglycosides did not change significantly (15). A 10-year survey on the prevalence of drug-resistant *A. baumannii* strains between 2004 and 2014 demonstrated a 64% increase in the frequency of imipenem-resistant strains and a 49% increase in the frequency of extensively drug-resistant strains

in 2014 (16). These findings highlight the eminent need for discovery of novel antimicrobial agents against drug-resistant bacteria. In this regard, probiotics have been suggested as promising and economically feasible alternatives to conventional antibiotics. In the present study, we isolated and identified *L. bulgaricus*, *L. plantarum*, *L. raffinolactis*, *L. piscium* and *L. lactis* from Pakistani-Turkmen goat milk samples. In 2004, a study in Algeria reported *Streptococcus thermophilus*, *Lactobacillus helveticus*, *L. plantarum*, *Lactobacillus delbrueckii* and *Lactobacillus lactis* as the most common lactic acid bacteria in raw milk samples from four goat breeds (17).

In a study in 2015, *Lactobacillus fermentum* (48%), *Lactobacillus acidophilus* (34%), *Lactobacillus viridescens* (8%), *Lactobacillus brevis* (5%) and *Lactobacillus gasseri* (4%) were detected in 40 milk samples collected from the Aarey Milk Colony in India (18). The results of these studies indicate the great diversity of probiotic bacteria in goat milk.

In the present study, *L. bulgaricus* and *L. lactis* exhibited the highest and lowest antagonistic activity against resistant *A. baumannii* isolates, respectively. Similar to our findings, a study in Mexico demonstrated the inhibitory activity of bacteriocinogenic probiotic bacteria from goat cheese against *Staphylococcus aureus*, *Bacillus cereus*, *Escherichia coli*, *Listeria monocytogenes*, *Pseudomonas aeruginosa*, *Shigella flexneri*, *Serratia marcescens*, *Enterobacter cloacae* and *Klebsiella pneumoniae* (19).

In 2004, a study confirmed antimicrobial activity of lactic acid bacteria against a standard strain of *S. aureus* (20). Among the probiotic bacteria isolated from goat milk, we

observed that *Lactococcus* spp. could exert a more promising antibacterial activity compared to *Lactobacillus* isolates against resistant *A. baumannii* strains.

Conclusion

Our findings revealed the alarmingly high prevalence of drug resistance among *A. baumannii* clinical isolates from hospitals of the Golestan Province, Iran. Given the favorable antimicrobial activity of goat milk-derived probiotics against resistant *A. baumannii* isolates, we suggest conducting future studies on the potential antimicrobial activity of these probiotics or their metabolites against other clinically important bacteria.

Funding source(s):

This research has been accomplished by the support of Microbiology lab, I.A.U, Gorgan branch and it is extracted from the MSc thesis of Maryam Yaghoobpour.

Conflict of interest:

The authors declare that there is no conflict of interest.

References

1. Curtis LT. Prevention of Hospital Acquired Re view of Non-Pharmacological Intervention. *JHosp Infect.*2008; 69(3):204-219. [DOI:10.1016/j.jhin.2008.03.018]
2. Eveillard M, Kempf M, Belmonte O, Pailhoriès H, Joly-Guillou ML. Reservoirs of *Acinetobacterbaumannii* outside the hospital and potential involvement in emerging human community-acquired infections. *Int Journal Infect Dis.*2013; 17(10) (:802-805. [DOI:10.1016/j.ijid.2013.03.021]
3. Tang SS, Apisarnthanarak A, Hsu LY. Mechanisms of B-lactam antimicrobial resistance and epidemiology of major community and healthcare-associated multidrug-resistant bacteria. *Adv DrugDelivRev.*2014; 7: 3-13. [DOI:10.1016/j.addr.2014.08.003]
4. Kim Uj, Kim Hk, An JH, Cho SK, Park Kh, Jang HC. Update on the epidemiology, treatment, and outcomes of carbapenem-resistant *Acinetobacter* infections. *chonnam Med Journal.*2014; 50(2):37-44. [DOI:10.4068/cmj.2014.50.2.37]
5. Nhu NTK, Lan NPH, Campbell JI, Parry CM, Thompson C, Tuyen HT, Howamg MVM, et al. Emergence of carbapenem-resistant *Acinetobacterbaumannii* as the major cause of ventilator-associated pneumonia in intensive care unit patients at an infectious disease hospital in southern Vietnam. *Journal Med Microbial.*2014; 63(10):1386-1394. [DOI:10.1099/jmm.0.076646-0]
6. Li J, Rayner CR, Nation RL, Owen RJ, Spelman D, Tan KE, Liolios L. Heteroresistance to colistin in multidrug-resistant *Acinetobacter baumannii*. *Antimicrob Agents Chemother.*2006; 50(9): 2946-2950. [DOI:10.1128/AAC.00103-06]
7. Garcia-Quintanilla M, Pulido MR, Lopez-Roja R, Pachon J, McConnell MJ. Emerging therapies for multidrug resistant *Acinetobacterbaumannii*. *Trends Microbial.* 2013;21(3):157-163. [DOI:10.1016/j.tim.2012.12.002]
8. Oh Y.J, Song SH, Baik SH, Lee HH, Han IM, Oh DH. A case of fulminant community-acquired *Acinetobacterbaumannii* pneumonia in Korea. *Korean Journal Intern Med.* 2013;28(4):486-490. [DOI:10.3904/kjim.2013.28.4.486]
9. Kirjavainen PV, Salminen SJ, Isolauri E. Probiotic bacteria in the management of atopic disease : underscoring the importance of viability. *Journal Pediatr Gastroenterol Nutr.*2003; 36(2): 223-227. [DOI:10.1097/00005176-200302000-00012]
10. Carvalho Eunice B, Maga Elizabeth A, Quetz Josiane S, Lima Ila FN, Magalhaes Hemerson YF, et al. Goat Milk With and Without Increased Concentrations of Lysozyme Improves Repair of Intesyinal Cell Damage Induced by Enteroaggregative *Escherichia Coli*. *BMC Gastroenterol.*2012; 12:106. [DOI:10.1186/1471-230X-12-106]
11. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fifth Informational Supplement. M100-S25. Wayne, PA.2015; 3(35):56-58.
12. Tsakris A, Poulou A, Pournaras S, Voulgari E, Vrioni G, Themeli-Digalaki K, et al. A simple phenotypic method for the differentiation of metallo-beta-lactamases and class A KPC carbapenemases in Enterobacteriaceae clinical isolates. *Journal Antimicrob Chemother.*2010; 65(8): 1664-1671. [DOI:10.1093/jac/dkq210]
13. Jafari S, Najafipour S, Kargar M, Abdollahi A, Mardaneh J, Fasihy Ramandy M, et al. Phenotypical Evaluation of Multi-Drug Resistant *Acinetobacter Baumannii*. *J Fasa Univ Med Sci.* 2013; 2 (4) :254-258.
14. Nourmohammadi Z, Zamanzad B, Shavarzi A, Kiani P. Evaluation antimicrobial resistance of *Acinetobacter baumannii* isolated from shahrekord teaching hospitals in 2013. *Journal Shahrekord Univ Med Sci.* 2015; 16(6): 1-8.

15. Moradi J, Hashemi Farhad B, Bahador A. Antibiotic Resistance of *Acinetobacter baumannii* in Iran: A Systemic Review of the Published Literature. *Osong Public Health Res Perspect.* 2015; 6(2): 79-86. [DOI:10.1016/j.phrp.2014.12.006]
16. Lei Gao, Yuan Lyu, and Yun Li. Trends in Drug Resistance of *Acinetobacter baumannii* over a 10-year period: Nationwide Data from the China Surveillance of Antimicrobial Resistance program. *Chin Med Journal.*2017; 130(6): 659-664. [DOI:10.4103/0366-6999.201601]
17. Badis A, Guetarni D, Moussa Boudjema B, Henni D.E, Kihal M. Identification and Technological properties of lactic acid bacteria isolated from raw goat milk of four Algerian races. *Food Microbiology.*2004; 21(5): 579-588. [DOI:10.1016/j.fm.2003.11.006]
18. Mithun S, Dipak V, Sheela S. Isolation and Identification of lactobacilli from raw milk samples obtained from Aarey Milk Colony. *IJSRP.*2015; 5(4): 2250-2253..
19. Oscar F. Hernández-Saldaña, Mauricio Valencia-Posadas, Norma M. de la Fuente-Salcido, Dennis K. Bideshi, and José E. Barboza-Corona. Bacteriocinogenic Bacteria Isolated from Raw Goat Milk and Cheese Produced in the Center of Mexico. *Indian J Microbiol.*2016;56(3):301-308. [DOI:10.1007/s12088-016-0587-3]
20. Savadogo A, Ouattrra CA, Bassole IHN, Traore AS. Antimicrobial activities of lactic acid bacteria strains isolated from Burkina faso fermented milk. *Pakistan Journal of nutrition.*2004; 3 (3):174-179. [DOI:10.3923/pjn.2004.174.179]

How to cite:

Fozouni L, Yaghoobpour M, Ahani A. Probiotics in Goat Milk: A Promising Solution for Management of Drug-Resistant *Acinetobacter baumannii*. *Jorjani Biomedicine Journal.* 2019; 7(2): 31-38.