

Prevalence of maxillofacial fractures in hospital patients: A five year retrospective study

Iman Shirinbak^{1,2}, Ali Baradaran Bagheri³, Mohammad Javad Kharazifard⁴, Peiman Goharshenasan⁵, Mohammad Pirouzan^{6*}

1. Department of Oral and Maxillofacial Surgery, Alborz University of Medical Sciences, Karaj, Iran
2. Dental Clinical Research Development Unit, Alborz University of Medical Sciences, Karaj, Iran
3. Department of neurosurgery, Shahid madni hospital, Alborz University of Medical Sciences, karaj, Iran
4. Department of Epidemiology and Biostatistics, Dental School, Tehran University of Medical Sciences, Tehran, Iran
- 5 Department of Surgery, Shahid Madni hospital, Alborz University of Medical Sciences, karaj, Iran
6. Student Research Committee, Alborz University of Medical Sciences, Karaj, Iran

Article Type:

Brief Report

Article History:

Received: 16 Jul 2020

Revised: 1 Aug 2020

Accepted: 5 Aug 2020

*Correspondence:

Mohammad Pirouzan,
Dental Student, Member of the
Student Research Committee
Faculty of Dentistry, Alborz
University of Medical Sciences,
Karaj, Iran

mohammadpirouzan@gmail.com



[DOI: 10.29252/jorjanibiomedj.8.3.75](https://doi.org/10.29252/jorjanibiomedj.8.3.75)

Abstract

Background and objective: Damages to the oromaxillofacial region, if not diagnosed and treated in a timely manner, will cause permanent, serious clinical problems because of the characteristics of this anatomical region. Accordingly, the present study was performed on a 5-year investigation of epidemiology of oromaxillofacial fractures in patients admitted to Shahid Madani Hospital, Karaj, Iran.

Material And Method: In this descriptive cross-sectional study, 235 medical files of patients with damages to the oromaxillofacial region available in the archive of Shahid Madani Hospital, Karaj from 2013 to 2018 were chosen as census and examined. Demographic variables including site and cause of fracture were recorded for each patient on information forms. The collected data were analyzed by SPSS 17 software and presented as descriptive statistics.

Result: In this study, out of 235 patients with oromaxillofacial fractures, 178 (75.7%) were male and 97 (41.3%) were female, respectively. The mean age of the patients was 30.96 ± 14.91 years. The main affected anatomical regions were as follows: Mandible 269 cases (49.17%), maxilla 117 cases (21.39%), and cheekbone 51 cases (9.32%). Accidents occurring with motor vehicles was the main cause of these fractures in 132 patients (56.2%).

Conclusion: The results of the present study indicated that the fractures of oromaxillofacial regions were more common in men, young people, and middle-aged individuals, and mostly occurred in the mandible, maxilla, and cheekbone, with the main cause of these fractures being accidents happening with motor vehicles.

Keywords: Facial Injuries; Jaw Fractures; Fractures, Bone; Accidents

Introduction

Maxillofacial trauma is presented as skeletal, dental, and soft tissue (1). Development of abnormalities in the growth of some jawbones, impairments in the temporomandibular (TMJ) joint and occlusion, displacement and degradation of bones, as well as oral deformity plus loss of teeth are among the complications of oromaxillofacial fractures (2). These fractures lead to incidence of severe complications, facial deformity, as well as problems in oral functions (3). The psychological aspects of damage to the oromaxillofacial region are also very important, since facial fractures often cause impaired aesthetics in the injured person (4).

Studies have shown that age and gender are important factors affecting incidence of oromaxillofacial trauma (5-7). The greatest incidence has been observed in the second decade of life, while the lowest has been seen in patients below five years of age and above sixty. The male to female ratio in the world has been reported at 4:1 (8).

The main cause of oromaxillofacial fractures worldwide is accidents with motor vehicles, falling, personal fights, and damages resulting from firearms, sports, and occupational accidents (9). These causes may differ given the geographical region, socioeconomic status, as well as cultural characteristics (10). In developing countries, oromaxillofacial fractures mostly occur in response to accidents with motor vehicles (11).

Researchers have examined the incidence and prevalence of oromaxillofacial fractures across different provinces and regions of Iran (12-15). However, since there is no adequate information about the epidemiology of these fractures in Alborz Province, the present

research was performed to investigate the frequency of oromaxillofacial fractures in patients hospitalized in Shahid Madani Hospital in Karaj.

Materials and Methods

In this descriptive cross-sectional study, which was performed retrospectively, 749 medical files as well as radiological documents, CT scans, and 3D facial reconstructions related to patients with oromaxillofacial damages available in the archive of Shahid Madani Hospital in Karaj, from 2013 to 2018, were chosen through census method and examined after approval by the Ethics Committee of Alborz University of Medical Sciences, code: IR.ABZUMS.REC.1398.104, also receiving an introduction letter from the head of the dentistry faculty, and receiving permission from the head of the Hospital.

Demographic information was recorded and kept confidential on the information form for every patient, such as age and gender, site of fracture; mandible (condyle, ramus, body, and symphysis), middle face (maxilla, cheekbone, nose, frontal sinus), and dentoalveolar, as well as cause of fracture; accident with motor vehicle, falling, intimate partner violence, sporting event, occupational event, and stumbling. The collected data were analyzed by SPSS 17 software and presented as descriptive statistics (Tables and Figures).

Result

In the present study, 235 patients with oromaxillofacial fracture were studied. Out of the 235 patients, 178 (75.7%) were male and 97 (41.3%) were female. The fracture ratio was 1.8 times higher in men than in women. Most of the injured individuals were young people (54%). The mean age of the patients was 30.96 ± 14.91 years. The minimum and

maximum age of patients was 1 and 90 years old, respectively (Table 1). The total number of fractures was 547. The results of the study indicated that the main fractured oromaxillofacial anatomical regions were mandible (269 cases), maxilla (117 cases), and cheekbone (51 cases). The highest mandibular fracture was observed in the body (48.9) and the lowest in the ramus (11.5) (Table 2).

The results of investigating the causes of fractures in the study patients suggested that in 56.2% of the cases (n = 132) accidents with motor vehicles was the main cause of fractures in the oromaxillofacial region, followed by other accidents including falling from heights (17/4%) and physical disputes (14%) (Figure 1).

Table 1. Frequency of maxillofacial fractures in patients' age and gender

Fractures		Yes n (%)	No n (%)
Mandible	condyle	52(22.1)	183(77.9)
	ramus	27(11.5)	208(88.5)
	Body	115(48.9)	120(51.1)
	symphysis	75(31.9)	160(68.1)
middle face	maxilla	117(49.8)	118(50.2)
	nose	34(14.5)	201(85.5)
	cheek	51(21.7)	184(78.3)
	frontal sinus	30(12.8)	105(87.2)
dentoalveolar		46(19.6)	189(80.4)

Table 2. Frequency of maxillofacial fractures according to anatomic site

demographic data		N	%
Age groups	Child (1-11 years)	17	7.2
	Adolescents (17-12 years)	14	6
	Young (34-18 years)	127	54
	Middle-aged (64-35 years)	73	31.1
	Older (over 65 years old)	4	1.7
gender	Female	97	41.3
	male	178	75.7




Fig 1. Frequency of maxillofacial fractures in patients according to the causes of fractures

Discussion

In this study, the research sample was chosen from patients hospitalized in one of the Hospital in Karaj, which is one of the important health care centers for different types of oromaxillofacial region fractures in the province, whereby a considerable number of people with such fractures are treated by the experienced health care team of this center.

The results of the present study indicated that the oromaxillofacial region fractures occur twice more frequently in men than in women (2:1 ratio), almost concurring with the results of Lee et al. (16) in Korea and Ellis et al. (17) in Scotland. This figure is low compared to similar studies including those of Kadkhodaie (13) in Rasht (1:12), Al Ahmed et al. (4) in UAE (1:11), and Adekeye (18) in Nigeria (1:16.9).

The higher prevalence of oromaxillofacial injuries in men in the present study has been confirmed in various studies (13, 19-22). This is possibly due to the fact that in many families in Iran men mostly work outside home for generating income for the family, which in turn increases the risk of accidents or violent conflicts (3). In addition, men in Iran mostly drive more than women do, and also participate in battle-involving sports such as soccer and basketball. These men are also more likely than women to consume alcohol and drugs before driving, which all increase the risk of oromaxillofacial fractures (23).

In the present study, most patients (42.3%) were young which is in line with similar studies (1, 4, 12, 14, 24-31). This is possibly due to behavioral, social, economic changes as well as emotional conflicts, mostly occurring in this age group. At these ages, the youth are in the stage of personality independence, social excitement, high

activity, recklessness in driving, and participation in physical conflicts (16, 22, 25, 28).

In the present study, the main fractured oromaxillofacial anatomical regions were the mandible, maxilla, and cheekbone. The fractures of facial bones (especially the mandible because of its prognathism in the face) are among the common injuries in patients admitted to hospital emergencies rooms (32). The incidence of maxillofacial fractures in the mandible, in this study, concurred with most similar studies (14, 24, 27, 28, 30, 31, 33, 34). The damages to oromaxillofacial region are among the most common injuries caused by accidents (35). In the present study, accidents with motor vehicles (58.8%) was the most common cause of fracture, which is in line with the results of other studies conducted in Iran (12, 14, 15, 36) and in other countries (29, 37, 38). However, the studies performed in Finland (39) and Australia (20) indicated that violent conflicts and daily activities are the most important causes of maxillofacial trauma. The large difference between the frequencies of fracture inducing accidents, in this study, compared to other causes can be due to poorer adherence to safety principles such as neglecting to use safety belts or helmets by drivers, and not taking driving rules seriously. Observing safety principles can considerably reduce the incidence of accidents (16). Incompleteness of some medical files of patients was the main limitation.

Conclusion

The results of the present study indicated that maxillofacial fractures are mostly common in men and the youth, and mostly occur in the mandible, maxilla, and cheekbone, with the main cause of these fractures being accidents. Planners, policymakers, private and

governmental companies should take measures to correct the infrastructures, promote rules and regulations, and train citizens by way of more educational programs and media advertisements in order to reduce the rate of accidents and their undesired consequences.

References

1. Bakardjiev A, Pechalova P. Maxillofacial fractures in Southern Bulgaria - a retrospective study of 1706 cases. *J Craniomaxillofac Surg* 2007;35(3):147-50. [DOI:10.1016/j.jcms.2007.01.005]
2. Anbiaee N, Ahmadian-Yazdi A, Bagherpour A, Ghaziani M. Two Year Evaluation of Maxillofacial Fractures in Conventional Radiographs of Patients Referring to Radiology Department of Mashhad Dental School. *J Mashhad Dent Sch* 2014;38(1):1-8.
3. Pandey S, Roychoudhury A, Bhutia O, Singhal M, Sagar S, Pandey RM. Study of the pattern of maxillofacial fractures seen at a tertiary care hospital in north India. *J Maxillofac Oral Surg* 2015;14(1):32-9. [DOI:10.1007/s12663-013-0578-4]
4. Al Ahmed HE, Jaber MA, Abu Fanas SH, Karas M. The pattern of maxillofacial fractures in Sharjah, United Arab Emirates: a review of 230 cases. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2004;98(2):166-70. [DOI:10.1016/j.tripleo.2004.01.020]
5. Elarabi MS, Bataineh AB. Changing pattern and etiology of maxillofacial fractures during the civil uprising in Western Libya. *Med Oral Patol Oral Cir Bucal* 2018;23(2):e248-e255. [DOI:10.4317/medoral.22268]
6. van den Bergh B, Karagozolu KH, Heymans MW, Forouzanfar T. Aetiology and

incidence of maxillofacial trauma in Amsterdam: a retrospective analysis of 579 patients. *J Craniomaxillofac Surg* 2012;40(6):e165-9.

[DOI:10.1016/j.jcms.2011.08.006]

7. Zhou HH, Liu Q, Yang RT, Li Z, Li ZB. Maxillofacial Fractures in Women and Men: A 10-Year Retrospective Study. *J Oral Maxillofac Surg* 2015;73(11):2181-8.

[DOI:10.1016/j.joms.2015.07.022]

8. Zhou HH, Ongodia D, Liu Q, Yang RT, Li ZB. Changing pattern in the characteristics of maxillofacial fractures. *J Craniofac Surg* 2013;24(3):929-33.

[DOI:10.1097/SCS.0b013e3182587f86]

9. Bali R, Sharma P, Garg A, Dhillon G. A comprehensive study on maxillofacial trauma conducted in Yamunanagar, India. *J Inj Violence Res* 2013;5(2):108-16.

[DOI:10.5249/jivr.v5i2.331]

10. Bataineh AB. Etiology and incidence of maxillofacial fractures in the north of Jordan. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1998;86(1):31-5.

[DOI:10.1016/S1079-2104(98)90146-9]

11. Khalil AF, Shaladi OA. Fractures of the facial bones in the eastern region of Libya. *Br J Oral Surg* 1981;19(4):300-4.

[DOI:10.1016/0007-117X(81)90051-2]

12. Ansari MH. Maxillofacial fractures in Hamedan province, Iran: a retrospective study (1987-2001). *J Craniomaxillofac Surg* 2004;32(1):28-34.

[DOI:10.1016/j.jcms.2003.07.010]

13. Kadkhodaie MH. Three-year review of facial fractures at a teaching hospital in northern Iran. *Br J Oral Maxillofac Surg* 2006;44(3):229-31.

[DOI:10.1016/j.bjoms.2005.06.016]

14. Motamedi MH. An assessment of maxillofacial fractures: a 5-year study of 237 patients. *J Oral Maxillofac Surg* 2003;61(1):61-4. [DOI:10.1053/joms.2003.50049]
15. Zargar M, Khaji A, Karbakhsh M, Zarei MR. Epidemiology study of facial injuries during a 13 month of trauma registry in Tehran. *Indian J Med Sci* 2004;58(3):109-14.
16. Lee JH, Cho BK, Park WJ. A 4-year retrospective study of facial fractures on Jeju, Korea. *J Craniomaxillofac Surg* 2010;38(3):192-6. [DOI:10.1016/j.jcms.2009.06.002]
17. Ellis E, 3rd, Moos KF, el-Attar A. Ten years of mandibular fractures: an analysis of 2,137 cases. *Oral Surg Oral Med Oral Pathol* 1985;59(2):120-9. [DOI:10.1016/0030-4220(85)90002-7]
18. Adekeye EO. The pattern of fractures of the facial skeleton in Kaduna, Nigeria. A survey of 1,447 cases. *Oral Surg Oral Med Oral Pathol* 1980;49(6):491-5. [DOI:10.1016/0030-4220(80)90068-7]
19. Aksoy E, Unlu E, Sensoz O. A retrospective study on epidemiology and treatment of maxillofacial fractures. *J Craniofac Surg* 2002;13(6):772-5. [DOI:10.1097/00001665-200211000-00012]
20. Gassner R, Tuli T, Hachl O, Rudisch A, Ulmer H. Cranio-maxillofacial trauma: a 10 year review of 9,543 cases with 21,067 injuries. *J Craniomaxillofac Surg* 2003;31(1):51-61. [DOI:10.1016/S1010-5182(02)00168-3]
21. Kieser J, Stephenson S, Liston PN, Tong DC, Langley JD. Serious facial fractures in New Zealand from 1979 to 1998. *Int J Oral Maxillofac Surg* 2002;31(2):206-9. [DOI:10.1054/ijom.2002.0208]
22. Lone P, Singh AP, Kour I, Kumar M. A 2-year retrospective analysis of facial injuries in patients treated at department of oral and maxillofacial surgery, IGGDC, Jammu, India. *Natl J Maxillofac Surg* 2014;5(2):149-52. [DOI:10.4103/0975-5950.154817]
23. Ungari C, Filiaci F, Riccardi E, Rinna C, Iannetti G. Etiology and incidence of zygomatic fracture: a retrospective study related to a series of 642 patients. *Eur Rev Med Pharmacol Sci* 2012;16(11):1559-62.
24. Brasileiro BF, Passeri LA. Epidemiological analysis of maxillofacial fractures in Brazil: a 5-year prospective study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2006;102(1):28-34. [DOI:10.1016/j.tripleo.2005.07.023]
25. Chandra Shekar BR, Reddy C. A five-year retrospective statistical analysis of maxillofacial injuries in patients admitted and treated at two hospitals of Mysore city. *Indian J Dent Res* 2008;19(4):304-8. [DOI:10.4103/0970-9290.44532]
26. Gomes PP, Passeri LA, Barbosa JR. A 5-year retrospective study of zygomatico-orbital complex and zygomatic arch fractures in Sao Paulo State, Brazil. *J Oral Maxillofac Surg* 2006;64(1):63-7. [DOI:10.1016/j.joms.2005.09.012]
27. Kamulegeya A, Lakor F, Kabenge K. Oral maxillofacial fractures seen at a Ugandan tertiary hospital: a six-month prospective study. *Clinics (Sao Paulo)* 2009;64(9):843-8. [DOI:10.1590/S1807-59322009000900004]
28. Paes JV, de Sa Paes FL, Valiati R, de Oliveira MG, Pagnoncelli RM. Retrospective study of prevalence of face fractures in southern Brazil. *Indian J Dent Res* 2012;23(1):80-6. [DOI:10.4103/0970-9290.99045]

29. Ugboko VI, Odusanya SA, Fagade OO. Maxillofacial fractures in a semi-urban Nigerian teaching hospital. A review of 442 cases. *Int J Oral Maxillofac Surg* 1998;27(4):286-9. [[DOI:10.1016/S0901-5027\(05\)80616-2](https://doi.org/10.1016/S0901-5027(05)80616-2)]
30. Akrami S, Navab-Azam A, Akaberi F. Epidemiologic investigation of maxillofacial fractures in admitted patients in Yazd trauma centers (2005-2011). *Yazd J Dent Res* 2014;2(1):46-60.
31. Anvari A. Evaluation of Injury patterns of Oral and Maxillofacial Fractures in Yazd, Iran from the Years 2014 to 2015.[thesis]. Yazd: Shahid Sadoughi University of Medical Science; 2016.
32. Dongas P, Hall GM. Mandibular fracture patterns in Tasmania, Australia. *Aust Dent J* 2002;47(2):131-7. [[DOI:10.1111/j.1834-7819.2002.tb00316.x](https://doi.org/10.1111/j.1834-7819.2002.tb00316.x)]
33. de Almeida OM, Alonso N, Fogaca WC, Rocha DL, Ferreira MC. [Facial fractures. Analysis of 130 cases]. *Rev Hosp Clin Fac Med Sao Paulo* 1995;50 Suppl:10-2.
34. Kamath RA, Bharani S, Hammannavar R, Ingle SP, Shah AG. Maxillofacial trauma in central karnataka, India: an outcome of 95 cases in a regional trauma care centre. *Craniofacial Trauma Reconstr* 2012;5(4):197-204. [[DOI:10.1055/s-0032-1322536](https://doi.org/10.1055/s-0032-1322536)]
35. Ardekian L, Samet N, Shoshani Y, Taicher S. Life-threatening bleeding following maxillofacial trauma. *J Craniomaxillofac Surg* 1993;21(8):336-8. [[DOI:10.1016/S1010-5182\(05\)80493-7](https://doi.org/10.1016/S1010-5182(05)80493-7)]
36. Khosravi H, Kazem-Nejad K. The pattern of jaw and face fractures in 5th Azar hospital in Gorgan - Iran (2003-04). *J Gorgan Univ Med Sci* 2007;8(4):42-45.
37. Adebayo ET, Ajike OS, Adekeye EO. Analysis of the pattern of maxillofacial fractures in Kaduna, Nigeria. *Br J Oral Maxillofac Surg* 2003;41(6):396-400. [[DOI:10.1016/S0266-4356\(03\)00165-7](https://doi.org/10.1016/S0266-4356(03)00165-7)]
38. de Matos FP, Arnez MF, Sverzut CE, Trivellato AE. A retrospective study of mandibular fracture in a 40-month period. *Int J Oral Maxillofac Surg* 2010;39(1):10-5. [[DOI:10.1016/j.ijom.2009.10.005](https://doi.org/10.1016/j.ijom.2009.10.005)]
39. Kontio R, Suuronen R, Ponkkonen H, Lindqvist C, Laine P. Have the causes of maxillofacial fractures changed over the last 16 years in Finland? An epidemiological study of 725 fractures. *Dent Traumatol* 2005;21(1):14-9. [[DOI:10.1111/j.1600-9657.2004.00262.x](https://doi.org/10.1111/j.1600-9657.2004.00262.x)]

How to cite:

Shirinbak I, Baradaran Bagheri A, Kharazifard M.J, Goharshenasan P, Pirouzan M. Prevalence of maxillofacial fractures in hospital patients: A five year retrospective study. *Jorjani Biomed J.* 2020; 8 (3) :75-81