Research Journal of Pharmacognosy (RJP) 3(4), 2016: 51-58 Received: Mar 2016 Accepted: July 2016



Original article

Chemical composition and antibacterial properties of *Ocimum basilicum*, *Salvia officinalis* and *Trachyspermum ammi* essential oils alone and in combination with nisin

T. Mehdizadeh¹, M.S. Hashemzadeh², A. Nazarizadeh³, M. Neyriz-Naghadehi⁴, M. Tat², M. Ghalavand², R. Dorostkar^{2*}

¹Department of Food Hygiene and Quality Control, Faculty of Veterinary Medicine, Urmia University, Iran. ²Applied Virology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran. ³Department of Clinical Pathology and Internal Medicine, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

⁴Department of Food Hygiene and Quality Control, Veterinary Faculty of Islamic Azad University, Urmia Branch, Urmia, Iran.

Abstract

Background and objectives: Plant essential oils are sometimes considered for use as antimicrobial agents in foods and medicines and they could be combined with other antimicrobial agents to strengthen the effect and/or reduce the required dose. This study was conducted to determine the chemical composition of the Ocimum basilicum, Salvia officinalis and Trachyspermum ammi essential oils and evaluate their antibacterial efficiency, alone and in combination with nisin, against Escherichia coli O 157 and Staphylococcus aureus. Methods: The chemical composition of three essential oils (Ocimum basilicum, Salvia officinalis and Trachyspermum ammi) were determined by gas chromatography/mass spectrometry. Further, their antibacterial properties and the synergistic effect of the combination of three essential oils and nisin were also assessed against Escherichia coli and Staphylococcus aureus. The antibacterial activity was determined by evaluation of minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) by broth dilution method in 96-well microplates. The synergistic effects were tested by the checkerboard method and the fractional inhibitory concentration (FIC) index was calculated. Results: The major components of O. basilicum, S. officinalis and T. ammi were linalool (35.99%), 1,8-cineole (22.91%) and p-Cymene (35.5%), respectively. In general, all of the essential oils as well as nisin exerted more considerable antibacterial effects against Gram-positive bacteria than Gram-negative one. The essential oil of T. ammi showed the highest activity against S. aureus with MIC (≤ 0.078 mg/mL) and MBC (≤ 0.156 mg/mL). The combined application showed synergistic activity against E. coli but no change in activity was observed against S. aureus. The most synergistic effect was observed for the combination of nisin and S. officinalis (FIC 0.03). Conclusion: It can be concluded that nisin could enhance the antibacterial potential of the essential oils.

Keywords: antibacterial activity, nisin, Ocimum basilicum, Salvia officinalis, synergy, Trachyspermum ammi

Available at: http://rjpharmacognosy.ir Copy right[©] 2014 by the Iranian Society of Pharmacognosy **Corresponding author: r.dorost@yahoo.com*, Tel/Fax: +9821-88617711

Introduction

Food materials from livestock such as meat and dairy products can be easily contaminated with microorganisms, leading to food spoilage, economic losses and more seriously public health problems. Food spoilage includes physical damage, chemical changes such as oxidation, color changes, or appearance of off-flavors and off-odors resulting from microbial growth and metabolism in the product [1]. Although, novel technologies for preservation and shelf life extension have been introduced recently, unfortunately, these steps do not completely eliminate undesirable pathogens from these products or delay microbial spoilage entirely. Hence, alternative preservation techniques such as novel non-thermal technologies and naturally derived antimicrobial ingredients are under investigation for their application to food products [2]. In the resent years, plant materials have been considered as an excellent alternative of currently available synthetic drugs and preservatives due to their lower or no side effects, broad spectrum antimicrobial activity and economic benefits [3]. Essential oils are rich sources of biologically active compounds and have been assessed to possess antimicrobial properties in vitro and in vivo [4-6]. Basil is the common name for the culinary herb Ocimum basilicum L. of the family Lamiaceae. It has been cultivated from many years ago and it is a wellknown plant worldwide. It is half-hardy and annual, depending on the species and cultivar; the leaves may taste somewhat like anise, with a strong, pungent, often sweet smell. Basil oil has also found a wide application in perfumery, as well as in dental and oral products [7,8]. Salvia officinalis L. (sage) is a perennial, evergreen subshrub, with woody stems, grayish leaves, and blue to purplish flowers. It is a member of the family Lamiaceae and is native to the Mediterranean region, though it has been naturalized in many places throughout the world. The plant has been used for treatment of a number of diseases such as, obesity, diabetes, depression and cancer from ancient times [9,10]. Trachyspermum ammi L. belonging to

Apiaceae family, commonly known as 'Ajwain', is widely grown in arid and semi-arid regions such as Egypt, Iraq, Iran, Afghanistan, Pakistan, and India [11]. There are a number of studies, which shows the potential anti-bacterial effects of the three mentioned plants against various including Escherichia bacteria coli and Staphylococcus aureus [8,12-16]. As a rule, herbal extracts show antimicrobial effects in considerably high concentrations. That is thought to be the major limitation for the application of essential oils in foods. The the high concentrations of essential oils can negatively influence the flavors and odors of foods and result in unacceptability. Therefore, research in this area should be focused on optimizing the essential oil combinations and applications to obtain effective antimicrobial activity at sufficiently low concentrations so as not to adversely affect the organoleptic acceptability of foods [2]. Bacteriocins from lactic acid bacteria such as nisin have been known as effective antibacterial peptides and have been proposed as food preservative since several years ago [2]. In the recent years, some researchers have demonstrated that the combined application of plants essential oils and bacteriocins exerts stronger antibacterial effects than the individual compounds [2,17]. The aim of the present study was to determine the chemical composition of the essential oils of three selected plants (O. basilicum, S. officinalis and T. ammi) and evaluation of the antibacterial efficiency of the essential oils, alone and in combination with nisin, against Escherichia coli O 157 and Staphylococcus aureus.

Experimental

Plant material

Ocimum basilicum and *Salvia officinalis* were purchased freshly from the local market at the end of spring 2013 and *T. ammi* seeds were collected from the rural area of West Azerbaijan at the same period. Pharmacognostic verification of the plants was performed by the Department of Botany and Herbal Medicine, Faculty of Agriculture and Technology, Urmia University, Urmia, Iran. The voucher specimens were deposited at the institute Herbarium (No. 77231, No. 77286 and No. 77317, respectively).

Chemicals and reagents

Nisin was purchased from Sigma–Aldrich Chemical Co. (USA) that contained 2.5% nisin with minimum potency of 10 4 IU/g. Nisin (20 mg) dissolved in 1 mL 0.02 N HCl (Merck, Germany). It was centrifuged for 20 min at 1500 rpm. The supernatant was collected and sterilized and kept in 4 °C until analysis onset.

Extraction of essential oils

Arial parts of *O. basilicum and S. officinalis* as well as the seeds of *T. ammi* were air-dried and powdered (250 g), then subjected to hydrodistillation for 3 h using a Clevenger type apparatus. The oil was dried over anhydrous Na_2SO_4 and preserved in a sealed vial at 4 °C until further analysis.

Gas chromatography/mass spectroscopy (GC/MS) analysis of essential oils

GC/MS analysis was carried on a Varian Gas Chromatograph (Thermofinigan, USA) series 3800 fitted with a VF-5 ms fused silica capillary column (60 m \times 0.25 mm, film thickness 0.25 µm) coupled with a 4000 series mass detector under the following conditions: injection volume 0.5 µL with split ratio 1:60, helium as the carrier gas at 1.0 mL/min constant flow mode, injector temperature 260 °C, oven temperature was programmed from 70 to 180 °C at 3 °C/min. Mass spectra: electron impact (EI+) mode, 70 eV and ion source temperature 290 °C. Mass spectra were recorded over 50–500 a.m.u range.

Identification of components

Identification of the essential oil constituents was performed on the basis of Retention Index [RI, determined with respect to homologous series of *n*-alkanes (C_8-C_{40} , Polyscience Corp., Niles IL) under the same experimental conditions], coinjection with standards and MS Library search (NIST 05 and Wiley) and by comparing with the

MS literature data [18].

Preparation and maintenance of bacteria

The stock culture of *Escherichia coli* O 157 (ATCC 25922) and *Staphylococcus aureus* (ATCC 29213) were obtained from department of Food Hygiene & Quality Control, Faculty of Veterinary Medicine, the University of Urmia, Urmia, Iran. The bacteria were maintained on BHI agar (brain heart infusion) at 4 °C and was grown in BHI broth at 37 °C for 24 h.

Antibacterial assays

The minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of the essential oils were assessed by the micro dilution method, recommended by the National Committee for Clinical Laboratory Standards (NCCLS) [19]. The oils were dissolved in dimethyl sulphoxide 5% (2.50 mg/mL) (Merck, Germany), added to the medium, and then diluted two fold to obtain concentrations in the range of 2.5-0.078 mg/mL. Inoculum suspension with a final concentration of 5×10^5 CFU/mL was added to a 96-well micro plate. The MIC was defined as the lowest concentration of the essential oil at which the microorganism did not demonstrate any visible growth after incubation at 37 °C for 24 h. The MBC was defined as the lowest concentration of the oils at which incubated microorganisms were completely killed [20]. All determinations were performed in triplicate.

Synergy studies, checkerboard method

In combination assays, a checkerboard procedure described by Davidson and Parish [21] was carried out. The checkerboard method was performed using 96-well microplates to obtain the fractional inhibitory concentration (FIC) index of the essential oils and nisin against each pathogenic bacterium. Briefly, the method allows different concentrations of each antimicrobial agent in each well of the assay plate in two axes (x and y) of an 8×8 matrix and it ensures that each well of the assay plate represents a different combination of two antimicrobial agents. Plates consisted of columns containing 25 μ L of an

essential oil (a) which was two-fold diluted in a model medium along the x-axis. Rows of the same plates contained 25 µL of nisin (b) which was also two-fold diluted in the same medium along the y-axis. Subsequently, 90 µL of BHI broth media containing 2.0×10^{-6} CFU/mL of the indicator strain was added to all wells. Plates were then incubated at 37 °C for 24 h and the optical density (600 nm) was measured. The tested compounds consisted of three essential oils of different plants (designated as antimicrobial 'a') (O. basilicum, S. officinalis and T. ammi) and nisin (designated as antimicrobial 'b'). The MIC of each essential oil alone or in combination with nisin was taken as the lowest concentration that inhibited bacterial growth completely after 24 h [4].

The fractional inhibitory concentration (FIC) index was calculated by adding the FIC values of antimicrobial compounds (a) and (b) (FICa+FICb). The FICa and FICb values were represented as the lowest concentrations of the essential oil and nisin, respectively, that caused the inhibition of bacterial growth in the combination tests. Calculations were performed as follows:

FICa = (MICa combined/MICa alone) FICb = (MICb combined/MICb alone) FIC = FICa + FICb

For interpretation of the results, FIC ≤ 0.5 was assigned as a synergistic effect, $0.5 < FIC \leq 1$ represented an additive effect, $1 < FIC \leq 4$ represented no interactive effect and FIC >4 showed antagonistic effect between the two tested antimicrobial agents [22].

Statistical analysis

All the experiments were performed twice in triplicate. The data were analyzed using one-way analysis of variance (ANOVA) tests using SPSS program (version 15, USA). Duncan's multiple range test was conducted to compare the mean values of the tested compounds alone or in combination. A significant difference was presumed at a p value <0.05.

Results and Discussion

Various chemical constituents were identified in the essential oil of O. basilicum, S. officinalis and T. ammi which have been shown in tables 1-3 respectively. As seen from table 1, the major components of O. basilicum were linalool (35.99%), estragole (28.56%) and eucalyptol (7.57%). Table 2 shows GC/MS analysis for S. officinalis; twenty-seven constituents were identified in the essential oil. The major component was 1,8-cineole (22.91%). The detailed chemical composition of T. ammi essential oil has been presented in the table 3; the principal chemical constituent was found to be pcymene (35.5%). In general, the essential oils displayed similar antibacterial effects against Gram-negative bacterium. Besides, Grampositive bacterium, S. aureus was found to be more sensitive to the essential oils than the Gram-negative one. The essential oil of T. ammi showed the highest activity against S. aureus with MIC (≤ 0.078 mg/mL) and MBC (≤ 0.156 mg/mL). Nisin exerted antibacterial activity in similar way to the essential oils. Table 4 also shows the antibacterial activity of O. basilicum, S. officinalis and T. ammi in combination with nisin. The FIC indices of nisin in combination with the essential oils against the bacteria can be found in table 4. The results show that the combined nisin and the different essential oils displayed a synergic activity against E. coli and an indifference effect against S. aureus.

 Table 1. Chemical composition of Ocimum basilicum

 essential oil

03501			
No.	Compounds	KI	Percentage
1	β –pinene	980	0.25
2	Eucalyptol	1043	7.57
3	linalool	1082	35.99
4	Camphor	1121	1.8
5	Estragole	1172	28.56
6	Methyleugenol	1361	6.37
7	Eugenol	1392	2.2
8	Aromadendrene	1419	0.56
9	Caryophyllene	1440	6.54
10	γ-Muurolene	1455	2.24
11	β-Caryophyllene oxide	1557	0.64
12	Tau-Cadinol	1619	7.27

The dominant compounds are indicated in bold.

 Table 2. Chemical composition of Salvia officinalis

 essential oil

No.	compounds	KI	Percentage
1	cis-Salvene	850	0.40
2	a-Thujene	922	13.96
3	Tricyclene	925	0.09
4	α-Pinene	931	12.91
5	Camphene	938	4.74
6	β- Thujene	967	8.91
7	β-Pinene	978	5.93
8	β-Myrcene	980	0.69
9	α-Terpinolene	1086	0.20
10	Camphor	1108	3.28
11	Borneol	1141	6.18
12	l-Phellandrene	1148	0.15
13	α-Terpinene	1155	0.31
14	1,8-Cineole	1191	22.91
15	β-Ocimene	1216	0.10
16	γ-Terpinene	1231	0.41
17	Bornyl acetate	1259	0.39
18	Endobornyl acetate	1283	0.77
19	Aromadendrene	1413	0.56
20	Trans-Caryophyllene	1419	7.41
21	α-Humulene	1430	3.19
22	α-Gurjunene	1432	0.10
23	Trans-Sabinene hydrate	1439	0.13
24	∆-Cadinene	1499	0.24
25	Veridiflorol	1587	3.08
26	1-Naphthalenepropanol	2026	0.11
27	Others	-	1.03

The dominant compounds are indicated in bold.

Food material contamination with microorganisms is considered a major concern and can lead to public health problems such as poisoning and even death. Considering the food spoilage, different aspects of the development of novel preservatives with the ability of protection against wide spectrum of microorganisms, with no impact on odor and flavor, is attractive. In recent years, the demand for chemical preservatives has considerably decreased because of their undesired effects on food and also on human health. Plants essential oils have been interesting agents as excellent replacements for the synthetic preservatives. Since higher concentrations of essential oils are generally required to ensure their antimicrobial activity for food preservation as compared to in vitro system, their application may be limited due to changes in organoleptic and textural quality of food or interactions of essential oils with food components [22].

Table 3. Chemical composition of *Trachyspermum ammi* essential oil

No.	Compounds	KI	Percentage
1	α-Thujene	924	1.34
2	α-Pinene	930	1.75
3	Camphene	940	0.22
4	Sabinene	965	0.08
5	2-β-Pinene	976	5.10
6	β-Thujone	978	0.48
7	β-Myrcene	982	2.17
8	Cis-Sabinene Hydrate	1065	0.17
9	α-Terpinene	1080	0.42
10	p-Cymene	1088	35.5
11	α-Terpinolene	1096	0.12
12	Thujone	1105	0.30
13	Camphor	1109	0.09
14	Pulegone	1208	0.14
15	γ-Terpinene	1230	0.09
16	γ-Terpinene	1235	32.52
17	Thymol	1259	18.32
18	Trans-Caryophyllene	1405	0.29
19	Aromadendrene	1415	0.02
20	α-Humulene	1435	0.13
21	Caryophyllene Oxide	1553	0.017
22	Others	-	0.489

The dominant compounds are indicated in bold.

The current study targeted this limitation and was conducted to evaluate the synergistic effects of the essential oils and a well-known bacteriocin, nisin, as a promising natural alternative. The major components of O. basilicum were linalool and estragole. Our data confirmed previous study, which reported that the most abundant component of O. basilicum was linalool. Further, it has been claimed that the season of O. basilicum cultivation could influence the chemical composition and antimicrobial activity of the plant. Generally, essential oils from winter and autumn crops exhibited greater antimicrobial activity, which might be attributed to the high contents of linalool and other oxygenated compounds in these samples [23]. Based on a previous comprehensive study [24], four dominant constituents of O. basilicum essential oil are methyl chavicol, linalool, methyl eugenol methyl cinnamate. The predominant and constituents of S. officinalis were found to be 1,8cineole and α -tujone. This finding is similar to the previous researches [16,25] but partly different with those of reported from Brazil [26].

Mehdizadeh T. et al.

Table 4. Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of nisin (I.U/mL) and *Ocimum basilicum, Salvia officinalis* and *Trachyspermum ammi* essential oil (mg/mL) against *E. coli* and *S. aureus*. The fractional inhibitory concentration (FIC) was calculated for determination of the antibacterial relationship between the plants essential oil and nisin.

				Bacteriun	n			
Test compound	Escherichia coli O 157			Staphylococcus aureus				
	MIC	MBC	FIC	Result	MIC	MBC	FIC	Result
O. basilicum	10	10<	-	-	0.62	1.25	-	-
S. officinalis	10<	10<	-	-	1.25	2.5	-	-
T. ammi	10	10<	-	-	≤ 0.078	≤0.156	-	-
Nisin	1000	1000	-	-	7.8≥	15.6≥	-	-
Nisin+ O. basilicum	31.25+0.31 ^a	$62.5 + 0.62^{a}$	0.06	\mathbf{S}^*	$7.8 + 0.078^{a}$	15.6+0.156 ^a	1.06	Ι
Nisin+ S. officinalis	15.6+0.156 ^b	31.2+0.31 ^b	0.03	S	$7.8 + 0.078^{a}$	15.6+0.156 ^a	1.06	Ι
Nisin+ T. ammi	31.25+0.31 ^a	62.5+0.62 ^a	0.06	S	7.8+0.078 ^a	15.6+0.156 ^a	2	Ι

*S: synergism, I: indifference within each column, the same small letters are not significantly different (*p*>0.05)

In this regard, Santos-Gomes and Fernandes-Ferreira have reported organ and season dependent variation in the essential oil composition of S. officinalis [27]. Our results show that S. officinalis essential oil showed great antibacterial activity against S. aureus but very little effects on E. coli. This aspect of the work is inconsistent with a previous study, which showed limited effects on both bacteria [16]. Phytochemical analysis of T. ammi essential oil revealed that the principal compositions were pcymene and γ -terpinene. This was in agreement with previous works [28]. In our study, T. ammi possessed the highest antibacterial activity. It had been reported that, T. ammi essential oil had better or equal efficacy as compared to the standard antibiotics [29]. Our results showed that the essential oils were more active against the Gram-positive bacterium than the Gram-negative one. This aspect of the work is also consistent with the previous reports [2,30]. The different interactions, permeability and diffusion of various hydrophobic agents across the cell membrane may influence the extent of the antimicrobial effect. Hydrophobic components can usually diffuse across lipid bilayers easily. whereas hydrophilic agents are often transported passively via channels present in the membranes. In the case of Gram-negative bacteria, which are less susceptible to various antimicrobials, the presence of lipopolysacharide in the outer membrane protects against penetration of hydrophobic antimicrobial agents; thus, higher doses of antimicrobial agents are required compared to those needed for Gram-positive bacteria [2,20]. Nisin is effective against the Gram-positive bacteria and has also shown its efficacy against the Gram-negative bacteria when it is used as an additional technological barrier (in the application of multiple processes or barriers) for food safety and preservation. It is approved as a food preservative in more than 40 countries worldwide [4]. In the current work, nisin showed considerable antibacterial activity against S. aureus, however, it was ineffective against E. coli. In the recent years, the combined application of bacteriocin and essential oils has attracted great interest. As can be seen from table 4 all of the combination showed synergistic effect (FIC≤0.5). The mechanisms on Ε. coli responsible for the synergetic effects have not yet clearly been established, but it has been claimed that, the essential oils usually act on cell membrane and result in increase in permeability and subsequent the cell death [31,32]. The use of essential oils and bacteriocins together play an important role in the formation of membrane pores, which alter the permeability of the membrane, the proton motive force, the efflux of amino acid and the pH gradient of bacteria [4]. Taking together, the results show that the combined use of the essential oils and nisin, exert synergetic antibacterial effect on E. coli. This would allow the lower concentration use of

essential oils and thereby lower or no impact on food quality. Adjustment of precise minimal concentration of essential oils and investigation of efficacy of this method on shelf life extension should be investigated in further researches

Acknowledgments

We would like to express our sincere gratitude and appreciation to Mrs. F. Golmoradi and Mr. H. Jafari-Namin, for their tremendous support.

Declaration of interest

The authors declare that there is no conflict of interest. The authors alone are responsible for the content of the paper.

References

- [1] Gram L, Ravn L, Rasch M, Bruhn JB, Christensen AB, Givskov M. Food spoilage interactions between food spoilage bacteria. *Int J Food Microbiol*. 2002; 78(1-2): 79-97.
- [2] Gutierrez J, Barry-Ryan C, Bourke P. Antimicrobial activity of plant essential oils using food model media: efficacy, synergistic potential and interactions with food components. *Food Microbiol.* 2009; 26(2): 142-150.
- [3] Nazarizadeh A, Mikaili P, Moloudizargari M, Aghajanshakeri S, Javaherypour S. Therapeutic uses and pharmacological properties of *Plantago major* L. and its active constituents. *J Basic Appl Sci Res.* 2013; 3(9): 212-221.
- [4] Turgis M, Vu KD, Dupont C, Lacroix M. Combined antimicrobial effect of essential oils and bacteriocins against foodborne pathogens and food spoilage bacteria. *Food Res Int.* 2012; 48(2): 696-702.
- [5] Joy B, Rajan A, Abraham E. Antimicrobial activity and chemical composition of essential oil from *Hedychium coronarium*. *Phytother Res.* 2007; 21(5): 439-443.
- [6] Bergonzelli GE, Donnicola D, Porta N, Corthesy-Theulaz IE. Essential oils as components of a diet-based approach to management of *Helicobacter* infection.

Antimicrob Agents Ch. 2003; 47(10): 3240-3246.

- [7] Fandohan P, Gnonlonfin B, Laleye A, Gbenou JD, Darboux R, Moudachirou M. Toxicity and gastric tolerance of essential oils from *Cymbopogon citratus*, *Ocimum gratissimum* and *Ocimum basilicum* in Wistar rats. *Food Chem Toxicol*. 2008; 46(7): 2493-3497.
- [8] Suppakul P, Miltz J, Sonneveld K, Bigger SW. Antimicrobial properties of basil and its possible application in food packaging. J Agric Food Chem. 2003; 51(11): 3197-3207.
- [9] Hamidpour M, Hamidpour R, Hamidpour S, Shahlari M. Chemistry, pharmacology, and medicinal property of sage (*Salvia*) to prevent and cure illnesses such as obesity, diabetes, depression, dementia, lupus, autism, heart disease, and cancer. *J Tradit Complement Med.* 2014; 4(2): 82-88.
- [10] Clebsch B, Barner CD. The new book of salvias: sages for every garden. New York: Timber Press, 2003.
- [11] Bairwa R, Sodha RS, Rajawat BS. Trachyspermum ammi. Pharmacogn Rev. 2012; 6(11): 56-60.
- [12] Horiuchi K, Shiota S, Hatano T, Yoshida T, Kuroda T, Tsuchiya T. Antimicrobial activity of oleanolic acid from *Salvia officinalis* and related compounds on vancomycin-resistant *Enterococci* (VRE). *Biol Pharm Bull*. 2007; 30(6): 1147-1149.
- [13] Kaya I, Yigit N, Benli M. Antimicrobial activity of various extracts of *Ocimum* basilicum L. and observation of the inhibition effect on bacterial cells by use of scanning electron microscopy. *Afr J Tradit Complement Altern Med.* 2008; 5(4):363-369.
- [14] Stefanovic OD, Stanojevic DD, Comic LR. Synergistic antibacterial activity of *Salvia* officinalis and Cichorium intybus extracts and antibiotics. Acta poloniae pharmaceutica. 2012; 69(3): 457-463.
- [15] Adigüzel A, Güllüce M, ŞENGÜL M. Antimicrobial effects of *Ocimum basilicum*

(Labiatae) extract. *Turk J Biol*. 2005; 29(3): 155-160.

- [16] Longaray Delamare AP, Moschen-Pistorello IT, Artico L, Atti-Serafini L, Echeverrigaray S. Antibacterial activity of the essential oils of *Salvia officinalis* L. and *Salvia triloba* L. cultivated in South Brazil. *Food Chem.* 2007; 100(2): 603-608.
- [17] Govaris A, Solomakos N, Pexara A, Chatzopoulou PS. The antimicrobial effect of oregano essential oil, nisin and their combination against *Salmonella enteritidis* in minced sheep meat during refrigerated storage. *Int J Food Microbiol*. 2010; 137(2-3): 175-180.
- [18] Adams RP. Identification of essential oil components by gas chromatography/quadrupole mass spectroscopy. J Am. Soc Mass Spectrom. 2005; 16(11): 1902-1903.
- [19] Ashour ML, El-Readi M, Youns M, Mulyaningsih S, Sporer F, Efferth T, Wink M. Chemical composition and biological activity of the essential oil obtained from *Bupleurum marginatum* (Apiaceae). J *Pharm Pharmacol.* 2009; 61(8): 1079-1087.
- [20] Rashid S, Rather MA, Shah WA, Bhat BA. Chemical composition, antimicrobial, cytotoxic and antioxidant activities of the essential oil of *Artemisia indica* Willd. *Food Chem.* 2013; 138(1): 693-700.
- [21] Davidson PM, Parish ME. Methods for testing the efficacy of food antimicrobials. *Food Technol.* 1989; 52(1): 148-154.
- [22] Gutierrez J, Barry-Ryan C, Bourke P. The antimicrobial efficacy of plant essential oil combinations and interactions with food ingredients. *Int J Food Microbiol.* 2008; 124(1): 91-97.
- [23] Hussain AI, Anwar F, Hussain Sherazi ST, Przybylski R. Chemical composition, antioxidant and antimicrobial activities of basil (*Ocimum basilicum*) essential oils depends on seasonal variations. *Food Chem*. 2008; 108(3): 986-95.

- [24] Lawrence BW. A further examination of the variation of Ocimum basilicum L. In: Lawrence BM, Mokherjee BD, Willis BJ, Eds. Flavors and Fragrances. New York: Elsevier, 1988.
- [25] Marino M, Bersani C, Comi G. Impedance measurements to study the antimicrobial activity of essential oils from Lamiaceae and Compositae. *Int J Food Microbiol*. 2001; 67(3): 187-195.
- [26] Porte A, Godoy R, Maia-Porte L. Chemical composition of sage (*Salvia officinalis* L.) essential oil from the Rio de Janeiro State (Brazil). *Rev Bras Plant Med.* 2013; 15(3): 438-441.
- [27] Santos-Gomes PC, Fernandes-Ferreira M.
 Organ- and season-dependent variation in the essential oil composition of *Salvia officinalis* L. cultivated at two different sites. *J Agric Food Chem.* 2001; 49(6):2908-2916.
- [28] Moein MR, Zomorodian K, Pakshir K, Yavari F, Motamedi M, Zarshenas MM. *Trachyspermum ammi* (L.) sprague: chemical composition of essential oil and antimicrobial activities of respective fractions. *Evid-Based Complement Altern Med.* 2015; 20(1): 50-56.
- [29] Kaur GJ, Arora DS. Antibacterial and phytochemical screening of *Anethum* graveolens, Foeniculum vulgare and *Trachyspermum ammi. BMC Complement Altern Med.* 2009; 9(1): 1-10.
- [30] Burt S. Essential oils: their antibacterial properties and potential applications in foods-a review. *Int J Food Microbiol*. 2004; 94(3): 223-253.
- [31] Lambert RJ, Skandamis PN, Coote PJ, Nychas GJ. A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *J Appl Microbiol*. 2001; 91(3): 453-462.
- [32] Ultee A, Kets EPW, Smid EJ. Mechanisms of action of carvacrol on the food-borne pathogen *Bacillus cereus*. *Appl Environ Microbiol*. 1999; 65(10): 4606-4610.