

Burden of Disease Attributable to Suboptimal Breastfeeding in Iran during 1990-2010; Findings from the Global Burden of Disease Study 2010

Roya Kelishadi¹, *Hossein Ansari², *Mostafa Qorbani³, Mohsen Jari¹, Shirin Djalalinia⁴, Asal Ataie-Jafari⁵, Mina Poorrezaeian⁶, Morteza Mansourian⁷, Saeid Safiri⁸, Omid Safari⁹, Mehdi Noorozi¹⁰, Hamid Asayesh¹¹, Rasool Mohammadi¹²

¹Child Department of Pediatrics, Child Growth and Development Research Center, Research Institute for Primordial Prevention of Non-communicable Disease, Isfahan University of Medical Sciences, Isfahan, Iran. ²Health Promotion Research Center, Department of Epidemiology and Biostatistics, Zahedan University of Medical Sciences, Zahedan, Iran. ³Department of Community Medicine, Alborz University of Medical Science, Karaj, Iran & Non-communicable Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran. ⁴Development of Research & Technology Center, Deputy of Research and Technology, MOHME, Tehran, Iran. ⁵Department of Nutrition, Science and Research Branch, Islamic Azad University, Tehran, Iran. ⁶Chronic Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran. ⁷Department of Health Education and Promotion, School of health, Iran University of Medical Sciences, Tehran, Iran. ⁸Managerial Epidemiology Research Center, Department of Public Health, School of Nursing and Midwifery, Maragheh University of Medical Sciences, Maragheh, Iran. ⁹Department of Pediatrics, Alborz University of Medical Science, Karaj, Iran. ¹⁰Substance Abuse and Dependence Research Centre, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran. ¹¹Department of Medical Emergencies, Qom University of Medical Sciences, Qom, Iran. ¹²Department of Epidemiology, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Abstract

Background: This study uses data of the global burden of diseases (GBD) study 2010 to report death, disability-adjusted life year (DALYs), years of life lived with disability (YLDs) and years of life lost due to premature mortality (YLLs), attributed to suboptimal breastfeeding by age and gender during 1990 to 2010 in Iran. **Materials and Methods:** The GBD assessments were used, together with estimates of death and DALYs due to specific risk factors to calculate the attributed burden of each risk factor exposure compared with the theoretical-minimum-risk exposure. Uncertainties in the distribution of exposure, relative risks, and relevant outcomes were incorporated into estimates of mortality attributable and burden and were presented as 95 % uncertainty interval (UI). **Results:** In both genders, the age standardized DALYs rates and the age standardized death rate [(from 5 (95% UI: 2-8) to 1 (95% UI: 0-2) per 100,000 populations], attributed to breastfeeding, had a decreasing trends. The age standardized YLD rate increased from 7 (95% UI: 2-15) to 10 (95% UI: 3-23) per 100,000 populations in boys and, from 7(95% UI: 2-16) to 11(95% UI: 3-26) per 100,000 populations in girls. The YLD changes showed some variation according to age categories. For both genders, the age standardizes YLL rate decreased from 395 (95% UI: 185-681) per 100,000 populations to 111(95% UI: 42-213) per 100,000 populations. **Conclusion:** The burden attributed to suboptimal breastfeeding had a considerable reduction rate from 1990 to 2010. Additional studies on burden of exclusive breastfeeding with more accurate data are recommended for policies make decision.

Key Words: Breastfeeding, Burden, Disability-adjusted life year, Iran, Years lost due disability.

*Please cite this article as: Kelishadi R, Ansari H, Qorbani M, Jari M, Djalalinia Sh, Ataie-Jafari A, et al. Burden of Disease Attributable to Suboptimal Breastfeeding in Iran during 1990-2010; Findings from the Global Burden of Disease Study 2010. Int J Pediatr 2016; 4(9): 3535- 49. DOI: 10.22038/ijp.2016.7185

*Corresponding Authors:

Hossein Ansari, Department of Epidemiology and Biostatistics, Health Promotion Research Center, Zahedan University of Medical Sciences, Hesabi Square, Zahedan, Iran & **Mostafa Qorbani**, Departments of Community Medicine, School of Medicine, Alborz University of Medical Sciences, Karaj, Iran.

Email: ansarih88@gmail.com & mqorbani1379@yahoo.com

Received date Mar 10, 2016; Accepted date: Jun 22, 2016

1- INTRODUCTION

Previous studies have shown that more than 56 million people around the world died in 2001. Of these, more than 10.6 million were children (1). About 99% of child deaths occurred in low- and middle-income countries. In addition, more than half of deaths in this year were due to respiratory diseases, diarrhea, malaria, AIDS, measles and perinatal conditions (1). Infectious diseases, particularly respiratory and gastrointestinal infections, are major causes of hospitalization and death in infants and children (1-3).

The risk of infectious diseases in children depends on many factors; most notably gender, gestational age, birth weight, nutritional status, socio-economic level of the family, family size, parental educational level and smoking status, as well as children care status (3-7). Infectious diseases are still a main health problem for children. For instance, 1.1 episodes of diarrhea per child-year for children under 5 years of age are reported in the USA (8). Respiratory infection in infants is still a common problem in American children, where about 3% of infants are hospitalized with moderate to severe respiratory infections every year (9). Several studies show that the prevalence of overweight and obesity in adults and children is increasing. Overweight and obesity also have been shown to be underlying or aggravating cause of many physical and mental diseases. Therefore, weight control has been proposed as a public health priority (10, 11). Chronic non-communicable diseases (NCDs), the most common of them are cardiovascular diseases, diabetes mellitus, hypertension, hypercholesterolemia and cancer, are the main public health problems (12,13). Several studies show a protective effect of breastfeeding in infancy against infectious and allergic diseases during infancy, childhood, or even throughout the life (14-

16). This is a protective effect against most non-communicable diseases like cardiovascular diseases (17-20), hypercholesterolemia (21, 23), diabetes mellitus 1, 2 (22, 24), hypertension (25, 26) and cancer (27, 29) during childhood, adolescence and adulthood.

Due to the numerous benefits of breastfeeding for infants, this paper aimed to present the age-and gender-specific DALYs and deaths rate attributed to suboptimal breastfeeding in Iran from 1990 to 2010. Data were obtained from global burden of diseases (GBD) 2010, which have been gathered and calculated by the Institute for Health Metrics and Evaluation (IHME), providing results for the years 1990, 1995, 2000, 2005, and 2010. We also, discuss the data, methods, and limitations of the GBD study 2010 in terms of different aspects of policy making.

2- MATERIALS AND METHODS

In brief, burden of diseases, injuries and risk factors are expressed in DALYs, a summary measurement of population health gap. DALYs reveal the sum of the years of life lost due to premature mortality (YLL) in the population and years of life lost due to disability (YLD) (30). More practical details about the data and methods used for specific risk factors including breast feeding are also available (31). In this approach also suboptimal breastfeeding include a third level to distinguish between nonexclusive breastfeeding during the first 6 months (31).

2-1. Study Design and Population

Using data of the GBD study 2010, this paper presents the trends in deaths and DALYs attributed to breast-feeding by gender and age from 1990 to 2010 for Iranian children.

2-2. Methods

This project was a systematic effort for data gathering and estimation processing of 291 types of diseases and injuries and 67 risk factors in 187 countries to calculate the global and regional comparative risk assessment of deaths and DALYs caused by different risk factors and diseases as described previously (31-37).

2-3. Ethical considerations

Ethical issues (Including plagiarism, Informed Consent, misconduct, data fabrication and/or falsification of data, double publication and/or submission, redundancy, etc) have been completely observed by the authors.

2-4. Data analyses

Using the Population Attributable Fraction (PAF), considered the fraction by which the occurrence of interested outcomes changed under a sustained alternative and more favorable exposure distribution. Death and disease burden attributable to risk factors (PAF) were computed by comparing the current distribution of exposure to the theoretical minimum risk counterfactual distribution of the exposure for each year, gender, age group, and cause. Uncertainty Interval (UI) was also calculated in the simulation analysis.

$$\text{PAF} = \frac{\text{Counterfactual population risk} - \text{Factual population risk}}{\text{Factual population risk}}$$

To assess the full effects of exposure distributions of breast feeding, five main steps were used (38). First, risk-outcome pairs were selected to be included in the analysis based on causal association criteria. The cause-specific mortality extracted from available published cohort studies (31). Estimation of distribution of exposure and statistical modeling such as space-time/Gaussian process regression model or meta-regression were followed through a wide range of covariates in databases that were generated in the GBD

2010 project was verified. In the third step, the effect size of breastfeeding outcomes' specific mortality considered. In this regard for each risk-outcome pair, relative risk estimated for each unit of exposure. The distribution of exposure has been compared to an alternative (counterfactual) distribution (theoretical-minimum-risk) and uncertainty in the estimates is presented as 95% uncertainty interval (95% UI).

3- RESULTS

Table.1 presents the cause specific DALYs rates per 100,000 populations attributed to breastfeeding in different age groups of children from 1990 to 2010. In both genders, the age standardized DALYs rates attributed to breastfeeding had a decreasing trend from 1990 to 2010. In another words 492 (95% UI: 191- 689) life year that were estimates in 1990 as life years that are spending with disability, through a decreasing trend reached to 121 (95% UI: 50- 224) in 2010.

The age standardized death rate attributed to breast feeding decreased 4 cases/100,000 population from 1990 to 2010 (from 5 (95% UI: 2-8) to 1 (95% UI: 0-2 respectively).

Table.2 presents the cause- specific death rates per 100,000 populations attributed to breastfeeding in different age groups between 1990 and 2010. The age standardized YLD rate or years of life lived with disability for both gender due to breast feeding increased from 1990 to 2010 [from 7(95% UI: 2-15) to 10(95% UI: 3-23) in boys, and from 7(95% UI: 2-16) to 11(95% UI: 3-26) in girls). The YLD changes or years of life that lost due to premature mortality showed some variation according to age categories (**Table.3**).

It is estimated that YLL follow a decreasing rate during the time between the 1990 and 2010. For both gender, age

standardizes rate decreased from 395(95% UI: 185-681) in 1990 to 111(95% UI: 42-213) in 2010. The highest rate of YLL for both sexes was observed in neonatal age group (**Table.4**).

4- DISCUSSION

This study showed that the DALY, death, and YLL rates attributed to suboptimal breastfeeding in Iran had a decreasing trend from the year 1990 to 2010 in infants and different age groups of children, as well as in all other age groups.

Results of previous studies showed that if 90% of USA families could comply with the recommendations of exclusive breastfeeding for 6 months, the United States would prevent a yearly excess 911 deaths of infants (39). Several studies also have shown more than 10 million child deaths and 44 million DALYs per year (40). Exclusive breastfeeding in the first 6 months of life is associated with lower incidence of infectious diseases, and lower deaths and DALY rate in children (39-42). Also, previous studies show that exclusive breastfeeding is associated with lower risk of obesity and high blood pressure in children and adolescents (43-45).

These effects assigned to factors such as Lactoferrin (LF), Immune Globulin A (IgA), secretion polysaccharides and oligosaccharides (3, 14). This protective effect was also dependent on the duration of breastfeeding (3, 7, 14). Related studies confirmed that exclusive breastfeeding significantly reduces the risk of neonatal morbidity such as; sepsis and severe infections (especially in full term infants) (3, 4, 15), acute otitis media (AOM) and viral upper respiratory tract infection and eustachian tube dysfunction (46).

Several factors such as the family socio-economic status, religious beliefs and cultural customs, lifestyle, mother's employment, parents' education, parents' age at marriage, maternal age,

advertisements by milk powder producing companies and the promotion of exclusive breast-feeding by mass media, are effective on increasing the prevalence of exclusive breastfeeding in children under 6 months and on continuation of breastfeeding (47-50).

This has been shown in many parts of the world, as the studies show an increase of the prevalence of exclusive breastfeeding after 2002 (51-53). The Demographic and Health Survey (DHS) conducted in the years 2000, 2005 and 2010 In Iran, showed an upward trend in the prevalence of exclusive breastfeeding in the first 6 months of life (54).

4-1. Study limitations and strengths

There are some limitations to the GBD Study 2010. In Iran, exposure estimations for many risk factors, including suboptimal breastfeeding, were affected by data source limitations. Estimation of exposure with the approach used here is another weakness of the study because of the data gaps for specific regions such as Iran. Finally, the GBD findings were at a national level, while sub-national information about the burden of diseases and risk factors are required for policy making. So, the burden of diseases, injuries, and risk factors should be evaluated at sub-national level. Also, to the best of our knowledge, present study is the first study, which report burden of diseases attributed to suboptimal breastfeeding in Iran using the GBD data.

5- CONCLUSIONS

Present study, revealed that burden attributed to suboptimal breastfeeding had a considerable reduction rate from 1990 to 2010. Similar to other part of world, results of the reduction of child mortality and DALY in Iran, comply with the upward trends in exclusive breastfeeding. Additional studies on burden of exclusive breastfeeding with more accurate data at

national and subnational level are recommended for policies make decision.

6- CONFLICT OF INTEREST: None.

7- REFERENCES

1. Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray C. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *The Lancet* 2006; 367(9524):1747-57.doi:10.1016/S0140-6736(06)68.
2. Eliminate Disparities in Infant Mortality. Office of Minority Health: Centers for Disease Control and Prevention. 2009. Available at: <http://www.cdc.gov/omhd/AMH/factsheets/infant.htm>.
3. Liessbeth D, Vincent WV, Albert H, Henriette AM. Prolonged and exclusive breastfeeding reduces the risk of infectious diseases in infancy. *Pediatrics* 2010;126:188-25.
4. Koopmanlp, Smit HA, Heijnen ML. Respiratory infections in infants: interaction allergy, child care, and siblings – the PIAMA study. *Pediatrics* 2001;108(4):943-48.
5. Caudri D, Wijga A, Gehring U. Respiratory symptoms in the first 7 years of life and birth weight at term; the PIAMA Birth Cohort. *Am J Respir Crit Care Med* 2007; 175(10): 1078-85.
6. Nafstad P, Jaakkola JJ, Hagen JA, Botten G, KongerudJ. Breastfeeding , maternal smoking and lower respiratory tract infections. *Eur Respir J* 1996;9(12): 2623-29.
7. Rylander E, Eriksson M, Pershagen G, Nordvall L, Ehrnst A, Ziegler T. Wheezing bronchitis in children: incidence, viral infections, and other risk factors in a defined population. *Pediatr Allergy Immunol* 1996; 7(1); 6-11.
8. Imhoff B, Morse D, Shiferaw B. Hawkins M, Vugia D, Lance-Parker S, et al. Burden of self-reported acute diarrheal illness in FoodNet surveillance areas, 1998-1999. *Clin Infect Dis* 2004; 38(Suppl.3):S219–S226.
9. Shay DK, Holman RC, Newman RD, Liu LL, Stout JW, Anderson LJ. Bronchiolitis-associated hospitalizations among US children, 1980-1996. *JAMA* 1999; 282(15):1440–46.
10. Kelishadi R. Childhood overweight, obesity, and the metabolic syndrome in developing countries. *Epidemiol Rev* 2007; 29:62-76
11. Kelishadi R, Haghdoost AA, Sadeghirad B, Khajehkazemi R. Trend in the prevalence of obesity and overweight among Iranian children and adolescents: A systematic review and meta-analysis. *Nutrition* 2014; 30(4):393-400.
12. Alwan A. World Health Organization. Global status report on non communicable diseases 2010. Geneva: World Health Organization; 2011. Available at: <http://apps.who.int/iris/handle/10665/44579>.
13. Kelishadi R, Farajian S. The protective effects of breastfeeding on chronic non-communicable diseases in adulthood: A review of evidence. *Adv Biomed Res* 2014; 3:3-7.
14. Kramer MS, Guo T, Platt RW. Infant growth and health outcomes associated with 6 mo of exclusive breastfeeding. *Am J Clin Nutr* 2003; 78(2): 291- 95.
15. Pardo – Crespo R, Perez – Iglesias R, Llorca J. Breast-feeding and risk of hospitalization for all causes and fever of unknown origin. *Eur j public Health* 2004; 14(3): 230-34.
16. Saarinen UM, Kajosaari M. Breastfeeding as prophylaxis against atopic disease: prospective follow-up study until 17 years old. *Lancet* 1995; 21: 1065-69.
17. Poulton R, Williams S. Breastfeeding and risk of overweight. *JAMA* 2001; 286:1449-50.
18. Owen CG, Martin RM, Whincup PH, Smith GD, Cook DG. Effect of infant feeding on the risk of obesity across the life course: a quantitative review of published evidence. *Pediatrics* 2005;115(5):1367–77.
19. Dewey KG. Is breast feeding protective against child obesity? *J Hum Lact* 2003;19:9-18.
20. Izadi V, Kelishadi R, Qorbani M, Esmaeilmotlagh M, TaslimiM, Heshmat R, Ardalan G, et al. Duration of breast-feeding and cardiovascular risk factors among Iranian children and adolescents: The CASPIAN III study. *Nutrition* 2013; 29: 744-51.
21. Owen CG, Whincup PH, Odoki K, Gilg JA, Cook DG. Infant feeding and blood

- cholesterol: a study in adolescents and a systematic review. *Pediatrics* 2002;110(3):597–608.
22. Taylor JS, Kacmar JE, Nothnagle M, Lawrence RA. A systematic review of the literature associating breastfeeding with type 2 diabetes and gestational diabetes. *J Am Coll Nutr* 2005;24(5):320–6.
23. Salmenpera L, Perheentupa J, Siimes MA, Adrian TE, Bloom SR, Aynsley-Green A. Effects of feeding regimen on blood glucose levels and plasma concentrations of pancreatic hormones and gut regulatory peptides at 9 months of age: Comparison between infants fed with milk formula and infants exclusively breast-fed from birth. *J Pediatr Gastroenterol Nutr* 1988;7:651-6.
24. Ravelli AC, van der Meulen JH, Osmond C, Barker DJ, Bleker OP. Infant feeding and adult glucose tolerance, lipid profile, blood pressure, and obesity. *Arch Dis Child* 2000; 82:248-52.
25. Owen CG, Whincup PH, Gilg JA, Cook DG. Effect of breast feeding in infancy on blood pressure in later life: systematic review and meta-analysis. *BMJ* 2003;327(7425):1189–95.
26. Martin RM, Ness AR, Gunnell D, Emmett P, Davey Smith G. Does breast-feeding in infancy lower blood pressure in childhood? The Avon Longitudinal Study of Parents and Children (ALSPAC). *Circulation* 2004; 109:1259-66.
27. Guise JM, Austin D, Morris CD. Review of case-control studies related to breastfeeding and reduced risk of childhood leukemia. *Pediatrics* 2005;116(5):e724–e731.
28. Higgins JP, Whitehead A, Turner RM, Omar RZ, Thompson SG. Meta-analysis of continuous outcome data from individual patients. *Stat Med* 2001; 20(15):2219–41.
29. Davis MK. Review of the evidence for an association between infant feeding and childhood cancer. *Int J Cancer Suppl* 1998; 11: 29-33.
30. Petersen PE, Ogawa H. The global burden of periodontal disease: towards integration with chronic disease prevention and control. *Periodontology* 2012; 60(1):15.
31. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380(9859): 2224.
32. Murray CJ, Ezzati M, Flaxman AD, Lim S, Lozano R, Michaud C, et al. GBD 2010: design, definitions, and metrics. *The Lancet* 2013; 380(9859): 2063-66.
33. Wang H, Dwyer-Lindgren L, Lofgren KT, Rajaratnam JK, Marcus JR, Levin-Rector A, Levitz CE, et al. Age-specific and sex-specific mortality in 187 countries, 1970-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380(9859): 2071.
34. Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet* 2013; 380(9859): 2163-2196.
35. Salomon JA, Vos T, Hogan DR, Gagnon M, Naghavi M, Mokdad A, et al., Common values in assessing health outcomes from disease and injury: disability weights measurement study for the Global Burden of Disease Study 2010. *The Lancet* 2013; 380(9859): 2129-43.
36. Salomon JA, Wang H, Freeman MK, Vos T, Flaxman AD, Lopez AD, et al., Healthy life expectancy for 187 countries, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380(9859): 2144.
37. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet* 2012; 380(9859): 2095.
38. Burnett RT, Pope CA, Ezzati M, Olives C, Lim SS, Mehta S, et al. An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate

- matter exposure. 2014. Available at: <http://ehp.niehs.nih.gov/1307049/>.
39. Bartick M, Reinhold A. The Burden of Suboptimal Breastfeeding in the United States: A Pediatric Cost Analysis. *pediatrics* 2010; 125(5):1048-56. doi: 10.1542/peds.2009-1616.
40. Black RE, Morris SS, Bryce J. Where and why are 10 million children dying every year? *The Lancet* 2003; 361: 2226-34. doi: 10.1016/s0140-6736(03)13779-8.
41. Jones G, Steketee RW, Black RE, Bhutta ZA, Morris SS. How many child deaths can we prevent this year? *The Lancet* 2003; 362: 65-71. doi: 10.1016/s0140-6736(03)13811-1.
42. Bland RM, Little KE, Coovadia HM, Coutoudis A, Rollins NC, Newell ML. Intervention to promote exclusive breastfeeding for the first 6 months of life in a high HIV prevalence area. *AIDS* 2008; 22: 883-91. doi: 10.1097/qad.0b013e3282f768de.
43. Kelishadi R, Rezaei F, Djalalinia S, Asadi M, Miranzadeh S, Motlagh ME, et al. Is duration of breastfeeding associated with anthropometric measures in children and adolescents? the weight disorders survey of the CASPIAN-IV study. *International Journal of Pediatrics* 2016;4(8):3299-313.
44. Djalalinia S, Qorbani M, Heshmat R, Motlagh M, Ardalan G, Bazyar N, et al. Association of Breast Feeding and Birth Weight with Anthropometric Measures and Blood Pressure in Children and Adolescents: The CASPIAN-IV Study. *Pediatrics and neonatology* 2015; 56(5):324-33.
45. Amorim RD, Coelho AF, Lira PI, Lima MD. Is breastfeeding protective for blood pressure in schoolchildren? A cohort study in northeast Brazil. *Breastfeeding Medicine* 2014;9(3):149-56.
45. Auinger P, Lanphear BP, Kalkwarf HJ. Trends in otitis media among children in the United States. *Pediatrics* 2003; 112: 514-20.
46. AnneMerewood MA, Supriya D. Mehta, Laura Beth Chamberlain BA, Barbara L. Philipp, Bauchner H. Breastfeeding Duration Rates and Factors Affecting Continued Breastfeeding Among Infants Born at an Inner-City US Baby-Friendly Hospital. *Journal of Human Lactation* 2007; 23:157-64.
47. Heidari Z, Keshvari M, Kohan S. Breastfeeding promotion, challenges and barriers: a qualitative research. *International Journal of Pediatrics* 2016;4(5):1687-95.
48. Sohrabi Z, Momenzadeh F, Aemmi SZ, Tabibi M, Musavi Z, Savabi M. Socio-demographic and lifestyle factors in breastfeeding mothers, referring to Isfahan Health Centers. *International Journal of Pediatrics* 2016;4(2):1331-7.
49. Haghighi M, Taheri E. Factors associated with breastfeeding in the first hour after birth, in baby friendly hospitals, Shiraz, Iran. *International Journal of Pediatrics* 2015;3(5.1):889-96.
50. World Health Organization. Global Strategy for Infant and Young Child Feeding. 54th World Health Assembly (Report no. A54/INF. DOC./4). Geneva, World Health Organization, 2001.
51. Lee WT, Wong E, Lui SS, Chan V, Lau J. Decision to breast feed and early Cessation of breast feeding in infants below 6 months old – a population-based study of 3,204 infants in Hong kong. *Asia Pac J Clin Nutr* 2007;16(1):163-71.
52. Ruowei Li, Zhen Z, Mokdad A, Barker L, Grummer Strawn L. Prevalence of Breastfeeding in the United States: The 2001 National Immunization Survey. *Pediatrics* 2003; 111: 1198- 1201.
53. Rashidian A, Khosravi A, Arab M, Beheshtian M, Khabiri R, Karimi-Shahanjarini A, et al. IrMIDHS - I.R. Iran Multiple-Indicator and Demographic Health Survey 2010: questionnaires, guides and protocols. Tehran: National Institute of Health Research and Deputy for Health, Ministry of Health and Medical Education; 2011.

Table-1: DALYs rate [95%UI] per 100,000 population attributed to suboptimal breastfeeding in Iran in different age and sex groups from 1990 to 2010					
Variables	1990 Rate [95% UI]	1995 Rate [95% UI]	2000 Rate [95% UI]	2005 Rate [95% UI]	2010 Rate [95% UI]
Both Genders					
Under 5 years	4345[2058,7444]	2623[1186,4604]	1742[748,2986]	1658[742,2846]	1356[564,2502]
Neonatal	63571[29321,109129]	37490[17245,62784]	21112[9286,35368]	17380[7274,30337]	14599[5359,27474]
Post Neonatal	19396[8604,34807]	12487[5318,23159]	8064[3298,14676]	7222[2958,13201]	5798[2236,11457]
1 to 4 years	146[11,445]	121[9,347]	117[9,343]	124[10,335]	123[11,325]
All Ages	725[343,1241]	344[156,604]	171[73,293]	140[63,241]	113[47,208]
Age standardized	402[191,689]	257[116,453]	166[71,285]	148[66,254]	121[50,224]
Girl					
Under 5 years	4242[1909,7444]	2591[1154,4596]	1780[779,3071]	1715[778,3076]	1397[561,2773]
Neonatal	59913[25709,105612]	35724[16868,63845]	20591[9165,36111]	17462[6349,33554]	14625[5029,32594]
Post Neonatal	19015[7852,35892]	12386[5003,23040]	8299[3460,14935]	7514[3081,14097]	6006[2207,13159]

1 to 4 years	154[13,46]	125[9,375]	120[9,34]	126[10,345]	126[11,347]
All Ages	698[314,1226]	335[149,594]	172[75,297]	144[66,259]	115[46,228]
Age standardized	393[177,687]	254[113,452]	169[73,293]	153[69,275]	125[50,248]
Boy					
Under 5 years	4444[1988,7957]	2654[1154,4891]	1706[720,3160]	1603[628,2973]	1316[499,2657]
Neonatal	67072[27885,130272]	39179[15583,74449]	21608[8533,39942]	17301[6718,31987]	14575[4614,31974]
Post Neonatal	19762[8119,3713]	12584[4964,24930]	7840[3161,15504]	6943[2589,13747]	5599[1915,12339]
1 to 4 years	139[10,429]	117[9,357]	114[9,340]	122[10,327]	121[11,322]
All Ages	750[336,1343]	353[153,651]	170[72,315]	136[53,252]	110[42,222]
Age standardized	412[184,738]	260[113,481]	162[68,301]	143[56,266]	118[45,237]

Table-2: Deaths rate [95% UI] per 100,000 population attributed to suboptimal breastfeeding in Iran in different age and sex groups from 1990 to 2010					
Variables	1990 Rate [95% UI]	1995 Rate [95% UI]	2000 Rate [95% UI]	2005 Rate [95% UI]	2010 Rate [95% UI]
Both Genders					
Under 5 years	50[23,86]	30[13,53]	19[8,34]	18[7,32]	14[6,28]
Neonatal	735[339,1266]	432[195,727]	241[105,409]	198[81,349]	166[59,317]
Post Neonatal	224[98,404]	143[60,266]	91[37,167]	81[31,151]	64[23,129]
1 to 4 years	1[0,4]	1[0,3]	1[0,2]	1[0,2]	0[0,1]
All Ages	8[4,14]	4[2,7]	2[1,3]	2[1,3]	1[0,2]
Age standardized	5[2,8]	3[1,5]	2[1,3]	2[1,3]	1[0,2]
Girls					
Under 5 years	48[22,85]	29[13,52]	20[8,35]	19[8,34]	15[6,31]
Neonatal	693[294,1223]	411[192,738]	235[103,416]	199[71,386]	166[55,376]
Post Neonatal	219[88,415]	142[56,265]	94[37,171]	84[33,162]	67[23,150]
1 to 4 years	1[0,4]	1[0,3]	1[0,2]	1[0,2]	0[0,1]
All Ages	8[4,14]	4[2,7]	2[1,3]	2[1,3]	1[0,3]
Age standardized	4[2,8]	3[1,5]	2[1,3]	2[1,3]	1[1,3]

Boys					
Under 5 years	51[22,92]	30[12,56]	19[8,36]	17[7,33]	14[5,29]
Neonatal	776[321,1509]	452[175,860]	247[96,460]	197[75,368]	166[50,368]
Post Neonatal	228[92,430]	144[56,288]	88[34,176]	78[28,157]	62[19,140]
1 to 4 years	1[0,4]	1[0,3]	1[0,2]	1[0,2]	1[0,2]
All Ages	9[4,15]	4[2,7]	2[1,4]	1[1,3]	1[0,2]
Age standardized	5[2,8]	3[1,6]	2[1,3]	2[1,3]	1[0,3]

Table-3: YLDs rate [95%UI] per 100000 population attributed to suboptimal breastfeeding in Iran in different age and gender groups from 1990 to 2010					
Variables	1990 Rate [95% UI]	1995 Rate [95% UI]	2000 Rate [95% UI]	2005 Rate [95% UI]	2010 Rate [95% UI]
Both Genders					
Under 5 years	78[26,174]	83[27,189]	100[33,229]	113[38,257]	120[38,268]
Neonatal	333[110,684]	333[113,643]	347[114,660]	337[122,654]	337[121,635]
Post Neonatal	210[73,441]	222[81,446]	249[90,502]	263[99,522]	269[109,538]
1 to 4 years	45[3,134]	51[4,155]	65[5,199]	75[6,203]	82[6,231]
All Ages	13[4,29]	11[4,25]	10[3,22]	10[3,22]	10[3,22]
Age standardized	7[2,16]	8[3,17]	9[3,20]	10[3,23]	11[3,24]
Girls					
Under 5 years	81[24,182]	86[27,193]	104[33,237]	117[35,261]	124[37,296]
Neonatal	335[102,691]	338[107,682]	354[114,708]	344[122,695]	343[114,685]
Post Neonatal	212[73,458]	225[80,472]	254[89,532]	268[98,567]	275[100,552]
1 to 4 years	48[4,144]	54[4,165]	68[5,208]	79[6,208]	87[7,255]
All Ages	13[4,30]	11[4,25]	10[3,23]	10[3,22]	10[3,24]
Age standardized	7[2,16]	8[3,17]	9[3,21]	10[3,23]	11[3,26]
Boys					

Under 5 years	76[24,172]	80[26,185]	97[31,225]	109[38,253]	116[36,253]
Neonatal	330[99,707]	328[104,651]	340[105,659]	331[111,678]	331[119,669]
Post Neonatal	208[67,460]	218[76,445]	244[86,495]	258[92,528]	264[97,552]
1 to 4 years	43[3,135]	49[4,145]	62[5,189]	72[6,208]	78[6,219]
All Ages	13[4,29]	11[3,25]	10[3,22]	9[3,21]	10[3,21]
Age standardized	7[2,15]	7[2,17]	9[3,20]	10[3,22]	10[3,23]

Table-4: YLLs rate [95% UI] per 100000 population attributed to suboptimal breastfeeding in Iran in different age and sex groups from 1990 to 2010					
Variables	1990 Rate [95% UI]	1995 Rate [95% UI]	2000 Rate [95% UI]	2005 Rate [95% UI]	2010 Rate [95% UI]
Both Genders					
Under 5 years	4267[1998,7357]	2540[1142,4556]	1642[692,2893]	1545[627,2739]	1236[475,2379]
Neonatal	63239[29117,108847]	37157[16750,62509]	20765[9069,35137]	17042[6967,29998]	14263[5082,27241]
Post Neonatal	19187[8451,34653]	12266[5143,22843]	7815[3171,14355]	6959[2633,13009]	5529[1995,11097]
1 to 4 years	101[8,306]	69[5,214]	52[4,159]	49[4,140]	41[4,121]
All Ages	711[333,1227]	333[150,597]	161[68,284]	131[53,232]	103[39,197]
Age standardized	395[185,681]	250[112,448]	156[66,276]	138[56,245]	111[42,213]
Girls					
Under 5 years	4161[1847,7292]	2504[1089,4501]	1676[688,2980]	1598[676,2952]	1272[485,2641]
Neonatal	59578[25257,105218]	35386[16481,63496]	20238[8864,35770]	17118[6086,33192]	14282[4702,32312]
Post Neonatal	18803[7595,35657]	12161[4785,22760]	8044[3148,14644]	7246[2825,13907]	5732[1992,12871]
1 to 4 years	106[9,333]	71[5,226]	52[4,158]	47[4,144]	39[3,120]
All Ages	685[304,1201]	324[141,582]	162[67,288]	135[57,248]	105[40,218]
Age standardized	385[171,675]	246[107,442]	160[65,284]	143[60,264]	114[43,236]

Boys					
Under 5 years	4369[1928,7867]	2573[1071,4829]	1609[647,3075]	1493[567,2837]	1201[422,2510]
Neonatal	66742[27583,129810]	38851[15092,73980]	21268[8234,39586]	16970[6416,31679]	14244[4296,31654]
Post Neonatal	19554[7857,36919]	12366[4769,24705]	7596[2925,15143]	6685[2379,13519]	5335[1656,12014]
1 to 4 years	96[7,307]	68[5,217]	52[4,166]	50[4,139]	43[4,129]
All Ages	737[325,1328]	342[142,642]	160[64,306]	127[48,241]	100[35,210]
Age standardized	405[179,729]	253[104,476]	153[62,293]	133[51,253]	107[38,224]