

Autosomal Recessive Hypohidrotic Ectodermal Dysplasia Caused by a Novel Mutation in *EDAR* Gene

Nader Ebadi¹, Sepehr Javadi¹, TayyebAli Salmani¹, Mohammad Miryounesi², Vahid Reza Yassae², *Soudeh Ghafouri-Fard¹

¹Department of Medical Genetics, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

²Genomic Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Abstract

Backgrounds

Hypohidrotic ectodermal dysplasia (HED) is a rare genetic disorder, distinguished by hypotrichosis, hypohidrosis, and hypodontia. HED can be inherited in X-linked recessive manner as a result of mutations in the *ectodysplasin A (EDA)* gene as well as autosomal dominant and autosomal recessive manners both of them caused by mutations in *EDA receptor (EDAR)* and *EDAR-associated death domain (EDARADD)* genes.

Results

In this report, we investigated a consanguineous Iranian family with autosomal recessive form of HED. A homozygous missense mutation was detected in exon 1 of *EDAR* gene in the proband (c.278C>G) resulting in p.C93S that alters the sequence of the EDAR protein.

Conclusion

We facilitated the effective genetic counseling and prenatal diagnosis in this family through detection of the disease causing mutation.

Key Words: Ectodermal dysplasia, EDAR, Mutation.

*Please cite this article as: Ebadi N, Javadi S, Salmani TA, Miryounesi M, Yassae VR, Ghafouri-Fard S. Autosomal Recessive Hypohidrotic Ectodermal Dysplasia Caused by a Novel Mutation in EDAR Gene. Int J Pediatr 2018; 6(1): 6899-6902. DOI: **10.22038/ijp.2017.28529.2480**

*Corresponding Author:

Soudeh Ghafouri-Fard, Department of Medical Genetics, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Email: s.ghafourifard@sbmu.ac.ir

Received date: Oct.23, 2017; Accepted date: Nov.22, 2017

1- INTRODUCTION

Hypohidrotic ectodermal dysplasia (HED) is a rare genetic disorder, distinguished by hypotrichosis (abnormal development of scalp and body hair), hypohidrosis (Impaired sweating), and hypodontia (congenital absence of teeth) (1, 2). HED is mostly inherited in X-linked recessive manner (OMIM: 305100) as a result of mutations in the *ectodysplasin A (EDA)* gene (3). However, HED can be inherited in autosomal dominant (OMIM: 129490), and autosomal recessive manners (OMIM: 224900) both of them caused by mutations in *EDA receptor (EDAR)* and *EDAR-associated death domain (EDARADD)* genes (4, 5).

Proteins that encoded by these three genes are essential for the activation of the Nuclear factor- κ B (NF- κ B) signaling pathway, a necessary pathway for normal development of ectodermal organs both in humans and in mice (6). Moreover, mutation in two other genes named *WNT10A* and *TRAF6* have been shown to cause HED (7).

2- CASE REPORT

In this report, we investigated a consanguineous Iranian family with autosomal recessive form of HED. The proband was a male child aged 7 years with light-colored, brittle, and slow-growing hair. Eyebrows were scanty and skin was dry and scaly. He suffered from hypohidrosis and hypodontia. The teeth were small and pointed. He showed a characteristic facial dysmorphology including a prominent forehead, a flattened bridge of the nose, and thick lips (**Figure.1**). Other features include thin, dark-colored skin around the eyes and chronic skin problems such as eczema. In order to find the underlying genetic cause, genomic DNA was isolated from blood sample of the patient after informed consent using the standard salting out technique.

Whole exome sequencing was performed in Laboratory of Molecular Diagnosis, University of Leuven using Illumina sequencer. A homozygous missense mutation was detected in exon 1 of *EDAR* gene (c.278C>G) resulting in p.C93S that alters the sequence of the EDAR protein. This mutation has not been reported in generalist polymorphism databases (ExaC or exome variant server [EVS]) or the 1000 genome database (<http://www.1000genomes.org/1000-genomes-browsers>). *In silico* functional analysis of the sequencing results by Polyphen -2 (Polymorphism Phenotyping v2) showed that it is possibly damaging.

In addition, Combined Annotation Dependent Depletion (CADD) which applies a uniform, genome-wide, variant scoring metric (C-score) through combining Sorting Intolerant From Tolerant (SIFT) method and PolyPhen to predict the pathogenicity of any variant (8), showed that this nucleotide change is deleterious with a score of 27. The mutation was confirmed by Sanger DNA sequencing in patients. Segregation analysis showed that parents were heterozygous for the detected mutation.

3- DISCUSSION

In this study we detected a novel homozygous missense mutation in *EDAR* gene in an Iranian consanguine family affected to autosomal recessive HED. *EDAR* belongs to tumor necrosis factor receptor superfamily (TNFRSF) and is a transmembrane protein with three cysteine-rich domain (CRD) in the ectodomain. These CRDs are rich in cysteines, are constrained by disulfide bridges and are involved in the receptor-ligand interaction (1, 2). According to Human Gene Mutation Database (HGMD) (www.hgmd.cf.ac.uk/), 42 different HED-causing mutations in *EDAR* gene have been reported up to now, 36 of which are missense mutations

(<http://www.hgmd.cf.ac.uk/ac/gene.php?gene=EDAR>). The detected mutation in the present study (c.278G>C) is a missense mutation in CRD2 that disrupts a disulfide bridge existing between cys93 and cys113 (<http://www.uniprot.org/uniprot/Q9UNE0>) and most probably alters the protein's secondary structure and /or the ligand

binding interaction site resulting in impairment of NF- κ B signaling which is important in the development of ectoderm. Interestingly, two other missense mutation in cys87 and cys113 has been reported in CRD2 disrupting its two disulfide bridges and causing HED phenotype (3, 4).




Fig. 1: The pedigree of family with hypohidrotic ectodermal dysplasia (HED).

4- CONCLUSION

In conclusion, we have reported the sixth missense mutation in CRD2 and added to *EDAR* mutation repository. Identification of underlying genetic cause of inherited disorders would facilitate preimplantation genetic diagnosis as well as prenatal diagnosis. In addition, detection of carrier status of parents enables genetic counselor to provide recurrence risk in each family.

5- CONFLICT OF INTEREST: None.

6- ACKNOWLEDGEMENT

The authors are grateful to the patient's family for their participation in the study.

7- REFERENCES

1. Zeng B, Xiao X, Li S, Lu H, Lu J, Zhu L, et al. Eight Mutations of Three Genes (EDA, EDAR, and WNT10A) Identified in Seven Hypohidrotic Ectodermal Dysplasia Patients. *Genes (Basel)*. 2016;7(9):65.
2. Bluschke G, Nusken KD, Schneider H. Prevalence and prevention of severe complications of hypohidrotic ectodermal dysplasia in infancy. Early human development. 2010 Jul;86(7):397-9. PubMed PMID: 20682465. Epub 2010/08/05. eng.
3. Kere J, Srivastava AK, Montonen O, Zonana J, Thomas N, Ferguson B, et al. X-linked anhidrotic (hypohidrotic) ectodermal dysplasia is caused by mutation in a novel transmembrane protein. *Nature genetics*. 1996 Aug;13(4):409-16. PubMed PMID: 8696334. Epub 1996/08/01. eng.
4. Monreal AW, Ferguson BM, Headon DJ, Street SL, Overbeek PA, Zonana J. Mutations in the human homologue of mouse *dl* cause autosomal recessive and dominant hypohidrotic ectodermal dysplasia. *Nature genetics*. 1999 08//print;22(4):366-9.
5. Headon DJ, Emmal SA, Ferguson BM, Tucker AS, Justice MJ, Sharpe PT, et al. Gene defect in ectodermal dysplasia implicates a death domain adapter in development. *Nature*. 2001 Dec 20-27;414(6866):913-6. PubMed PMID: 11780064. Epub 2002/01/10. eng.
6. Laurikkala J, Pispala J, Jung HS, Nieminen P, Mikkola M, Wang X, et al. Regulation of hair follicle development by the TNF signal ectodysplasin and its receptor Edar. *Development (Cambridge, England)*. 2002 May;129(10):2541-53. PubMed PMID: 11973284. Epub 2002/04/26. eng.
7. Bohring A, Stamm T, Spaich C, Haase C, Spree K, Hehr U, et al. WNT10A mutations are a frequent cause of a broad spectrum of ectodermal dysplasias with sex-biased manifestation pattern in heterozygotes. *American journal of human genetics*. 2009 Jul;85(1):97-105. PubMed PMID: 19559398. Pubmed Central PMCID: Pmc2706962. Epub 2009/06/30. eng.
8. Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. *Nature genetics*. 2014 Mar;46(3):310. PubMed PMID: WOS:000332036700017. English.