

Case Report (Pages: 7569-7573)

# Abdominal Splenosis: An Unusual Cause of Bowel Obstruction in a 7-Year-Old Child

Mohamed Zouari<sup>1</sup>, \*Hamdi Louati<sup>1</sup>, Senda Houidi<sup>1</sup>, Mahdi Ben Dhaou<sup>1</sup>, Mohamed Jallouli<sup>1</sup>, Riadh Mhiri<sup>1</sup>

<sup>1</sup>Department of Pediatric Surgery. Hedi Chaker Hospital. 30219 Sfax, Tunisia.

#### Abstract

Splenosis is a rare condition defined as a heterotopic auto-transplantation of splenic tissue, typically after blunt abdominal trauma. It can occur anywhere in the abdominal cavity, or even the chest. Splenosis found in clinical practice is relatively rare because most patients are asymptomatic. We report a rare case of abdominal splenosis in a 7-year-old boy, 3 years after an abdominal blunt trauma.

Key Words: Bowel obstruction, Children, Splenosis; Surgery, Trauma.

<u>\*Please cite this article as</u>: Zouari M, Louati H, Houidi S, Ben Dhaou M, Jallouli M, Mhiri R. Abdominal Splenosis: An Unusual Cause of Bowel Obstruction in a 7-Year-Old Child. Int J Pediatr 2018; 6(5): 7569-73. DOI: **10.22038/ijp.2018.30404.2673** 

\*Corresponding Author:

Hamdi Louati, Department of Pediatric Surgery, Hedi Chaker Hospital, 30219 Sfax, Tunisia.

Email: drhamdilouati85@yahoo.com

Received date: Jan.22, 2018; Accepted date: Feb.22, 2018

### **1- INTRODUCTION**

Splenosis is a rare condition defined as a heterotopic auto-transplantation of splenic tissue, typically after blunt abdominal trauma (1). It can occur anywhere in the abdominal cavity, or even the chest (2, 3). Splenosis seeds onto exposed vascularized peritoneal surfaces following splenic trauma or surgery. However, splenosis found in clinical practice is relatively rare because most patients are asymptomatic (4).We report a rare case of abdominal splenosis in a 7-year-old boy, 3 years after an abdominal blunt trauma.

### 2- CASE REPORT

A 7-year-old boy, with a history of a trauma to the left upper quadrant at the age of four, presented to the pediatric emergency department with a 2-day

history of abdominal pain and vomiting. On physical examination, the patient was lethargic, his temperature was 37.9°C and his abdomen was distended. Vital signs were normal. Plain abdominal radiograph showed air-fluid levels (**Figure.1**). Computed tomography (CT) scan findings were suggestive of mechanical small bowel obstruction. However, the cause of this obstruction was not detected.

Urgent exploratory laparotomy was exploration performed. Upon and adhesiolysis, a well-contoured, dark blue mass was found in the small bowel mesentery in tight contact with the rectum and the abdominal wall. This mass was resected (Figure.2). The pathological examination of the operative specimen revealed a normal splenic tissue, consistent with a diagnosis of splenosis.



Fig.1: Plain abdominal radiograph showing air-fluid levels.



Fig.2: Intraoperative view of the resected mass.

### **3- DISCUSSION**

Splenosis is a benign condition among patients with a history of splenic trauma or surgery (4-6). The majority of splenosis cases occur after traumatic splenectomies, whereas a few are the result of elective splenectomies done for hematological diseases (6). The actual incidence of splenosis is not known due to the incidental nature of the diagnosis (7). Splenosis is not a rare disease. Abdominal splenosis has been shown to occur in 16-17% of patients who undergo elective splenectomy for a hematological disease, and in 44-76% of patients with a posttraumatic splenectomy due to spillage after traumatic insult (8).

Splenic fragments could heterotopically auto transplant onto vascularized intra or extra peritoneal surfaces, deriving its blood supply from adjacent tissues and organs (2, 9). Splenic fragments usually spread by a direct seeding process onto adjacent surfaces, mainly the small intestine serosa, the greater omentum, the parietal peritoneum, the undersurface of the diaphragm, and the pancreatic tail (10). A hematological spread of splenic fragments to distant organs such as the liver, the breast, and the brain has also been described (11). Most of patients with splenosis are asymptomatic and splenic deposits are usually found incidentally on imaging or at laparotomy (4, 12). However, splenosis may be implicated in recurrent episodes of abdominal pain, small bowel obstruction, and bleeding by invasion into the gut or its own rupture (7, 13). Standard CT and Magnetic resonance imaging (MRI scans could be able to identify and describe the anatomical locations of these deposits (13).

However, these imaging tools are of limited value and are nonspecific in this entity. Splenosis could mimic malignancy on imaging posing a diagnostic dilemma. Standard CT and MRI scans cannot distinguish splenosis nodules from other conditions such as metastatic disease, pheochromocytoma, lymphoma, carcinomatosis, hepatic or renal malignancy, endometriosis, or simple adenopathy (14, 15). In the present case, CT scan was not helpful for the diagnosis of abdominal splenosis. Technetium-99mlabelled erythrocytes scintigraphy has been demonstrated to be the most sensitive and specific diagnostic technique for detecting ectopic splenic tissue (15-17). When imaging findings alone do not provide a definitive diagnosis of splenosis, surgery is needed to reach the final diagnosis (17). The treatment of splenosis depends on the patient's symptoms (18). Asymptomatic splenosis pose no risk and the resection of splenic deposits is not required (6). required Surgery is in cases of symptomatic or complicated splenosis, in patients with hematological disease for whom splenectomy is beneficial, and when scintigraphic methods are not available (19). At present, the development of laparoscopic surgical techniques, provides a port for minimally invasive entry for the visualization of suspected abdominal masses, and allows access for eventual biopsy or resection (20).

### **4- CONCLUSION**

Abdominal Splenosis should be included as a potential source for abdominal pain or bowel obstruction in splenectomized patients or those with abdominal blunt trauma to prevent unnecessary surgeries.

#### 5- CONFLICT OF INTEREST: None.

## **6- REFERENCES**

1. Gupta K, Ahluwalia A, Jain T, Saggar K. Abdominal splenosis mimicking peritoneal deposits - A case report. Pan Afr Med J 2014;17:269.

2. Akay S, Ilica AT, Battal B, Karaman B, Guvenc I. Pararectal mass: An atypical location of splenosis. J Clin Ultrasound 2012;40(7):443-7.

3. Baldolli A, Coeuret S, Le Pennec V, Agostini D, Verdon R. Thoracic splenosis mimicking a pleuropneumonia: A case report. Medicine (Baltimore) 2017;96(29):e7552.

4. He ZL, Xu X, Ke QH, Zheng SS. Successful diagnosis of intrahepatic splenosis mimicking hepatic tumor. Kaohsiung J Med Sci 2016; 32: 224-5.

5. Neufeld EA, Sekhon S, Gerscovich EO. Intra-renal splenosis mimicking a solid renal mass. Ultrasound 2017; 25(3):173-6.

6. Younan G, Wills E, Hafner G. Splenosis: A Rare Etiology for Bowel Obstruction-A Case Report and Review of the Literature. Case Rep Surg 2015;2015:890602.

7. Carman KA, Walters BS, Kennedy AP. Abdominal splenosis. J Pediatr Surg Case Rep 2018 Jan 9. doi.org/10.1016/j.epsc.2018.01.006.

8. Nicolas G, Schoucair R, Shimlati R, Rached L, Khoury G. Laparoscopic gastric band removal complicated by splenosis. Clin Case Rep 2016;4(8):807-11.

9. Sikov WM, Schiffman FJ, Weaver M, Dyckman J, Shulman R, Torgan P. Splenosis presenting as occult gastrointestinal bleeding. Am J Hematol 2000;65(1):56-61.

10. Lin WC, Lee RC, Chiang JH, et al. MR features of abdominal splenosis. AJR Am J Roentgenol 2003;180(2):493-6.

11. Kwok CM, Chen YT, Lin HT, Su CH, Liu YS, Chiu YC. Portal vein entrance of splenic erythrocytic progenitor cells and local hypoxia of liver, two events cause intrahepatic splenosis. Med Hypotheses 2006; 67(6):1330-2.

12. Gincu V, Kornprat P, Thimary F, Jahn S, Mischinger HJ. Intestinal obstruction caused by splenosis at the rectosigmoid junction, mimicking malignant pelvic tumor. Endoscopy 2011;43 Suppl 2 UCTN:E260.

13. Lake ST, Johnson PT, Kawamoto S, et al. CT of splenosis: patterns and pitfalls. AJR Am J Roentgenol 2012; 199(6):W686-93.

14. Deng Y, Jin Y, Li F. Splenosis mimicking an extramural duodenal mass: A case report. Oncol Lett 2014;8(6):2811-3.

15. Felice M, Tourojman M, Rogers C. Right retroperitoneal splenosis presenting as an adrenal mass. Urol Case Rep. 2017; 16: 44-5.

16. Alang N. Splenosis: an unusual cause of massive gastrointestinal bleeding. R I Med J 2013; 96(11):48-9.

Nikam and Nagure

17. Tordjman M, Eiss D, Dbjay J, et al. Renal Pseudo-Tumor Related to Renal Splenosis: Imaging Features. Urology 2018 Jan 26. pii: S0090-4295(18)30062-1.

18. Gheonea IA, Săndulescu SM, Firulescu SC, Tudorașcu DR, Ciobanu MO, Badea O, et al. A rare case of ovarian splenosis. Rom J Morphol Embryol. 2016;57(2 Suppl):811-6.

19. Ksiadzyna D, Peña AS. Abdominal splenosis. Rev Esp Enferm Dig 2011;103(8):421-6.

20. Barbaros U, Dinccag A, Kabul E. Minimally invasive surgery in the treatment of splenosis. Surg Laparosc Endosc Percutan Tech 2006;16:187-9.