

## Psychotherapy for Postpartum Depression in Iranian Women: A Systematic Review and Meta-Analysis

Masumeh Ghazanfarpour<sup>1</sup>, Fatemeh Rajab Dizavandi<sup>2</sup>, Leila Kargarfard<sup>3</sup>, Elahe Heidari<sup>4</sup>,  
Talat Khadivzadeh<sup>1</sup>, \*Masumeh Saeidi<sup>5</sup>, Somayeh Abdollahian<sup>6</sup>, Minoos Safaei<sup>1</sup>

<sup>1</sup>Evidence - Based Care Research Center, Mashhad University of Medical Sciences, Mashhad, Iran.

<sup>2</sup>Faculty Member, Department of Community Health and Psychiatric Nursing, School of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, Iran. <sup>3</sup>Instructor of Fatemeh School and Midwifery, Shiraz University of Medical Sciences, Iran. <sup>4</sup>Assistant Professor, Department of Pediatrics, Faculty of Medicine, Mashhad university of Medical Sciences, Mashhad, Iran. <sup>5</sup>Student Research Committee, Department of Medical Education, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran. <sup>6</sup>Student Research Committee, Department of Midwifery and Reproductive Health, Nursing and Midwifery School, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

### Abstract

**Background:** Postpartum depression (PPD) is a type of disorder could have serious effects on the mother, the baby, and other family members, given the contradictory results of the previous studies about the effect of education programs, aim of the present study was to evaluate the effect of educational programs on postpartum depression in Iranian women.

**Materials and Methods:** English electronic information databases such as Medline (via PubMed), Scopus, ISI Web of Science, EMBASE, and Cochrane Library were searched until February 2018. In addition, Iran doc, Barakatskns, Magiran, Medlib, SID, and google scholar were searched using equivalent keywords in Persian until February 2018. Cochrane Q test ( $p < 0.05$  as statistically significant), and I<sup>2</sup> index were used to evaluate heterogeneity. Funnel plot and Egger's regression were conducted to detect publication bias.

**Results:** The finding of the Meta-analysis indicate that education interventions was found to be more effective and could significantly improve postpartum depression (Standard Mean deviation (SMD) =1.44;  $P < 0.001$ ; Confidence Interval [CI] =0.595 to 2.40; Heterogeneity  $p < 0.001$ ; I<sup>2</sup>=87%). The funnel plot appears asymmetric with an outlier study. However, this finding was not confirmed by Egger's test that was nonsignificant ( $p = 0.262$ ), and show no asymmetry. The level of depression was lower in women receiving problem-solving skills in compared to who received no intervention (SMD =1.83;  $P < 0.001$ ); the level of depression showed a significant borderedline decline in women receiving Mindfulness-based Cognitive Therapy compared to those who did not receive intervention (SMD =1.20;  $P = 0.089$ ).

**Conclusion:** Based on the finding of the current meta-analysis, it can be said that educational programs that teaches Mindfulness-based cognitive and problem-solving skills can improve of postpartum depression.

**Key Words:** Iran, Depression, Educational Program, Meta-analysis, Postpartum, Women.

\*Please cite this article as: Ghazanfarpour M, Rajab Dizavandi F, Kargarfard L, Heidari E, Khadivzadeh T, Saeidi M, et al. Psychotherapy for Postpartum Depression in Iranian Women: A Systematic Review and Meta-Analysis. Int J Pediatr 2018; 6(6): 7781-90. DOI: [10.22038/ijp.2018.30682.2690](https://doi.org/10.22038/ijp.2018.30682.2690)

### Corresponding Author:

Masumeh Saeidi, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

E-mail: Masumeh\_Saeidi@yahoo.com

Received date: Jan.17, 2018; Accepted date: Feb.22, 2018

## 1- INTRODUCTION

Postpartum depression (PPD) is a type of disorder that affects some women within the first 2 to 6 weeks following childbirth. Symptoms may include crying episodes, despondency, mood swings, feelings of worthlessness and inability to perform maternal role (1-3). Postpartum depression is a common disorder, affecting 10 to 30% of childbearing women (4-6). In a meta-analysis, a total of 41 studies met the inclusion criteria. The pooled prevalence of PPD in Iran was reported to be 25.3% (95% Confidence Interval [CI]: 22.7% -27.9%)(7). Psychological problems such as PPD account for more than 12.5% of referrals to women's healthcare centers (8). The cause of PPD has not been specifically determined so far (9-12). Some researchers consider this disorder as a response to hormonal imbalance after childbirth, particularly the loss of estrogen, estradiol, tryptophan, and endorphin, as well as elevated levels of prolactin and cortisol (1, 13, 14).

Other possible reasons could include disturbed adjustment of sleep and diet, and the baby's helplessness (1, 15, 16). Postpartum depression could have serious effects on the mother, the baby, and other family members, which leads to resultant complications for the mother-child relationship, and adversely affects the growth and development of the infant (5, 17-20). It can also cause behavioral, social and emotional problems in such children. In some severe cases of this disorder, depressed mother has been reported to even commit infanticide. In fact, the PPD has a significant negative impact on all aspects of the quality of life of the mothers. Women affected by the PPD may experience different problems in their social activities, individual performance, and household chores, exerting a negative influence on the mood of other family members. According to previous studies, depressed women's spouses are often more

likely to develop depression as well, which increases marital conflicts (16, 21-26). Studies and evidences demonstrate that psychological and educational interventions are effective in reducing postpartum depression, and can improve the quality of life of the mothers without having any negative influence or complication (1). Given the contradictory results of the previous studies about the effect of education programs on the postpartum depression, the inadequacy of the educations available for the mothers, the high prevalence of this disorder in Iran, and the adverse effects of depression on health of the mothers, of the family and of society, we aimed to conduct the present meta-analysis in this regard. The aim of the present study was to evaluate the effect of educational programs on postpartum depression.

## 2- MATERIALS AND METHODS

### 2-1. Method

To perform the systematic review, English electronic information databases such as Medline (via PubMed), Scopus, ISI Web of Science, EMBASE, and Cochrane Library were searched without publication date restrictions until February 2018, 2018. Following keywords were used to identify articles related to the effect of education program on postpartum depression (PPD). English keywords were used to search databases: "Depression", "Intervention", "Iran", "Postpartum" "Psychotherapy", and "Women".

In addition, IranDoc, Barakatks, Magiran, Medlib, Google Scholar, Magiran, and SID were searched using equivalent keywords in Persian till to February 2018. Also, the references of the relevant articles were investigated to find further related articles. Two authors independently reviewed the titles and abstracts of the articles. If the subject was relevant, the full text article was extracted, and reviewed by the

authors. Finally, articles meeting inclusion criteria were evaluated.

### 2-2. Inclusion criteria

Studies were included into systematic review if they had the following criteria:

- Experimental study (Quasi-experimental or clinical trials),
- Reported severity of postpartum depression with validated instrument tool,
- Used a psychotherapeutic intervention.

### 2-3. Data extraction

Two authors independently extracted following data based on a pre-defined checklist designed by authors: name of first author, year of publish, region of study (city), Measurement tool, whether randomization was done or not, whether randomization technique was suitable or not, a number of participants in intervention and control group, level of depression complaints, age, comparability of the treatment and control groups, design, and major relevant findings.

### 2-4. Quality assessment of the included studies

Studies quality was by two reviewers assessed using the Oxford Center for Evidence-based Medicine checklist for therapeutic studies (<http://www.cebm.net/index.aspx?o=5653>. Accessed in March 2013).

### 2-5. Statistical analyses

The main effect size was calculated as standardized difference in means (SMD). We reported the results based on a random effects model because high heterogeneity across studies. Cochrane Q test ( $p < 0.05$  as statistically significant), and  $I^2$  index were used to evaluate heterogeneity. We also conducted a funnel plot and Egger's regression to detect publication bias.

## 3- RESULTS

**Table.1** shows the characteristic of five studies included in current study (*Please see the end of paper*). Process of selection of included studies in the education interventions met analysis was shown in **Figure.1**. The finding of the Meta-analysis indicate that education interventions was found to be more effective and could significantly improve postpartum depression (Standard Mean deviation (SMD) =1.49;  $P=0.001$ ; 95%CI= 0.595 to 2.40, at five trial studies (1, 8, 27-29). However, Heterogeneity across studies were very high ( $p < 0.001$ ;  $I^2=87\%$ ). The funnel plot is shown in **Figure.2**.

We performed sensitivity analysis to detect potential source (outlier study) of heterogeneity. However, heterogeneity continued to remain after excluding one by one study. Also, sensitivity analysis indicated the robust of meta-analysis by showing similar SMD before and after excluding one by one study. We also performed subgroup analysis by type of intervention because of high heterogeneity.

At current systematic review, two studies (Parsa et al. and Nasiri et al.) assessed the effect of problem-solving skills on postpartum depression. Level of depression was lower in women receiving problem-solving skills in compared to who received no intervention (SMD =1.83;  $P < 0.001$ ; 95% CI= 1.43 to 2.33) (1, 28).

Heterogeneity was 0% among studies that assessed effect of Problem-solving skills on postpartum depression. In Ghadam pour et al. and Abad et al.'s studies, level of depression showed a significant borderline decline in women receiving Mindfulness-based Cognitive Therapy incompared to those who did not receive intervention (SMD =1.20;  $P=0.089$ ; 95% CI= 2.58 to 1.73) (17, 29). Heterogeneity was  $I^2=83\%$  among studies that assessed effect of Mindfulness-based cognitive on postpartum depression. Subgroup analysis

showed that typ of intervention (Mindfulness-based cognitive Therapy vs. Problem-solving skills) might be the source of heterogeneity. Naddap et al. conducted a Quasi-experimental. They divided subjects into two equal groups to receive Group psychotherapy (n=15) and control group (n=15). Post partum depression improved significantly in group psychotherapy compared to control group (27). Pooled effect sizes were larger in

studies assessed the effect of Problem-solving skill traning compared to Mindfulness –basees cognitive therapy ( $p < 0.001$ ). We made a funnel plot to evaluate publication bias (Figure.5). The funnel plot appears asymmetric with an outlier study. However, this finding was not confirmed by Egger's test that was non-significant ( $p = 0.262$ ), and show no asymmetry.






Fig.1: Flowchart of included studies.




Meta Analysis

**Fig.2:** Effects of educational programs on postpartum depression. The horizontal lines denote the 95% confidence interval, ■ point estimate (size of the square corresponds to its weight); ♦, combined Overall effect of treatment.




Meta Analysis

**Fig.3:** Effects of Mindfulness-based cognitive Therapy programs on postpartum depression. The horizontal lines denote the 95% confidence interval, ■ point estimate (size of the square corresponds to its weight); ♦, combined Overall effect of treatment



**Fig.4:** Effects of problem-solving skills programs on postpartum depression. The horizontal lines denote the 95% confidence interval, ■ point estimate (size of the square corresponds to its weight); ♦, combined Overall effect of treatment



CI: Confidence interval; SMD: Standardized mean difference.

**Fig.5:** Funnel plot of results from included published studies on the effects of education program on postpartum depression.

#### **4- DISCUSSION**

To best our knowledge, this is the first meta-analysis assessed the effect of psychotherapy treatment for postpartum depression. Five studies were included into meta-analysis. Two studies used Problem-solving skills (1, 28); one study utilized Group psychotherapy (27) and two studies used mindfulness-based cognitive therapy (17, 29). In Kordi et al., depressive symptoms decreased significantly in problem-solving skills group compared to control group (28). In Naddaf et al., post partum depression improved significantly in group psychotherapy compared to control group (27). In Ghadam pour et al.'s study, the Beck Depression Inventory-II (BDI-II) decreased significantly in Mindfulness-based cognitive therapy group compared to control group (17).

In Bakhtiari et al.'s study BDI decreased significantly in mindfulness-based cognitive group compared to control group (29). In Parsa et al.'s study, based on Edinburgh scale, postpartum depression improved significantly in Problem-solving skills group compared to control group. According to forest plot, three studies showed educational program were effective in decreasing severity of depression (1, 28, 29). In contrast in two studies, severity of postpartum depression decreased non-significantly. Discrepancy between studies may be related to difference in content from training programmers, sample size, severity of depression, different instrument tool and setting. After of combination of the result of five studies in a meta-analysis, educational intervention was found to be more effective and could significantly improve postpartum depression.

##### **4-1. Limitation of study**

This systematic review had several limitations that need to be noted and discussed. The first limitations are large heterogeneity among studies that may be

due to different instrument tool, content from training programmers, difference in severity of depression, difference in sample size and setting. The second, some of studies did not provide the information of education session. Third limitation, Higgins (30) suggested that a minimum of ten studies need to conduct a meta-regression. Therefore, the number of studies included in the systematic review was too small to perform a Meta regression to detect whether severity of depression impact on the result of meta-analysis. However, sensitivity analysis showed no impact of severity of depression on result. Fourth limitation, most of studies included into meta-analysis had a small sample size that larger sample size must be considered in future study. Fifth, the fact that all the studies performed in Iran may limit the generalizability of the finding of the meta-analysis to other countries. Last, despite of that fact that we performed a comprehensive search, but it is possible that some of studies be missed length of the systematic review is all studies used validated and reliable instrument tool. Also, studies were well-designed and described base on consort guideline.

#### **5- CONCLUSIONS**

Based on the finding of the current meta-analysis, it can be said that educational programs that teaches Mindfulness-based cognitive and Problem-solving skills, can improve postpartum depression. Psychotherapy and educational intervention can offer as an alternative to pharmacotherapy particularly for women who are concern about side effect of herbal and chemical drugs.

#### **6- CONFLICT OF INTEREST**

All the authors declare that they have no conflict of interest.

#### **7- REFERENCES**

1. Parsa P, Ahangpour P, Shobeiri F, Soltanian A, Rahimi A. The Effect of Group Counseling Based on Problem Solving on Postpartum Depression in Mothers Attending to Health Care Centers in Hamadan City. *Journal of Urmia Nursing and Midwifery Faculty*. 2017;15(6):440-8.
2. Chaaya M, Campbell O, El Kak F, Shaar D, Harb H, Kaddour A. Postpartum depression: prevalence and determinants in Lebanon. *Archives of women's mental health*. 2002;5(2):65-72.
3. O'Hara M, Gorman LL. Can postpartum depression be predicted ?Primary Psychiatry. 2004.
4. Moses-Kolko EL, Roth EK. Antepartum and postpartum depression: healthy mom, healthy baby. *Journal of the American Medical Women's Association (1972)*. 2004;59(3):181-91.
5. Gjerdingen DK, Yawn BP. Postpartum depression screening: importance, methods, barriers, and recommendations for practice. *The Journal of the American Board of Family Medicine*. 2007;20(3):280-8.
6. Fergerson SS, Jamieson DJ, Lindsay M. Diagnosing postpartum depression: can we do better? *American Journal of Obstetrics & Gynecology*. 2002;186(5):899-902.
7. Jafari-Shobeiri M, Ghojzadeh M, Azami-Aghdash S, Naghavi-Behzad M, Piri R, Pourali-Akbar Y, et al. Prevalence and risk factors of gestational diabetes in Iran: a systematic review and meta-analysis. *Iranian journal of public health*. 2015;44(8):1036.
8. Pour EG, Azizi A, Mohamadi J. The Efficacy of Detached Mindfulness in Meta-Cognitive Therapy on Postpartum Depression. *Journal of Nursing Education*. 2016;5(5):17-22.
9. Kwon SJ, Lee CW. Figurate Purpuric Eruptions on the Trunk: Acetaminophen-induced Rashes. *The Journal of dermatology*. 1998;25(11):756-8.
10. Corwin EJ, Murray-Kolb LE, Beard JL. Low hemoglobin level is a risk factor for postpartum depression. *The Journal of nutrition*. 2003;133(12):4139-42.
11. Stuart S, O'Hara M, Gorman L. The prevention and psychotherapeutic treatment of postpartum depression. *Archives of Women's Mental Health*. 2003;6(2):s57-s69.
12. Davis L, Edwards H, Mohay H, Wollin J. The impact of very premature birth on the psychological health of mothers. *Early human development*. 2003;73(1-2):61-70.
13. Kalantaridou SN, Makrigiannakis A, Mastorakos G, Chrousos GP. Roles of Reproductive Corticotropin-Releasing Hormone. *Annals of the New York Academy of Sciences*. 2003;997(1):129-35.
14. Abedini S, Golmakani N, Behdani F, Esmaeili H, Safa O. Comparing marital satisfaction between women with and without. *Bimonthly Journal of Hormozgan University of Medical Sciences*. 2006;10(3):251-6.
15. Sanford D. Postpartum depression: The most frequent complication of childbirth. *International Journal of Childbirth Education*. 2002;17(1):10.
16. Zare Z, Golmakani N, Shareh H, Khadem N. Factors related to marital satisfaction in primiparous women during postpartum period. *Journal of Midwifery and Reproductive Health*. 2014;2(2):120-7.
17. Ghadam Pour E, Azizi A, Mohamadi J. The Efficacy of Detached Mindfulness in Meta-Cognitive Therapy on Postpartum Depression. *Journal of Nursing Education*. 2016;5(5):17-22.
18. Milofsky J. Milofsky J. A Provider's handbook on culturally competent care. Page 70.
19. Weier KM, Beal MW. Complementary therapies as adjuncts in the treatment of postpartum depression. *Journal of Midwifery and Women's Health*. 2004;49(2):96-104.
20. Goodman JH, Santangelo G. Group treatment for postpartum depression: a systematic review. *Archives of women's mental health*. 2011;14(4):277-93.
21. Malta LA, McDonald SW, Hegadoren KM, Weller CA, Tough SC. Influence of interpersonal violence on maternal anxiety, depression, stress and parenting morale in the early postpartum: a community based



pregnancy cohort study. *BMC pregnancy and childbirth*. 2012;12(1):153.

22. Toopkanlooi S. The Effectiveness of Existential Group Psychotherapy on Postpartum Depression in Nulliparous Women. *Scientific Journal of Ilam University of Medical Sciences*. 2016;23(6):185-95.

23. Troductio I. Prevalence, quality of life and social support in postpartum depression: a longitudinal study. *Pain*.48:000. Conference: Conference: The Asian Conference on the Social Sciences. ACSS 2011, At Osaka (Japan).

24. Mancini F, Carlson C, Albers L. Use of the Postpartum Depression Screening Scale in a collaborative obstetric practice. *Journal of Midwifery and women's Health*. 2007;52(5):429-34.

25. de Mendonça JS, Bussab VS, Rodrigues A, Siqueira J, Cossette L. Postpartum depression, father's involvement, and marital and co-parental relationships from mothers' and fathers' perspectives in a low-income Brazilian sample. *Family Science*. 2012;3(3-4):164-73.

26. O'Mahony JM, Donnelly TT, Bouchal SR, Este D. Cultural background and socioeconomic influence of immigrant and

refugee women coping with postpartum depression. *Journal of immigrant and minority health*. 2013;15(2):300-14.

27. Naddap S, Toopkanlooi, Safoora. The Effectiveness of Existential Group Psychotherapy on Postpartum Depression in Nulliparous Women. *scientific journal of ilam university of medical sciences*. 2016;23(6):185-95.

28. Nasiri S, Kordi M, Ebrahimzadeh S, Modarres Gharavi M. The effect of problem solving skills training on severity of depression symptoms in postpartum period. *J Fundament Ment Health*. 2012;14(55):226-35.

29. Abad ANS, Bakhtiari M, Kashani FL, Habibi M. The comparison of effectiveness of treatment based on acceptance and commitment with cognitive-behavioral therapy in reduction of stress and anxiety in cancer patients. *International Journal of Cancer Research and Prevention*. 2016;9(3):229.

30. Higgins J. *Cochrane handbook for systematic reviews of interventions version 5.0. 2* (updated September 2009). The Cochrane Collaboration. Available at: <http://www.Cochrane-handbook.org>. 2009.

**Table-1:** Characteristics of 5 studies included in our systematic review.

Author, City, Year, Reference	Design	Age	Level of complaints and measurement tool	Measurement tool	Type of interventions	Participants Intervention	randomization /Suitable randomization technique	Basel Comparability of the treatment and control groups	Major relevant findings
Parsa et al. Hamadan of Iran, 2016 (1)	Randomised controlled trials (RCTS )	Age range 19-23 years	10-15 scores based on Edinburgh scale	Edinburgh Scale	Problem-solving skills /Eight sessions weekly	Intervention (n=41) and control (n=41)	Yes/no	Yes	Edinburgh Scale improved significantly in intervention group compared to control group.
Ghadam pour et al., Tabriz of Iran, 2016 (17)	Quasi-experimental	26 years	High depression according to BDI	Beck Depression Inventory-II	Mindfulness-based cognitive Therapy Eight sessions (two sessions a week)	Intervention (n=15), and control (n=15)	Convenience method/no	Yes	Beck Depression Inventory-II decreased significantly in intervention group compared to control group.
Naddap et al., Mashhad of Iran, 2012, (27)	Quasi-experimental	23.29 years	Edinburgh Scale $\geq 10$ Mild-moderate depression	Edinburgh Scale	Group psychotherapy ten session, control/ no intervention	Intervention (n=13), and control (n=13)	Yes/no	Yes	Edinburgh Scale improved significantly in intervention group compared to control group.
Kordi et al., Mashhad of Iran, 2012 (28)	Randomised controlled trials (RCTS )	Intervention =25.8 years; Control group=25.7 years	Edinburgh Scale $\geq 10$ Beck Depression I between 4 to 28	Edinburgh Scale	Problem-solving skills /six sessions of 45-50 min; control/ no intervention	Intervention (n=26), and control (n=28)	Yes/no	Yes	Depressive symptoms decreased significantly in problem-solving skills group compared to control group.
Bakhtiari et al., Esfahan, of Iran, 2012 (29)	Experimental	Intervention group =24 years, and control group=26 years	At least 5-7 scores based on Beck depression inventory (BDI)	Beck depression	Mindfulness-based cognitive Therapy Eight sessions weekly	Intervention (n=15) and control (n=15)	Yes/no	Yes	BDI decreased significantly in intervention group compared to control group.