

Tumor Necrosis Factor alpha, N-terminal Pro Brain Natriuretic Peptide and Interleukin-6 Correlations with Doppler Tissue Imaging findings in Major Thalassemia Patients

Noor Mohammad Noori¹, *Alireza Teimouri¹, Maryam Nakhaee Moghadam¹, Hossein Bagheri²

¹Children and Adolescents Health Research Center, Resistant Tuberculosis Institute, Zahedan University of Medical Sciences and Health Services, Zahedan, Iran.

²English Department, Medical School, Zahedan University of Medical Sciences and Health Services, Zahedan, Iran.

Abstract

Background

In respect to abnormalities in thalassemia patients and the effects of TNF α , IL $_6$, and NT-pro-BNP on echocardiography findings the study aimed to investigate TNF α , NT-pro-BNP and IL $_6$ correlations with Doppler tissue imaging findings in major thalassemia.

Materials and Methods: This case-control study was performed on 112 healthy children and 112 patients with thalassemia in Ali Asghar hospital, Zahedan, Iran, in 2017. Healthy children were selected from those that referred to the pediatric clinics for routine check-up. Fasting blood samples were taken to measure TNF α , NT-pro-BNP and IL $_6$ serums. Those thalassemia patients who had regular transfusions of hemoglobin higher than 10 g/dl and 48-72 h after packed red blood cell transfusion and those who were asymptomatic were entered to the study. Participants underwent echocardiography by pediatric cardiologist in Doppler method.

Results: NT-pro-BNP, TNF α and IL $_6$ were lower in controls ($P<0.001$). Height, weight and BMI were higher in controls ($P<0.001$). Right ET, left ET, IVSD, PWD, PWS, EF, FS, left ET', left ICT', left IRT', right ET', right IRT', right S', right Peak A', VE', VA', left and right MPI, LVMI were different in case and control ($P<0.05$). Right Peak A ($P=0.049$), PWD ($P=0.032$), Left MPI ($P=0.02$) and LVMI ($P=0.005$) were correlated with IL $_6$. IVSD ($P=0.01$), LVDS ($P=0.01$), EF and FS ($P<0.001$), left ET' ($P=0.02$), left ICT' ($P<0.001$), right ET' ($P=0.001$), right S' ($P=0.002$), left MPI' ($P<0.010$) were correlated with NT-pro-BNP.

Conclusion

NT-pro-BNP, TNF α , IL $_6$ were higher in thalassemia patients and NT-pro-BNP was correlated with some of the echocardiography findings. This suggests that NT-pro-BNP would be the best biomarker for evaluation of cardiac functions in thalassemia patients.

Key Words: Children, Tumor necrosis factor- α , NT-pro-BNP, Interleukin-6, Thalassemia.

*Please cite this article as: Noori NM, Teimouri A, Nakhaee Moghadam M, Bagheri H. Tumor Necrosis Factor alpha, N-terminal Pro Brain Natriuretic Peptide and Interleukin-6 Correlations with Doppler Tissue Imaging findings in Major Thalassemia Patients. *Int J Pediatr* 2019; 7(10): 10255-268. DOI: **10.22038/ijp.2019.40837.3445**

*Corresponding Author:

Alireza Teimouri: M.Phil, Ph.D in Demography, Children and Adolescent Health Research Center, Resistant Tuberculosis Institute, Zahedan University of Medical Sciences, Zahedan, Iran.

Email: Alirezateimouri260@gmail.com

Received date: Mar.15, 2019 ; Accepted date: Aug.22, 2019

1- INTRODUCTION

Thalassemia is blood disorders are inherited from parents and are characterized by abnormal hemoglobin. Newborn babies with thalassemia account for 60,000 to 100,000 cases every year globally (1, 2). It occurs in Mediterranean Sea area and extends in a line to regions in the Far East and has now become a global health problem because of migration (3, 4). Iran is located in Mediterranean belt and has approximately 25,000 thalassemia patients and three million carriers. In Iran, the Persian Gulf and the Caspian Sea areas have a prevalence of more than 10% and Sistan and Baluchistan (SB) province has 4 to 10% (1). The province of SB has thalassemia gene frequency in high numbers of patients who are receiving regular health services such as blood transfusion (1, 3, 5).

In thalassemia, one of the most important organs that is impressed by iron overload is heart (6). Natriuretic Peptides (NPs) have been considered as diagnostic biomarkers for cardiovascular diseases, and they contain B-type natriuretic peptide (BNP), prohormone brain natriuretic peptide (pro BNP). In comparison, N Terminal Pro B Type Natriuretic Peptide (NT-pro BNP), is more stable with longer half-life, about 2 h compared to BNP (7), and it is useful for screening asymptomatic patients who are high risk such as Thalassemia patients (2).

In the early phases of the cardiac involvement, the level of NT-pro-BNP increases before an increase in diastolic pressure and there is a strong relationship between plasma levels of NT-pro-BNP and iron overload (8, 9). A significant relationship between NT-pro-BNP and some diastolic dysfunctions has been observed (9), and approved that an increase can be used as a tool for primary detection of cardiac hemosiderosis (1). Cytokines have great immunologic actions

and are considered as important catabolic factors and it seems that cytokines such as tumor necrosis factor alpha (TNF- α), and interleukin IL-6 are important mediators of the cachectic process in cardiac patients and are secreted secondary to several chronic diseases. In addition, they play an important role in cardiac cachexia and exacerbate prognosis (5). In sickle cell anemia, TNF- α has been linked as a risk factor for the occurrence of painful crises, as well as participating in the occlusion of the microcirculation (7). Moreover, pro-resorptive levels of TNF- α and IL-6 are elevated in patients with thalassemia.

Several studies have demonstrated that there is an association between NT-Pro-BNP (1), TNF α (5), IL-6 (5, 7), and echocardiography findings in thalassemia, congenital heart defect (CHD), and sickle cell anemia patients (2). Tissue Doppler Echocardiography (TDE) has become an established arm of diagnostic ultrasound examination and uses frequency shift of ultrasound waves to calculate myocardial velocity. Two techniques are used to assess myocardial function: pulsed TDE and color-coded TDE. It has the advantage of increased spatial resolution and ability to evaluate multiple structures in a single view (10). The usefulness of TDE in detection of myocardial iron overload has been reported (11).

Kanazirev et al., found that in asymptomatic individuals' diastolic dysfunctions by TDE are proper and significant tools in the prognostic development of symptomatic iron-induced cardiac disease (11). Regarding abnormalities in thalassemia patients such as cardiac involvement and the association with TNF- α , IL-6 and NT-pro-BNP the present study aimed to investigate the possible relationship between these cytokines and NT-pro-BNP with DTE findings in thalassemia patients compared to healthy children.

2- MATERIALS AND METHODS

2-1. Study design

This case-control study was performed on 224 participants aged 7 to 18 years consisted of 112 healthy children as controls and 112 patients with thalassemia as case group in pediatric cardiac center in collaboration with Center for Specific Diseases in Ali Asghar hospital, Zahedan city, Sistan and Baluchestan province, Iran, during the year 2017. Controls were chosen from those healthy children that referred to the clinics for routine check-up.

2-2. Criteria

Those thalassemia patients who had regular transfusions of hemoglobin higher than 10 g/dl and 48-72 h after packed red blood cell transfusion were entered to the study. Exclusion criteria in both groups of participants were valvular heart disease, rhythm and structural abnormality, active infection, systemic inflammatory diseases, renal insufficiency, and these criteria were considered for both groups.

2-3. Echocardiography measurements

Major proceedings on patients were checking medical history, physical examination, chest X-ray and echocardiography which was performed by one pediatric cardiologist. Echocardiography was performed on patients by the same pediatric cardiologist using My lab 60 with transducer 3, 8 [made in Italy). In order to achieve high precision in echocardiography findings, measurement were repeated for 3 cycles and the average was considered. Echocardiogram was performed on participants without them holding their breath, and view was taken in the mitral valve surface in parasternal position (1). Echocardiography findings were as follows: LVDD: left ventricular end-diastolic dimension, Peak E: early mitral and tricuspid valve flow velocity, Peak A: late mitral and tricuspid valve flow

velocity, ET: ejection time (for Aorta and Pulmonary), PWD: posterior wall dimension in diastole, IVSD: interventricular septal dimension in Diastole, IVSS: interventricular septal dimension in systole, EF: ejection fraction, FS: fractional shortening, LVM: left ventricular mass, LVMI: left ventricular mass index, were measured using conventional echocardiography of the left side and estimated from three cardiac cycles. Myocardial performance index (MPI), isovolumic relaxation time (IRT), isovolumic contraction time (ICT) of both sides, were measured with echocardiography (12). The sample volume was positioned at the tips of the tricuspid and mitral valve leaflets in the apical four-chamber view to enable the measurement of (a): the time of interval between the end and the start of trans-mitral and trans-tricuspid flow.

The sample volume was thereafter relocated to the left ventricular outflow tract just below the aortic valve (apical five-chamber view), so as to measure (b): which is the left ventricular ejection time. The right ventricular outflow velocity pattern was also recorded from the parasternal short-axis view with the Doppler sample volume positioned just distal to the pulmonary valve for the measurement of (b). Myocardial Performance Index (Tei Index), was calculated as: $a-b/b = IRT + ICT/ET$. The left ventricular mass index (LVMI), was also calculated by the following formula: $LVM (g) = 0.8 (1.04 (LVEDD + PWTD + IVSTD) - LVEDD) + 0.6$, and $LVMI (g/m^2) = LVM / 2.7 (g/m^2)$. All the parameters in the above formula were measured in the M-mode view and in diastole and were utilized for left ventricular mass evaluation (1). Doppler tissue echocardiography (DTI) was another method that was performed from the apical four-chamber view and a 3 mm pulsed Doppler sample volume was placed at the

level of lateral mitral annulus. Myocardial velocity profiles of the lateral tricuspid annulus and lateral mitral annulus were obtained by placing the sample volume at the junction of the tricuspid annulus and the right ventricle (RV) free wall and at the junction of the mitral annulus and LV posterior wall, respectively. With this modality, the recorded values were the

early (E) and late (A) diastolic mitral and tricuspid annular velocities, and the ratio of E/A. Right ventricle and left ventricle myocardial performance index (MPI) was obtained by dividing the sum of isovolumic relaxation time (IRT) and isovolumetric contraction time (ICT) by the ejection time (ET) ($MPI = (ICT + IRT)/ET$) (Figure.1).




Fig.1: Diagram of Doppler Tissue Echocardiography waves: S', systolic wave; E', early diastolic wave; A', late diastolic wave (12).

Left and right S': Systolic myocardial velocity above the baseline in pulmonic vein, mitral and tricuspid Left and Right E': early diastolic myocardial relaxation velocity below the baseline in pulmonic vein, mitral and tricuspid. Left and Right A': myocardial velocity associated with atrial contraction in pulmonic vein, mitral and tricuspid.

2-3. NT-pro-BNP, TNF- α and Interlukin_6 measurements

To measure NT-pro-BNP, TNF- α and IL-6, 3 milliliters (ml) of blood was taken from all participants at 8.00 AM after an overnight fasting. The samples were centrifuged at 3,000 rpm/min for 10 min at 5 °C. Separated serum of NT-pro-BNP, TNF- α and IL-6 were kept in -70cc refrigerator until measuring. The samples were transferred to the biochemistry library of Zahedan University of Medical Sciences with considering the cold chain circumstances and then 250 microns isolated serum samples were selected for measuring the levels of NT-pro-BNP and as well as TNF- α and Interlukin_6 by ELISA kit (USA).

2-4. Anthropomorphic measurements

Children's height and weight were measured. The recumbent length for children under 2 years was graded using a flat wooden table and weight measurements for children under 2 years were performed by using Mika balance weights with difference of 100gr. Then BMI was calculated with (Kg / m^2) (weight in kilograms divided by height to the power of 2) formula.

2-5. Ethical Approval

Consent form was obtained from the participants or their guardians after the study approval. The study was approved as a project proposed to the Children and Adolescent Health Research Center by the Ethics Committee of Zahedan University of Medical Sciences, Zahedan, Iran (ID number: 6798).

2-6. Statistical Analysis

Statistical software of SPSS version 20.0 for windows was used for all statistical analyses. Data were reported as mean \pm standard deviation (SD) for continuous

variables or percentage for categorical variables. Differences in baseline characteristics between the two groups were evaluated with Mann-Whitney U test for all continuous variables, all categorical variables were compared using chi-square test. All p-values were two-sided and the results were considered statistically significant at the level of $p < 0.05$.

3- RESULTS

The study aimed to assess NT-pro-BNP, TNF $_{\alpha}$ and IL $_6$ correlations with echocardiography findings in thalassemia patients. In the results, due to normality analysis of Kolmogorov-Smirnov test, the null hypothesis stated that the data followed free distribution because of the p-value that was less than 0.05 in the majority of variables except a few (left Peak E and left Peak E' with p-value lower than [0.05]), especially considering all participants. Participants' sex distribution was similar between case and controls

(Chi-square=0.029 and $P=0.866$) (Table.1). Table.2 showed that controls and patients were matched by age ($p > 0.05$). The table also showed that NT-pro-BNP (119.55±199.84 vs. 606.21±993.34), TNF $_{\alpha}$ (23.18±13.43 vs 39.72±18.54) and IL $_6$ (3.01±1.00 vs 23.49±13.84) were lower in controls significantly compared to patients ($P < 0.001$). Height, weight and BMI were significantly higher in controls compared to patients ($P < 0.001$). Amongst echocardiography findings, Right ET ($P < 0.001$), Left ET ($P < 0.001$), IVSD ($P < 0.001$), PWD ($P=0.029$), PWS (0.014), EF ($P < 0.001$), FS ($P < 0.001$), left ET' ($P < 0.001$), left ICT' ($P=0.041$), Left IRT' ($P < 0.001$), right ET' ($P=0.004$), right IRT' ($P=0.004$), right S' ($P < 0.001$), right Peak A' ($P=0.049$), VE' ($P=0.010$), VA' ($P=0.011$), left and right MPI' ($P < 0.001$), LVMI ($P=0.004$) were different between case and control significantly.

Table-1: Sex distribution of participants in patients and controls.

| Gender | Statistics | Groups of participants | | Chi-square | P-value |
|--------|------------|------------------------|------------|------------|---------|
| | | Control | Case | | |
| Female | Number (%) | 37(50.74%) | 36(49.26%) | 0.029 | 0.866 |
| Male | Number (%) | 33(49.30%) | 34(50.70%) | | |

Table-2: Case-control comparison of the interested variables in the study.

| Variables | Participants | Mean | SD | Mean Rank | Sum of Ranks | M-W U | P- value |
|------------|--------------|--------|--------|-----------|--------------|--------|----------|
| Age | Control | 11.49 | 4.04 | 70.4 | 4928 | 2443 | 0.977 |
| | Case | 11.27 | 3.92 | 70.6 | 4942 | | |
| NT-pro-BNP | Control | 119.55 | 199.84 | 50.54 | 3538 | 1053 | <0.001 |
| | Case | 606.21 | 993.34 | 90.46 | 6332 | | |
| TNF | Control | 23.18 | 13.43 | 52.06 | 3644.5 | 1159.5 | <0.001 |
| | Case | 39.72 | 18.54 | 88.94 | 6225.5 | | |
| Interlukin | Control | 3.01 | 1 | 38.42 | 2689.5 | 204.5 | <0.001 |
| | Case | 23.49 | 13.84 | 102.58 | 7180.5 | | |
| Height | Control | 155.04 | 14.32 | 91.41 | 6399 | 986 | <0.001 |
| | Case | 134.07 | 18.83 | 49.59 | 3471 | | |
| Weight | Control | 44.7 | 12.92 | 90.56 | 6339 | 1046 | <0.001 |
| | Case | 30.27 | 11.25 | 50.44 | 3531 | | |

| | | | | | | | |
|--------------|---------|--------|-------|-------|--------|--------|--------|
| BMI | Control | 18.23 | 3 | 84.3 | 5901 | 1484 | <0.001 |
| | Case | 16.14 | 2.49 | 56.7 | 3969 | | |
| Left Peak E | Control | 102.09 | 17.91 | 71.51 | 5006 | 2379 | 0.767 |
| | Case | 99.7 | 19.98 | 69.49 | 4864 | | |
| Left Peak A | Control | 52.42 | 8.84 | 68.09 | 4766 | 2281 | 0.481 |
| | Case | 54.86 | 12.43 | 72.91 | 5104 | | |
| Right Peak E | Control | 70.13 | 11.28 | 72.54 | 5077.5 | 2307.5 | 0.552 |
| | Case | 70.22 | 13.85 | 68.46 | 4792.5 | | |
| Right Peak A | Control | 49.18 | 12.73 | 64.5 | 4515 | 2030 | 0.080 |
| | Case | 51.37 | 11.14 | 76.5 | 5355 | | |
| Right ET | Control | 288.15 | 28.97 | 90.68 | 6347.5 | 1037.5 | <0.001 |
| | Case | 260.39 | 21.16 | 50.32 | 3522.5 | | |
| Left ET | Control | 287.76 | 21.99 | 97.94 | 6855.5 | 529.5 | <0.001 |
| | Case | 248.14 | 26.76 | 43.06 | 3014.5 | | |
| IVSD | Control | 0.66 | 0.11 | 56.32 | 3942.5 | 1457.5 | <0.001 |
| | Case | 0.75 | 0.13 | 84.68 | 5927.5 | | |
| LVDD | Control | 3.9 | 0.4 | 68.09 | 4766.5 | 2281.5 | 0.482 |
| | Case | 3.92 | 0.73 | 72.91 | 5103.5 | | |
| PWD | Control | 0.36 | 0.05 | 63.06 | 4414 | 1929 | 0.029 |
| | Case | 0.38 | 0.09 | 77.94 | 5456 | | |
| IVSS | Control | 0.83 | 0.15 | 64.78 | 4534.5 | 2049.5 | 0.094 |
| | Case | 0.88 | 0.14 | 76.22 | 5335.5 | | |
| LVDS | Control | 2.1 | 0.29 | 66.57 | 4660 | 2175 | 0.252 |
| | Case | 2.21 | 0.52 | 74.43 | 5210 | | |
| PWS | Control | 0.36 | 0.05 | 62.11 | 4347.5 | 1862.5 | 0.014 |
| | Case | 0.39 | 0.08 | 78.89 | 5522.5 | | |
| EF | Control | 73.88 | 18.69 | 94.66 | 6626.5 | 758.5 | <0.001 |
| | Case | 42.07 | 34.51 | 46.34 | 3243.5 | | |
| FS | Control | 42.87 | 13.83 | 95.01 | 6651 | 734 | <0.001 |
| | Case | 22.7 | 18.93 | 45.99 | 3219 | | |
| LVM | Control | 51.19 | 15.59 | 66.59 | 4661.5 | 2176.5 | 0.254 |
| | Case | 55.4 | 22.29 | 74.41 | 5208.5 | | |
| Left ET' | Control | 361.84 | 86.78 | 91.41 | 6399 | 986 | <0.001 |
| | Case | 263.03 | 52.87 | 49.59 | 3471 | | |
| Left ICT' | Control | 94.33 | 15.41 | 77.49 | 5424 | 1961 | 0.041 |
| | Case | 89.69 | 28.98 | 63.51 | 4446 | | |
| Left IRT' | Control | 70.51 | 12.29 | 47.79 | 3345 | 860 | <0.001 |
| | Case | 96.03 | 38.38 | 93.21 | 6525 | | |
| Left S' | Control | 8.95 | 2.02 | 71.76 | 5023 | 2362 | 0.714 |
| | Case | 8.69 | 1.57 | 69.24 | 4847 | | |
| Left Peak E' | Control | 14.65 | 2.52 | 71.54 | 5007.5 | 2377.5 | 0.762 |
| | Case | 14.57 | 3.26 | 69.46 | 4862.5 | | |

| | | | | | | | |
|---------------|---------|--------|-------|-------|--------|--------|--------|
| Left Peak A' | Control | 6.42 | 1.85 | 67.52 | 4726.5 | 2241.5 | 0.384 |
| | Case | 6.67 | 1.49 | 73.48 | 5143.5 | | |
| Right ET' | Control | 280.03 | 58.98 | 80.4 | 5628 | 1757 | 0.004 |
| | Case | 255.89 | 38.81 | 60.6 | 4242 | | |
| Right ICT' | Control | 75.84 | 13.93 | 69.6 | 4872 | 2387 | 0.791 |
| | Case | 77.98 | 18.57 | 71.4 | 4998 | | |
| Right IRT' | Control | 96.14 | 21.69 | 60.84 | 4258.5 | 1773.5 | 0.004 |
| | Case | 107.61 | 22.57 | 80.16 | 5611.5 | | |
| Right S' | Control | 64.51 | 20.47 | 81.87 | 5731 | 1654 | 0.001 |
| | Case | 42.81 | 41.33 | 59.13 | 4139 | | |
| Right Peak E' | Control | 14.58 | 2.51 | 67.99 | 4759 | 2274 | 0.463 |
| | Case | 14.63 | 2.98 | 73.01 | 5111 | | |
| Right Peak A' | Control | 6.69 | 2.14 | 63.75 | 4462.5 | 1977.5 | 0.049 |
| | Case | 7.75 | 2.99 | 77.25 | 5407.5 | | |
| VS' | Control | 10.18 | 2.12 | 76.24 | 5337 | 2048 | 0.094 |
| | Case | 9.62 | 1.63 | 64.76 | 4533 | | |
| VE' | Control | 17.03 | 2.19 | 79.31 | 5551.5 | 1833.5 | 0.010 |
| | Case | 15.92 | 2.71 | 61.69 | 4318.5 | | |
| VA' | Control | 8.87 | 2.84 | 79.21 | 5545 | 1840 | 0.011 |
| | Case | 7.78 | 1.63 | 61.79 | 4325 | | |
| Left MPI' | Control | 0.48 | 0.11 | 45.07 | 3155 | 670 | <0.001 |
| | Case | 0.74 | 0.25 | 95.93 | 6715 | | |
| Right MPI' | Control | 0.63 | 0.12 | 57.45 | 4021.5 | 1536.5 | <0.001 |
| | Case | 0.74 | 0.18 | 83.55 | 5848.5 | | |
| LVMI | Control | 23.64 | 7.42 | 60.71 | 4250 | 1765 | 0.004 |
| | Case | 28.85 | 11.85 | 80.29 | 5620 | | |
| Left A/A' | Control | 9.79 | 7.64 | 70.41 | 4928.5 | 2443.5 | 0.978 |
| | Case | 8.84 | 4.68 | 70.59 | 4941.5 | | |
| Left E/E' | Control | 7.32 | 2.9 | 69.25 | 4847.5 | 2362.5 | 0.715 |
| | Case | 7.25 | 2.62 | 71.75 | 5022.5 | | |
| Right E/E' | Control | 5.03 | 1.72 | 72.56 | 5079.5 | 2305.5 | 0.547 |
| | Case | 5.12 | 2.07 | 68.44 | 4790.5 | | |
| Right A/A' | Control | 8.94 | 8.68 | 74.32 | 5202.5 | 2182.5 | 0.265 |
| | Case | 7.91 | 6.45 | 66.68 | 4667.5 | | |

M-W U: Mann-Whitney U, SD: Standard Deviation, Peak E: Early mitral and tricuspid valve flow velocity, Peak A: Late mitral and tricuspid valve flow velocity, ET: Ejection time (for Aorta and Pulmonary), IVSD: Interventricular septal dimension in diastole, LVDD: Left ventricular end-diastolic dimension, PWD: Posterior wall dimension in diastole, IVSS: Interventricular septal dimension in systole, LVDS: Left ventricular diameter at systole, PWS: Posterior wall dimension in systole, EF: Ejection fraction, FS: Fractional shortening, LVM: Left ventricular mass, ICT: Isovolumetric contraction time, ICT: Isovolumetric contraction time, S': Systolic myocardial velocity above the baseline in pulmonic vein, mitral and tricuspid, E': early diastolic myocardial relaxation velocity below the baseline in pulmonic vein, mitral and tricuspid. A': myocardial velocity associated with atrial contraction in pulmonic vein, mitral and tricuspid, MPI: Myocardial Performance Index (Tei Index), BMI: Body Mass Index, LVMI: Left Ventricular Mass Index.

Table.3 showed the correlation between Doppler finding and major variables of the study. Left Peak E was correlated with BMI ($P<0.001$). Right Peak A had correlation with IL_6 ($P=0.049$). Left ET correlated with height ($P=0.001$) and weight ($P=0.008$). IVSD correlated with NT-pro-BNP ($P=0.01$), height ($P<0.001$), weight ($P<0.001$), and age ($P<0.001$). LVDD correlated with height ($P<0.001$), weight ($P<0.001$), age ($P<0.001$) and BMI ($P=0.024$). PWD correlated with IL_6 ($P=0.032$), height ($P=0.032$), age ($P=0.001$). LVDS correlated with NT-Pro-BNP ($P=0.01$), height ($P=0.005$), weight ($P=0.009$), age ($P=0.001$), and BMI ($P=0.038$). PWS was correlated with height ($P=0.045$), age ($P=0.002$). EF and FS were correlated with NT-pro- BNP ($P<0.001$), height ($P<0.001$), weight ($P<0.001$), age ($P<0.001$), and BMI ($P<0.001$). Left ET' was correlated with NT-pro-BNP ($P=0.02$), weight ($P=0.024$), age ($P=0.001$), and BMI ($P=0.016$). Left ICT' was correlated with NT-pro- BNP ($P<0.001$), height ($P<0.001$), weight ($P<0.001$), and age ($P<0.001$).

Right ET' was correlated NT-pro- BNP ($P=0.001$), height ($P=0.003$), weight ($P<0.001$), age ($P=0.001$) and BMI ($P=0.004$). Right ICT' was correlated with age ($P=0.036$). Right IRT' correlated with age ($P<0.001$). Right S' was correlated with NT-pro- BNP ($P=0.002$), height ($P=0.001$), weight ($P=0.001$) and age ($P=0.001$). VE' was correlated with height ($P=0.04$). VA' was correlated with IL_6 ($P=0.002$). Left MPI' was correlated with NT-pro- BNP ($P<0.010$), height ($P=0.003$), weight ($P=0.02$), and age ($P<0.001$). Left MPI' correlated with IL_6 ($P=0.02$), height ($P<0.001$), weight ($P=0.003$) and age ($P<0.001$). LVMI correlated with IL_6 ($P=0.005$), height ($P<0.001$), weight ($P<0.001$), age ($P<0.001$) and BMI ($P=0.025$). Left E/E' was correlated with BMI ($P=0.009$).

(Please see the table.3 in the end of paper).

4- DISCUSSION

The results of the present study revealed that NT-pro-BNP, TNF- α , IL_6, were higher in case. As an adaptive mechanism of chronic anemia, heart rate is expected to be higher in thalassemia patients. Many studies indicated that NT-pro-BNP is a sensitive biomarker to detect asymptomatic LV dysfunction especially with an important role in diagnostic and prognostic thalassemia implications. These studies proposed an increase of NT-pro-BNP in thalassemia patients compared to controls that is in line with the present study results (1, 3, 13). In this line Noori et al., conducted a study on children and adolescents to find the correlation between NT-pro-BNP and cardiac involvement in patients with thalassemia and resulted that the NT-pro-BNP levels were significantly higher in thalassemia children compared to controls (1). Noori et al. (14), Kremastinos et al. (3), and Ozyoruk et al. (15), reported higher levels of NT-pro-BNP in the patients than controls, all these results were in line with the present study. Aessopos et al., found high levels of NT-Pro BNP in patients with overt heart failure, independent of severity of cardiac insufficiency (6, 16). In this line Kanazirev et al. (11) arrived at the same conclusion of higher level of NT-pro-BNP in thalassemia patients compared to controls. It was hypothesized that a relationship between serum chemokines and inflammatory responses in thalassemia patients may be established because of the presence of active inflammation in these patients. A broad range of cytokines such as, granulocyte-macrophage-colony-stimulating factor (GM-CSF), IL_1, IL_3, IL_6, IL_8 and TNF_ α are released by activated endothelium and these have been detected in the plasma of patients with sickle cell disease (SCD) (17). Abnormal IL_6 productions may play a significant

role in a number of diseases, such as autoimmune diseases, chronic inflammation, and lymphoid malignancies (18). The present study showed a low level of IL_6 for the asymptomatic thalassemia patients compared to controls that was in line with Asem Metwally Abo-Shanab et al. (19), Elzubeir et al. (18), Akcalı et al. (20), and El-Rasheidy et al. (21). It was suggested that an increase in TNF_α could be a cause for macrophage activation due to iron overload and the antigenic stimulation induced by chronic transfusion therapy. The activated macrophages were selectively phagocytosing apoptotic erythroid precursors, thereby contributing to ineffective erythropoiesis (22). Garadah et al. (23), and Ragab et al. (24) reported that TNF_α had higher level in patients compared to controls. All these results were similar to the present study.

Noori et al. (1, 25, 26) conducted some studies with the aims of echocardiography findings comparison between thalassemia patients and healthy group. In all studies, Noori et al., concluded that left heart findings of EF, FS and MPI were varied in groups that was consistent with the present study. EF and FS were higher and MPI was lower in controls compared to patients in all Noori et al.'s studies similar to the present study. The result of the LVMI was lower in control (25), consistent with the present study. LVDD was lower (1), and higher (25) in control while in the present study, it was higher in control. Right ejection time was lower in case resulted by Noori et al. (1, 26) in agreement with the present study but in another study they found similarity in case and control (25). Noori et al. presented that MPI and Peak E/A were different between case and control in the right heart that all were similar with the present study (1, 25, 26). Recently Noori et al. (14), performed a study on The Correlation of Ferritin and Leptin Serum Levels with Cardiac Involvement in Thalassemia Patients

Compared to Controls and concluded that PWD, PWS, EF, FS and right ET were different in patients and controls which were in line with the present study results. The ratio of E/E' as an important marker of diastolic dysfunction was either normal or increased, especially in cases of restrictive pattern of diastolic dysfunction and correlated with NT-pro-BNP (4). Kremastinos et al. (3, 13) found that NT-pro-BNP was correlated with E/E' in thalassemia patients and concluded that NT-pro-BNP may be an early marker of diastolic dysfunction. Similarly, Garadah et al. (23) reported higher NT-pro-BNP, greater E wave velocity with shorter deceleration time and a greater E/A ratio with a tendency to restrictive diastolic dysfunction in thalassemia patients. They establish a higher E/E' ratio and lower S' and E' values compared to controls. Sarray et al. (27) concluded that IL_6 was correlated with the duration of vaso-occlusive crisis, but TNF_α was not correlated. Compared to the present study, it is observed that IL_6 had significant correlation with LVSD and TNF_α had significant correlation with left MPI in thalassemia patients. Akpınar et al. (28) stated that NT-pro-BNP was correlated with systolic longitudinal myocardial velocities dissimilar with the present study.

The reason for the difference may be due to the type of thalassemia patients that asymptomatic patients considered in the present study. Noori et al., conducted a study to evaluate the correlation of NT-pro BNP and heart findings in thalassemia patients and found that with using a reference cut point for NT pro BNP, LAd, LVDS and RWT correlated with NT pro BNP levels patients (1). From the study it can be concluded that there is a correlation between Doppler finding and major variables of the study. Right Peak A had correlation with IL_6, IVSD correlated with NT-pro-BNP, PWD correlated with IL_6, LVDS correlated with NT-Pro-BNP,

EF and FS were correlated with NT-pro-BNP, Left ET' was correlated with NT-pro-BNP, Left ICT' was correlated with NT-pro-BNP, Right ET' was correlated with NT-pro-BNP, Right S' was correlated with NT-pro-BNP, VA' was correlated with IL_6. Left MPI' was correlated with NT-pro-BNP, Left MPI' correlated with IL_6, LVMI correlated with IL_6. Surprisingly, from the results concerned with correlation analysis it is resulted that TNF α was not correlated with any of echocardiography findings when NT-pro-BNP was correlated with some of the echocardiography findings compared to IL_6.

4-1. Study Limitations

The study limitations were lack of proper cooperation by participants especially controls and selecting asymptomatic patients amongst thalassemia.

5- CONCLUSIONS

NT-pro BNP, TNF- α , IL_6 were higher in thalassemia patients and they correlated with some of echocardiography findings. From the study it is also concluded that TNF α was not correlated with any of echocardiography findings when NT-pro-BNP was correlated with some of the echocardiography findings compared to IL_6. The findings of the study suggested that NT-pro BNP would be the best biomarker for evaluation of cardiac functions in thalassemia patients.

6- CONFLICT OF INTEREST: None.

7- ACKNOWLEDGMENTS

Authors would like to offer their warm thanks to all participants and their parents.

8- REFERENCES

1. Noori NM, Teimouri A, Anvari N. Diagnostic Value of N Terminal Pro B Type Natriuretic Peptide (NT-pro BNP) in Cardiac Involvement in Patients with Beta-

Thalassemia. *International Journal of Pediatrics*. 2017; 5(4):4641-62.

2. Noori NM, Mahjoubifard M, Mohammadi M, Fard AJ, Abassi A, Farzanegan B. Comparison of QT dispersion with left ventricular mass index in early diagnosis of cardiac dysfunction in patients with β -thalassemia major. *Iranian Red Crescent Medical Journal*. 2014; 16(5): e11698.

3. Kremastinos DT, Tsiapras DP, Kostopoulou AG, Hamodraka ES, Chaidaroglou AS, Kapsali ED. NT-proBNP levels and diastolic dysfunction in β -Thalassaemia major patients. *European journal of heart failure*. 2007; 9(5):531-6.

4. Noori NM, Teimouri A, Nakhaey Moghaddam M. Diagnostic Value of NT-pro BNP Biomarker and Echocardiography in Cardiac Involvements in Beta-thalassemia Patients. *International Journal of Pediatrics*. 2017; 5(11):6077-94.

5. Noori NM, Shahramian I, Teimouri A, Keyvani B, Mahjoubifard M. Serum Levels of Tumor Necrosis Factor- α and Interleukins in Children with Congenital Heart Disease. *The Journal of Tehran University Heart Center*. 2017; 12(1):15-22.

6. Aessopos A, Berdoukas V, Tsironi M: The heart in transfusion dependent homozygous thalassaemia today prediction, prevention and management. *Eur J Haematol* 2008; 80:93-106.

7. Laurentino MR, Maia Filho PA, Barbosa MC, Bandeira IC, da Silva Rocha LB, Gonçalves RP. Influence of β S-globin haplotypes hydroxyurea on tumor necrosis factor-alpha levels in sickle cell anemia. *Revista brasileira de hematologia e hemoterapia*. 2014; 36(2):121-5.

8. Shahramian I, Noori NM, Teimouri A, Akhlaghi E, Sharafi E. The correlation between serum level of leptin and troponin in children with major beta-Thalassemia. *Iranian journal of pediatric hematology and oncology*. 2015; 5(1):11.

9. Noori NM, Birjandi H, Nakhaey Moghaddam M, Teimouri A. Evaluation of Serum NT-Pro Brain Natriuretic Peptide Levels in Neonates with Respiratory Distress.

International Journal of Pediatrics. 2018;7769-80.

10.Kattamis A, Apostolakou F, Delaporta P, Papassotiriou I. Correlation of NT-ProBNP levels and cardiac iron concentration in patients with transfusion-dependent Thalassemia Major. *Blood*. 2011; 118:3192.

11.Kanazirev B, Dimova M, Kaleva V, Gercheva S, Petrova K, Zlateva V, Bocheva Y, Nikolova S, Hadjolyan K. Early identification of heart failure in patients with thalassemia major by NT-pro-BNP examination. Correlation with echocardiographic parameters of morphology and function. *International Journal of Medical and Health Research*. 2017; 3(12):22-9.

12.Noori N, Shahraki T, Teimouri A, Shahramian I. Cardiac Involvements in Patients with Celiac Disease by Doppler Tissue Echocardiography Compared to Conventional Echocardiography. *International Cardiovascular Research Journal*. 2018; 12(1):13-21.

13.Kremastinos DT, Hamodraka E, Parissis J, Tsiapras D, Dima K, Maisel A. Predictive value of B-type natriuretic peptides in detecting latent left ventricular diastolic dysfunction in β -thalassemia major. *American heart journal*. 2010; 159(1):68-74.

14.Noori NM, Birjandi H, Teimouri A. The Correlation of Ferritin and Leptin Serum Levels with Cardiac Involvement in Thalassemia Patients Compared to Controls. *International Journal of Pediatrics*. 2018;7623-38.

15.Ozyoruk D, Öner T, Oymak Y, Çelik HT. Comparison of Doppler echocardiographic and tissue Doppler velocity data in beta-thalassaemia major with high and normal NT-proBNP levels of children in the south-east region of Turkey. *Translational pediatrics*. 2014; 3(4):287.

16.Aessopos A, Farmakis D, Polonifi A, et al: Plasma B-type natriuretic peptide concentration in beta-thalassaemiapatient. *Eur J Heart Fail* 2007; 9: 537-54.

17.Zachariah M, Al Zadjali S, Bashir W, Al Ambusaidi R, Misquith R, Wali Y, Pathare A. Impact of mannose-binding protein gene

polymorphisms in Omani sickle cell disease patients. *Mediterranean journal of hematology and infectious diseases*. 2016; 8(1): e2016013.

18.Elzubeir AM, Alobied A, Halim H, Awad M, Elfaki S, Mohammed M, Saddig OA. Estimation of Serum Interleukin-6 level as a Useful Marker for Clinical Severity of Sickle Cell Disease among Sudanese Patients. *Int. J. of Multidisciplinary and Current research*. 2017; 5: 642-4.

19.Abo-Shanab AM, El-Desouky MA, Kholoussi NM, El-Kamah GY, Helwa IA, Fahmi AA. The relation between glutathione S-Transferase M1 null-genotype and cardiac problems in beta-thalassemia. *Acta Biochimica Polonica*. 2016; 63(2): 267-71.

20.Akcalı A, Kahraman Çeneli S, Gümüş P, Buduneli N, Lappin DF, Özçaka Ö. The association between thalassemia major and periodontal health. *Journal of periodontology*. 2015; 86(9):1047-57.

21.El-Rasheidy FH, Essa ES, Mahmoud AA, Nada AE. Elevated serum adiponectin is related to elevated serum ferritin and interleukin-6 in β -thalassaemia major children. *Journal of Pediatric Endocrinology and Metabolism*. 2016; 29(8):953-8.

22.Tanyong DI, Panichob P, Kheansaard W, Fucharoen S. Effect of tumor necrosis factor-alpha on erythropoietin-and erythropoietin receptor-induced erythroid progenitor cell proliferation in β -thalassemia/hemoglobin E patients. *Turkish Journal of Hematology*. 2015; 32(4):304.

23.Garadah TS, Jaradat AA, AlAlawi ME, Hassan AB, Sequeira RP. Pain frequency, severity and QT dispersion in adult patients with sickle cell anemia: correlation with inflammatory markers. *Journal of blood medicine*. 2016; 7: 255- 34.

24.Ragab SM, Safan MA, Obeid OM, Sherief AS. Lipoprotein-associated phospholipase A2 (Lp-PLA2) and tumor necrosis factor-alpha (TNF- α) and their relation to premature atherosclerosis in β -thalassemia children. *Hematology*. 2015; 20(4):228-38.

25.Noori NM, Mohamadi M, Keshavarz K, Alavi SM, Mahjoubifard M, Mirmesdagh Y. Comparison of Right and Left Side Heart

26. Functions in Patients with Thalassemia Major, Patients with Thalassemia Intermedia, and Control Group. *J The Univ Heart Ctr* 2013; 8(1):35-41.

27. Noori NM, Keshavarz K, Shahriar M. Cardiac and pulmonary dysfunction in asymptomatic beta-thalassaemia major. *Asian Cardiovascular and Thoracic Annals*. 2012 (5):555-9.

28. Sarray S, Saleh LR, Saldanha FL, Al-Habboubi HH, Mahdi N, Almawi WY. Serum

IL_6, IL-10, and TNF α levels in pediatric sickle cell disease patients during vasoocclusive crisis and steady state condition. *Cytokine*. 2015; 72(1):43-7.

29. Akpınar O, Acartürk E, Kanadaşı M, Unsal C, Başlamışlı F. Tissue doppler imaging and NT-proBNP levels show the early impairment of ventricular function in patients with beta-thalassaemia major. *Acta Cardiol*. 2007; 62(3):225-31.

Table-3: Correlation of NT-pro-BNP, TNF $_{\alpha}$, IL $_6$, and anthropometric indices with echocardiography findings in thalassemia patients.

| Variables | Statistics | NT-pro-BNP | TNF $_{\alpha}$ | Interleukin | Height | Weight | BMI | Variables | NT-pro-BNP | TNF $_{\alpha}$ | Interleukin | Height | Weight | BMI |
|--------------|------------|------------|-----------------|-------------|--------|--------|--------|---------------|------------|-----------------|-------------|--------|--------|--------|
| Left Peak E | P.C | -0.052 | -0.10 | -0.142 | 0.073 | -0.111 | -.323 | Left S' | 0.127 | 0.06 | -0.084 | 0.001 | 0.025 | 0.049 |
| | P-value | 0.669 | 0.411 | 0.24 | 0.55 | 0.359 | 0.006 | | 0.295 | 0.622 | 0.491 | 0.994 | 0.835 | 0.684 |
| Left Peak A | P.C | -0.097 | -0.027 | -0.078 | 0.082 | 0.062 | 0.004 | Left Peak E' | -0.002 | -0.16 | -0.126 | 0.055 | 0.099 | 0.096 |
| | P-value | 0.423 | 0.825 | 0.523 | 0.502 | 0.611 | 0.976 | | 0.986 | 0.185 | 0.297 | 0.653 | 0.415 | 0.427 |
| Right Peak E | P.C | -0.213 | -0.135 | -0.198 | 0.094 | 0.054 | -0.012 | Left Peak A' | 0.228 | 0.022 | -0.007 | -0.156 | -0.093 | 0.026 |
| | P-value | 0.076 | 0.264 | 0.1 | 0.437 | 0.655 | 0.92 | | 0.057 | 0.858 | 0.952 | 0.196 | 0.445 | 0.833 |
| Right Peak A | P.C | -0.118 | -0.056 | -0.236 | -0.074 | -0.034 | 0.006 | Right ET' | 0.446 | 0.132 | 0.026 | -0.347 | -0.374 | -.337 |
| | P-value | 0.331 | 0.644 | 0.049 | 0.543 | 0.78 | 0.962 | | <0.001 | 0.277 | 0.832 | 0.003 | 0.001 | 0.004 |
| Right ET | P.C | -0.108 | -0.087 | -0.002 | 0.199 | 0.15 | 0.033 | Right ICT' | -0.099 | 0.062 | -0.103 | 0.211 | 0.171 | 0.086 |
| | P-value | 0.374 | 0.473 | 0.987 | 0.099 | 0.216 | 0.787 | | 0.416 | 0.608 | 0.395 | 0.079 | 0.157 | 0.479 |
| Left ET | P.C | -0.055 | 0.04 | -0.073 | 0.394 | 0.313 | 0.074 | Right IRT' | -0.147 | 0.132 | -0.076 | 0.173 | 0.087 | -0.009 |
| | P-value | 0.649 | 0.744 | 0.551 | 0.001 | 0.008 | 0.544 | | 0.226 | 0.276 | 0.533 | 0.152 | 0.472 | 0.943 |
| IVSD | P.C | -.307 | 0.108 | -0.01 | 0.598 | 0.485 | 0.22 | Right S' | -0.368 | 0.087 | 0.086 | 0.506 | 0.398 | 0.145 |
| | P-value | 0.01 | 0.373 | 0.933 | <0.001 | <0.001 | 0.067 | | 0.002 | 0.473 | 0.477 | <0.001 | 0.001 | 0.230 |
| Age | P.C | -0.458 | 0.022 | 0.081 | 0.699 | 0.618 | .379 | Right Peak E' | 0.174 | 0.041 | 0.017 | -0.073 | -0.02 | 0.011 |
| | P-value | <0.001 | 0.86 | 0.504 | <0.001 | <0.001 | 0.001 | | 0.15 | 0.738 | 0.886 | 0.549 | 0.87 | 0.926 |
| LVDD | P.C | -0.231 | 0.026 | -0.024 | 0.419 | 0.39 | .270 | Right Peak A' | 0.223 | -0.058 | -0.125 | -0.216 | -0.187 | -0.094 |
| | P-value | 0.055 | 0.832 | 0.846 | <0.001 | 0.001 | 0.024 | | 0.063 | 0.636 | 0.302 | 0.073 | 0.121 | 0.438 |
| PWD | P.C | -0.143 | 0.072 | 0.256 | 0.257 | 0.158 | 0.005 | VS' | -0.028 | 0.075 | -0.108 | 0.115 | 0.055 | -0.054 |
| | P-value | 0.239 | 0.551 | 0.032 | 0.032 | 0.193 | 0.969 | | 0.816 | 0.535 | 0.372 | 0.343 | 0.653 | 0.655 |
| IVSS | P.C | -0.036 | -0.047 | -0.021 | 0.217 | 0.085 | -0.115 | VE' | 0.044 | -0.047 | -0.194 | -0.246 | -0.234 | -0.154 |
| | P-value | 0.768 | 0.702 | 0.866 | 0.071 | 0.485 | 0.345 | | 0.715 | 0.697 | 0.108 | 0.04 | 0.051 | 0.202 |

| | | | | | | | | | | | | | | |
|-----------|---------|--------|--------|--------|--------|--------|--------|------------|--------|--------|--------|--------|--------|--------|
| LVDS | P.C | -0.306 | 0.203 | 0.138 | 0.331 | 0.309 | .249 | VA' | -0.126 | -0.02 | -0.366 | 0.217 | 0.228 | 0.182 |
| | P-value | 0.01 | 0.092 | 0.254 | 0.005 | 0.009 | 0.038 | | 0.297 | 0.872 | 0.002 | 0.071 | 0.058 | 0.132 |
| PWS | P.C | -0.148 | 0.049 | 0.234 | 0.241 | 0.148 | 0.007 | Left MPI' | -0.307 | 0.096 | 0.057 | 0.354 | 0.277 | 0.149 |
| | P-value | 0.22 | 0.686 | 0.051 | 0.045 | 0.223 | 0.953 | | 0.01 | 0.43 | 0.64 | 0.003 | 0.02 | 0.217 |
| EF | P.C | -0.603 | -0.061 | -0.132 | 0.885 | 0.839 | .553 | Right MPI' | -0.362 | 0.001 | -0.106 | 0.402 | 0.355 | .236 |
| | P-value | <0.001 | 0.613 | 0.277 | <0.001 | <0.001 | <0.001 | | 0.002 | 0.994 | 0.384 | 0.001 | 0.003 | 0.05 |
| FS | P.C | -0.588 | -0.049 | -0.154 | 0.875 | 0.83 | .540 | LVMI | -0.334 | 0.119 | 0.101 | 0.579 | 0.494 | .268 |
| | P-value | <0.001 | 0.687 | 0.203 | <0.001 | <0.001 | <0.001 | | 0.005 | 0.325 | 0.405 | <0.001 | <0.001 | 0.025 |
| LVM | P.C | -0.295 | 0.133 | 0.074 | 0.518 | 0.45 | .255 | Left A/A' | -0.171 | -0.024 | -0.043 | 0.051 | -0.023 | -0.127 |
| | P-value | 0.013 | 0.272 | 0.545 | <0.001 | <0.001 | 0.033 | | 0.157 | 0.846 | 0.726 | 0.678 | 0.848 | 0.295 |
| Left ET' | P.C | 0.277 | -0.004 | -0.096 | -0.234 | -0.27 | -.288 | Left E/E' | -0.049 | 0.017 | -0.035 | 0.000 | -0.16 | -.309 |
| | P-value | 0.02 | 0.973 | 0.429 | 0.051 | 0.024 | 0.016 | | 0.686 | 0.888 | 0.774 | 0.998 | 0.185 | 0.009 |
| Left ICT' | P.C | -0.375 | 0.043 | 0.051 | 0.528 | 0.41 | 0.152 | Right E/E' | -0.229 | -0.085 | -0.125 | 0.079 | -0.015 | -0.11 |
| | P-value | 0.001 | 0.723 | 0.675 | <0.001 | <0.001 | 0.208 | | 0.057 | 0.482 | 0.302 | 0.516 | 0.904 | 0.365 |
| Left IRT' | P.C | 0.071 | 0.15 | 0.042 | -0.104 | -0.143 | -0.123 | Right A/A' | -0.185 | -0.015 | -0.029 | 0.075 | 0.018 | -0.07 |
| | P-value | 0.56 | 0.216 | 0.727 | 0.394 | 0.237 | 0.312 | | 0.126 | 0.9 | 0.81 | 0.536 | 0.881 | 0.564 |

P .C: Pearson Correlation, Peak E: Early mitral and tricuspid valve flow velocity, Peak A: Late mitral and tricuspid valve flow velocity, ET: Ejection time (for Aorta and Pulmonary), IVSD: Interventricular septal dimension in diastole, LVDD: Left ventricular end-diastolic dimension, PWD: Posterior wall dimension in diastole, IVSS: Interventricular septal dimension in systole, LVDS: Left ventricular diameter at systole, PWS: Posterior wall dimension in systole, EF: Ejection fraction, FS: Fractional shortening, LVM: Left ventricular mass, ICT: Isovolumetric contraction time, S': Systolic myocardial velocity above the baseline in pulmonic vein, mitral and tricuspid, E': early diastolic myocardial relaxation velocity below the baseline in pulmonic vein, mitral and tricuspid. A': myocardial velocity associated with atrial contraction in pulmonic vein, mitral and tricuspid, MPI: Myocardial Performance Index (Tei Index), BMI: Body Mass Index, LVMI: Left Ventricular Mass Index.