Iranian Journal of Numerical Analysis and Optimization Vol. 5, No. 2, (2015), pp 59-73

Application of modified simple equation method to Burgers, Huxley and Burgers-Huxley equations

Z. Ayati^{*}, M. Moradi and M. Mirzazadeh

Abstract

In this paper, modified simple equation method has been applied to obtain generalized solutions of Burgers, Huxley equations and combined forms of these equations. The new exact solutions of these equations have been obtained. It has been shown that the proposed method provides a very effective, and powerful mathematical tool for solving nonlinear partial differential equations.

Keywords: Modified simple equation method; Burgers equation; Huxley equation; Burger-Huxley equation.

1 Introduction

Mathematical modeling of many real phenomena leads to a non-linear partial differential equations in various fields of sciences and engineering. Many powerful methods have been presented for solving PDEs so far, such as tanh-function method [19] and [28], sine-cosine method [29], Homotopy Analysis method [17], Homotopy perturbation method [6], variational iteration method [9] and [10], Adomian decomposition method [1], Exp-function method [1], [11], [36] and [37], simplest equation method [7] and [4], and many others. Most recently, a modification of simplest equation method (MSE method) has been developed to obtain solutions of various nonlinear

M. Mirzazadeh

^{*}Corresponding author

Received 3 June 2014; revised 7 April 2015; accepted 17 June 2015 Z. Ayati

Department of Engineering Sciences, Faculty of Technology and Engineering East of Guilan, University of Guilan, Rudsar-Vajargah, Iran. e-mail: zainab.ayati@guilan.ac.ir; Ayati.zainab@gmail.com

M. Moradi

Department of Engineering Sciences, Faculty of Technology and Engineering East of Guilan, University of Guilan, Rudsar-Vajargah, Iran. email:

Department of Engineering Sciences, Faculty of Technology and Engineering East of Guilan, University of Guilan, Rudsar-Vajargah, Iran. email:

⁵⁹

evolution equations [14], [15], [21], [31], [32], [33] and [34]. The present paper is motivated by the desire to extend the MSE method to obtain generalized solutions of Burgers, Huxley, and Burgers-Huxley. The procedure of this method, by the help of Matlab, Maple or any mathematical package, is of utter simplicity.

2 The MSE method

Consider a nonlinear partial equation in two independent variables, say x and t, in the form of

$$P(u, u_t, u_x, u_{tt}, u_{xx}, ...) = 0,$$
(1)

where u = u(x,t) an unknown function, P is a polynomial in u = u(x,t)and its various partial derivatives, in which the highest order derivatives and nonlinear terms are involved. This method consists of the following steps.

Step 1. Using the transformation

$$\xi = x + wt, \tag{2}$$

where w is constant, we can rewrite equation (1) as a following nonlinear ODE:

$$Q(u, u', u'', ...) = 0.$$
(3)

Where the superscripts denote the derivatives with respect to ξ .

Step 2. Suppose that the solution of equation (2) can be expressed as follows

$$u(\xi) = \sum_{i=0}^{m} a_i (\frac{F'(\xi)}{F(\xi)})^i.$$
 (4)

Where a_i are constants to be determined later, with $a_m \neq 0$ and $F(\xi)$ is an unknown function to be determined later.

Step 3. The positive integer m can be determined by considering the homogeneous balance of the highest order derivatives and highest order nonlinear appearing in equation (2).

Step 4. Calculating all necessary derivatives u', u'', u''', \dots , and substituting equation (3) into equation (2) yields a polynomial of $\frac{F'(\xi)}{F(\xi)}$ and its derivatives. Equating the coefficients of same power of $F^{-i}(\xi)$ to zero gives a system of equations which can be used to solve for determining unknown constants, $F(\xi)$

and $F'(\xi)$. By substituting obtained results into equation (3), solutions of the equation (1) can be obtained.

3 Application of the MSE method

In this section, the modified simple equation method has been applied to obtain generalized solutions of Burgers, Huxley, and Burgers-Huxley.

3.1 Application MSE method to Burgers equation

The Burgers equation is a nonlinear partial differential equation of second order of the form

$$u_t + uu_x = \nu u_{xx}.$$
(5)

Where ν is the viscosity coefficient [2], [22], [23] and [27]. Many problems can be modeled by the Burgers' equation. This equation is one of the very few nonlinear partial differential equations which can be solved exactly for the restricted set of initial function. The study of the general properties of the equation has drawn considerable attention due to its place of application in some fields such as gas dynamics, heat conduction, elastically, etc.

To apply MSE method on equation (4), lets introduce a variable ξ , defined as

$$\xi = x - wt. \tag{6}$$

So, equation (4) turns to the following system of ordinary different equation,

$$-wu' + uu' = \nu u''. \tag{7}$$

Where w is constant to be determined. Integrating (7) and considering the integral constant to be zero, we obtain

$$-wu + \frac{1}{2}u^2 = \nu u'.$$
 (8)

Suppose that the solution of ODE equation (8) can be expressed by a polynomial in $\frac{F'}{F}$ as shown in (3). Balancing the terms u^2 and u' in equation (8), yields to m = 1. So we can write (3) as the following simple form

$$u(\xi) = a_1 \frac{F(\xi)}{F(\xi)} + a_0, a_1 \neq 0.$$
(9)

Z. Ayati, M. Moradi and M. Mirzazadeh

62 So

$$u' = a_1 \left(\frac{F''}{F} - \left(\frac{F'}{F}\right)^2\right). \tag{10}$$

Substituting (9) and (10) into equation (8) and equating each coefficient of $F^{-i}(\xi)$, (i=0,1,2) to zero, we derive

$$-wa_0 + \frac{1}{2}a_0^2 = 0, (11)$$

$$(-w+a_0)F'-\nu F''=0,$$
(12)

$$(\frac{1}{2}a_1^2 + \nu a_1)(F')^2 = 0.$$
(13)

By solving equations (11) and (13), the following results will be obtained

$$a_0 = 0, 2w, a_1 = -2\nu.$$

uation (12) turns to
 $wF' + \nu F'' = 0.$

So

So

and

Case 1. when eq

$$F' = A e^{\frac{-w}{\nu}\xi}.$$
 (14)

Where A is a arbitrary constant. Integrating (13) with respect ξ , $F(\xi)$ will be obtained as fallows

$$F = \frac{-A\nu}{w}e^{\frac{-w}{\nu}\xi} + B$$

where B is a constant of integration. Now, the exact solution of equation (4) has the form

$$u_1(x,t) = \frac{-2\nu A e^{\frac{-w}{\nu}(x-wt)}}{\frac{-A\nu}{w} e^{\frac{-w}{\nu}(x-wt)} + B}.$$

Case 2. when $a_0 = 2w$, equation (12) yields to

$$wF' - \nu F'' = 0.$$
$$F' = Ae^{\frac{w}{\nu}\xi},$$
$$F = \frac{A\nu}{w}e^{\frac{w}{\nu}\xi} + B$$

Now, the exact solution of equation (4) has the form

$$u_2(x,t) = 2w - \frac{2w\nu A e^{\frac{w}{\nu}(x-wt)}}{A\nu e^{\frac{w}{\nu}(x-wt)} + wB} = \frac{2w^2 B}{A\nu e^{\frac{w}{\nu}(x-wt)} + wB}.$$

Note that all obtained solutions have been checked with maple 13 by putting into the original equation and found correct.

3.2 Application MSE method to Huxley equation

Now we will bring to bear the MSE method to obtain exact solution to the Huxley equation [3], [7], [8], [12], [13], [18], [25] and [26] in the following form

$$u_t = u_{xx} + u(k - u)(u - 1).$$
(15)

The Huxley equation is an evolution equation that describes the nerve propagation in biology from which molecular CB properties can be calculated. It also gives a phenomenological description of the behaviour of the myosin heads II. This equation has many fascinating phenomena such as bursting oscillation [3], interspike [18], bifurcation, and chaos [35]. A generalized exact solution can gain an insight into these phenomena. There is no universal method for nonlinear equations. In this part, the exact solution will be obtained by the MSE method.

By considering (6), equation (15) turns to the following ordinary differential equation,

$$wu' + u'' + u(k - u)(u - 1) = 0.$$
 (16)

Balancing the terms u'' and u^3 in equation (16), yields to m = 1. So we can rewrite (3) as the following simple form

$$u(\xi) = a_1 \frac{F(\xi)}{F(\xi)} + a_0, a_1 \neq 0.$$
(17)

Now by substituting 19) into equation (16) and equating each coefficient of $F^{-i}(\xi)$, (i = 0, 1, 2, 3) to zero, the following result will be obtained

$$a_0^3 - (k+1)a_0^2 + ka_0 = 0, (18)$$

$${}_{1}F''' + wa_{1}F'' + (2(k+1)a_{0}a_{1} - 3a_{0}^{2}a_{1} - ka_{1})F' = 0,$$
(19)

$$-3a_1F'F'' + ((k+1)a_1 - wa_1 - 3a_0a_1^2)(F')^2 = 0,$$
 (20)

$$(2a_1 - a_1^3)(F')^3 = 0. (21)$$

Solving equations (15) and (22), we drive

 $a_0 = 0, 1, k,$ $a_1 = \pm \sqrt{2}.$

(24)

Case 1. when $a_0 = 0$, equations (16) and (17) yield

$$F''' + wF'' - kF' = 0, (22)$$

$$3F'' + (w - k - 1)F' = 0.$$
(23)

By substituting equation (23) into (22), we obtain

$$F''' + (w + \frac{3k}{w - k - 1})F'' = 0.$$

So

$$F'' = A e^{-\alpha\xi}. \tag{2}$$
 Where $\alpha = w + \frac{3k}{w-k-1}$ and A is a arbitrary constant. Therefore, we have

$$F' = -\frac{A}{\alpha}e^{-\alpha\xi} + B. \tag{25}$$

where A and B are arbitrary constants. By substituting (25) into equations (22) and (23), we get

$$w = \frac{k+1}{4} \pm \frac{3}{4}\sqrt{k^2 - 6k + 1}, B = 0.$$

Thus, (25) can be rewritten as follows

$$F' = -\frac{A}{\alpha}e^{-\alpha\xi}.$$
 (26)

Integrating (26) with respect ξ , $F(\xi)$ will be obtained as follows

$$F = \frac{A}{\alpha^2} e^{-\alpha\xi} + C,$$

where C is a constant of integration. Substituting the value of F and F' into equation (19), the following exact solution of equation (16) has been obtained

$$u_1(x,t) = \frac{\pm\sqrt{2}A\alpha e^{-\alpha\xi}}{Ae^{-\alpha\xi} + \alpha^2 C}$$

Case 2. when $a_0 = 1$, equations (16) and (17) turns to

$$F''' + wF'' + (k-1)F' = 0, (27)$$

$$3F'' + ((w - k - 1 \pm 3\sqrt{2})F' = 0.$$
⁽²⁸⁾

By substituting equation (8) into (27), we obtain

 $F''' + (w - \frac{3(k-1)}{w-k-1\pm 3\sqrt{2}})F'' = 0.$ $F'' = Ae^{-\alpha\xi},$ (29)

where $\alpha = w - \frac{3(k-1)}{w-k-1\pm 3\sqrt{2}}$ and A is a arbitrary constant. Therefore, we have

$$F' = -\frac{A}{\alpha}e^{-\alpha\xi} + B, \tag{30}$$

65

where A and B are arbitrary constants. By substituting (30) into equations (27) and (28), we get

$$w = \frac{k+1}{4} - \frac{3}{4}\sqrt{2} \pm \frac{3}{4}\sqrt{k^2 - 6k - 6k\sqrt{2} + 27 - 6\sqrt{2}}, B = 0,$$

$$k+1 - \frac{3}{4}\sqrt{2} \pm \frac{3}{4}\sqrt{k^2 - 6k - 6k\sqrt{2} + 27 - 6\sqrt{2}}, B = 0,$$

or

So

$$w = \frac{k+1}{4} + \frac{3}{4}\sqrt{2} \pm \frac{3}{4}\sqrt{k^2 - 6k + 6k\sqrt{2} + 27 + 6\sqrt{2}}, B = 0.$$

Thus, (30) can be rewritten as follows

$$F' = -\frac{A}{\alpha}e^{-\alpha\xi}.$$
(31)

Integrating (31) with respect, will be obtained as follows

$$F = \frac{A}{\alpha^2} e^{-\alpha\xi} + C,$$

where C is a constant of integration. Substituting the value of F and F' into equation (19), the following exact solution of equation (16) has been obtained

$$u_2(x,t) = 1 + \frac{\pm\sqrt{2}A\alpha e^{-\alpha\xi}}{Ae^{-\alpha\xi} + \alpha^2 C}.$$

Case 3. when $a_0 = k$, equations (16) and (17) turn to

$$F''' + wF'' + k(1 - k)F' = 0,$$

$$3F'' + ((w - 1 + k(-1 \pm 3\sqrt{2}))F' = 0)$$

By using the same method applied in case 1, the following solution will be obtained $\sqrt{2} 4 = -\alpha \xi$

$$u_3(x,t) = k + \frac{\pm\sqrt{2A\alpha}e^{-\alpha\xi}}{Ae^{-\alpha\xi} + \alpha^2 C},$$

www.SID.ir

Z. Ayati, M. Moradi and M. Mirzazadeh

where

$$\alpha = w - \frac{3k(1-k)}{w - 1 + k(-1 \pm 3\sqrt{2})}$$

and

$$w = \frac{1 - k(-1 \pm 3\sqrt{2})}{4} \pm \frac{3}{4}\sqrt{1 - 2k(-1 \pm 3\sqrt{2}) + k(-1 \pm 3\sqrt{2})^2 - 8k^2 + 8k}.$$

3.3 Application MSE method to Burgers-Huxley equation

The analysis presented in this part is based on the generalized nonlinear Burgers-Huxley equation,

$$u_t = u_{xx} + uu_x + u(k-u)(u-1),$$
(32)

which models the interaction between reaction mechanisms, convection effects and diffusion transports [20], and some special cases of the equation, which usually appear in mathematical modelling of some real world phenomena. It also gives a phenomenological description of the behaviour of the myosin heads II [30] and Fitzhugh-Nagoma equation, an important nonlinear reaction-diffusion equation used in circuit theory, biology and population genetics [5].

By considering (6), equation (32) turns to the following ordinary differential equation,

$$wu' + u'' + uu' + u(k - u)(u - 1) = 0.$$
(33)

Balancing the terms $u^{\prime\prime}$ and u^3 in Eq. (33), yields to m=1 . So we can rewrite (3) as the following simple form

$$u(\xi) = a_1(\frac{F'}{F}) + a_0, a_1 \neq 0.$$
(34)

Now by substituting (34) into equation (33) and equating each coefficient of $F^{-i}(\xi)$, (i = 0, 1, 2, 3) to zero, the following result will be obtained

$$a_0^3 - (k+1)a_0^2 + ka_0 = 0, (35)$$

$$a_1 F''' + (wa_1 + a_0 a_1) F'' + (2(k+1)a_0 a_1 - 3a_0^2 a_1 - ka_1) F' = 0, \qquad (36)$$

$$(-3a_1 + a_1^2)F'F'' + ((k+1)a_1 - wa_1 - a_0a_1 - 3a_0a_1^2)(F')^2 = 0, \quad (37)$$

$$(2a_1 - a_1^2 - a_1^3)(F')^3 = 0. (38)$$

Solving equation (35) and (38), we get

$$a_0 = 0, 1, k,$$

 $a_1 = 1, -2.$

Case 1. when $a_0 = 0$ and $a_1 = 1$ equations (36) and (37) yield

$$F''' + wF'' - kF' = 0, (39)$$

$$2F'' + (w - k - 1)F' = 0. (40)$$

By substituting equation (40) into (39), we obtain

$$F''' + (w + \frac{2k}{w - k - 1})F'' = 0.$$

$$F'' = Ae^{-\alpha\xi}$$
(41)

So

where $\alpha = w + \frac{2k}{w-k-1}$ and A is a arbitrary constant. Therefore, we have

$$F' = -\frac{A}{\alpha}e^{-\alpha\xi} + B, \qquad (42)$$

where A and B are arbitrary constants. By substituting (42) into equations (39) and (40), we get

$$w = \pm (k-1), B = 0,$$

 So

$$\alpha = -1, -k.$$

Thus, (42) can be rewritten as follows

1

$$F' = -\frac{A}{\alpha}e^{-\alpha\xi}.$$
(43)

Integrating (43) with respect ξ , $F(\xi)$ will be obtained as follows

$$F' = \frac{A}{\alpha^2} e^{-\alpha\xi} + C,$$

where C is a constant of integration. Substituting the value of F and F' into equation (34), the following exact solutions of equation (33) has been obtained

$$u_1(x,t) = \frac{Ae^{x-(k-1)t}}{Ae^{x-(k-1)t}+C},$$
$$u_2(x,t) = \frac{AKe^{k(x+(k-1)t)}}{Ae^{k(x+(k-1)t)}+k^2C},$$

Case 2. when $a_0 = 0$, and $a_1 = -2$ equations (36) and (37) yield

$$F''' + wF'' - kF' = 0, (44)$$

$$-5F'' + (w - k - 1)F' = 0.$$
(45)

By substituting equation (45) into (44), we obtain

$$F''' + (w + \frac{5k}{w - k - 1})F'' = 0.$$

By using the same method applied in case 1, the following solution will be obtained

$$u_{3,4}(x,t) = \frac{2A\alpha e^{-\alpha\xi}}{Ae^{-\alpha\xi} + \alpha^2 C}$$

where

$$\alpha = w + \frac{5k}{w - k - 1},$$

and

$$w = \frac{3}{8}(k+1) \pm \frac{5}{8}\sqrt{k^2 - 14k + 1}.$$

Case 3. when $a_0 = 1$ and $a_1 = 1$, equations (31) and (32) turn to

$$F''' + (w+1)F'' + (k-1)F' = 0,$$
(46)

$$2F'' + (w+3-k)F' = 0.$$
 (47)

By substituting equation (47) into (46), we obtain

$$F''' + (w + 1 - \frac{2(k-1)}{w+3-k})F'' = 0$$

So

$$F'' = Ae^{-\alpha\xi},$$

where $\alpha = w + 1 - \frac{2(k-1)}{w+3-k}$ and A is an arbitrary constant. Therefore, we have

$$F' = -\frac{A}{\alpha}e^{-\alpha\xi} + B, \tag{48}$$

where A and B are arbitrary constants. By substituting (48) into equations (46) and (47), we get

$$w = -1 \pm k, B = 0.$$

Thus, the following exact solutions of equation (33) has been obtained.

$$u_5(x,t) = 1 - \frac{Ae^{-x+(1-k)t}}{Ae^{-x+(1-k)t} + C},$$
$$u_6(x,t) = 1 - \frac{A(1-k)e^{-(1-k)(x+(1+k)t)}}{Ae^{-(1-k)(x+(1+k)t)} + (1-k)^2C}.$$

Case 4. when $a_0 = 1$ and $a_1 = -2$, by using the same method applied in case 1, the following solution will be obtained.

$$u_{7,8}(x,t) = 1 + \frac{2A\alpha e^{-\alpha\xi}}{Ae^{-\alpha\xi} + \alpha^2 C}.$$

$$\alpha = w + 1 - \frac{5(k-1)}{w - 6 - k},$$

and

Where

$$w = \frac{13+3k}{8}(k+1) \pm \frac{5}{8}\sqrt{k^2 + 30k + 33}.$$

Case 5. when $a_0 = k$ and $a_1 = 1$, equations (31) and (32) turn to

$$F''' + (w+k)F'' + k(1-k)F' = 0,$$
(49)

$$2F'' + (w + 3k - 1)F' = 0.$$
 (50)

By substituting equation (50) into (49), we obtain

$$F''' + (w + k - \frac{2k(1-k)}{w+3k-1})F'' = 0.$$

So

$$F'' = Ae^{-\alpha\xi},$$

where $\alpha = w + k - \frac{2k(1-k)}{w+3k-1}$ and A is an arbitrary constant. Therefore, we have

$$F' = -\frac{A}{\alpha}e^{-\alpha\xi} + B, \tag{51}$$

where A and B are arbitrary constants. By substituting (51) into equations (49) and (50), we get

$$w = -k \pm 1, B = 0.$$

Thus, the following exact solutions of equation (32) has been obtained

$$u_9(x,t) = k - \frac{Ake^{-k(x+(k-1)t)}}{Ae^{-k(x+(k-1)t)} + k^2C},$$

Z. Ayati, M. Moradi and M. Mirzazadeh

$$u_{10}(x,t) = k - \frac{A(k-1)e^{-(k-1)(x+(k+1)t)}}{Ae^{-(k-1)(x+(k+1)t)} + (k-1)^2C}.$$

Case 6. when $a_0 = k$ and $a_1 = -2$, by using the same method applied in case 5, the following solution will be obtained.

$$u_{11,12}(x,t) = k + \frac{2A\alpha e^{-\alpha\xi}}{Ae^{-\alpha\xi} + \alpha^2 C}.$$

Where

$$\alpha = w + k - \frac{5k(1-k)}{w - 1 - 6k},$$

and

$$w = \frac{13+3k}{8}(k+1) \pm \frac{5}{8}\sqrt{33k^2 + 30k + 1}.$$

4 Conclusion

In this paper, modified simple equation method has been applied to obtain the generalized solutions of some nonlinear partial differential equation. The results show that modified simple equation method is a powerful tool for obtaining the exact solutions of nonlinear differential equations. It may be concluded that, the method can be easily extended to all kinds of nonlinear equations. The advantage of this method over other methods is that in most methods applied for the exact solution of partial differential equations such as Exp-function method, $\frac{G'}{G}$ - expansion method, tanh-function method, and so on, the solution is presented in terms of some pre-defined functions, but in the MSE method, $F(\xi)$ is not pre-defined or not a solution of any pre-defined equation. Therefore, some new solutions might be found by this method. The computations associated in this work were performed by Maple 13.

References

- Biazar, J., Babolian, E., Nouri, A. and Islam, R. An alternate algorithm for computing Adomian Decomposition method in special cases, Applied Mathematics and Computation 38 (2003)523–529.
- 2. Burgers, M. The Nonlinear Diffusion Equation, Reidel, Dordtrecht, 1974.

- Duan, L. X. and Lu, Q. S. ursting oscillations near codimension-two bifurcations in the Chay Neuron model, International Journal of Nonlinear Sciences and Numerical Simulation 7(1) (2006) 59–64.
- 4. Ebaid, A. Exact solitary wave solutions for some nonlinear evolution equations via Exp-function method, Phys.Lett. A 365 (2007) 213–219.
- Hariharan, G. and Kannan, K. Haar wavelet method for solving FitzHugh-Nagumo equation, International Journal of Mathematical and Statistical Sciences 2 (2) (2010) 59–63.
- 6. He, J.H. Application of homotopy perturbation method to nonlinear wave equations, Chaos Solitons Fractals 26 (2005) 695–700.
- He, J. H. Modified Hodgkin-Huxley model, Chaos, Solitons, and Fractals 29(2) (2006) 303–306.
- He, J. H. Resistance in cell membrane and nerve fiber, Neuroscience Letters 373(1) (2005) 48–50.
- He, J. H. Variational iteration method for autonomous ordinary differential systems, Appl. Math. Comput. 114 (2000) 115–123.
- 10. He, J. H. Variational iteration method a kind of non-linear analytical technique: some examples, Int. J. Nonlinear Mech. 34 (1999) 699–708.
- 11. He, J. H. and Wu, X. H. Exp-function method for nonlinear wave equations, Chaos Solitons Fractals, 30 (2006) 700–708.
- He, J. H. and Wu, X. H. Modified Morris-Lecar model for interacting ion channels, Neurocomputing 64 (2005) 543–545.
- Hodgkin, A. L. and Huxley, A. F. A quantitative description of membrane current and its application to conduction and excitation in nerve, The Journal of Physiology 117(4) (1952) 500–544.
- Jawad, A. J. M., Petkovic, M. D. and Biswas, A. Modified simple equation method for nonlinear evolution equations, Appl. Math. Comput. 217 (2010) 869–877.
- khan, K. and Akbar, M. A. Exact and solitary wave solutions for the Tzitzeica-Dodd-Bullough and the modified Kdv-Zakharov-Kuznetsov equations using the modified simple equation method, Ain Shams Engineering Journal 4(4) (2013)903–909.
- Kudryashov, N. A. Nadejda B. Loguinova, Extended simplest equation method for nonlinear differential equations, Applied Mathematics and Computation 205 (2008) 396–402.

- 17. Liao, S. J. The proposed homotopy analysis techniques for the solution of nonlinear problems, Ph.D. dissertation. Shanghai: Shanghai Jiao Tong, University; 1992 [in English].
- Liu, S. Q., Fan, T. and Lu, Q. S. The spike order of the winnerless competition -WLC-model and its application to the inhibition neural system, International Journal of Nonlinear Sciences and Numerical Simulation 6(2) (2005) 133–138.
- Malfliet, W. and Hereman, W. The tanh method: I. Exact solutions of nonlinear evolution and wave equations, Phys Scr 54 (1996) 563-568.
- 20. Satsuma, J. Topics in soliton theory and exactly solvable nonlinear equations, Singapore:World Scientific (1987).
- Taghizadeh, N., Mirzazadeh, M., Samiei Paghaleh, A. and Vahidi, J. Exact solutions of nonlinear evolution equations by using the modified simple equation method, Ain Shams Engineering Journal 3 (2012) 321– 325.
- 22. Veksler, A. and Zarmi, Y. Wave interactions and the analysis of the perturbed Burgers equation, Physica D 211 (2005) 57-73.
- Veksler, A. and Zarmi, Y. Freedom in the expansion and obstacles to integrability in multiple-soliton solutions of the perturbed KdV equation, Physica D 217 (2006) 77–87.
- Vitanov N. K. Modified method of simplest equation:powerful tool for obtaining exact and approximate traveling-wave solutions of nonlinear PDEs, Commun Nonlinear Sci Numer Simulat 16 (2011) 1176–85.
- Wang, J., Chen, L. and Fei, X. Analysis and control of the bifurcation of Hodgkin-Huxley model, haos, Solitons, and Fractals 31(1) (2007) 247–256.
- Wang, J., Chen, L. and Fei, X. Bifurcation control of the Hodgkin-Huxley equations, Chaos, Solitons and Fractals 33(1) (2007) 217–224.
- Wazwaz, A. M. Analytic study on Burgers, Fisher, Huxley equations and combined forms of these equations, Applied Mathematics and Computation 195 (2008) 754–761.
- Wazwaz, A. M. The tanh method: Exact solutions of the Sine-Gordon and Sinh-Gordon equations, Appl. Math. Comput. 167 (2005) 1196–1210.
- Wazwaz A. M. The tanh and the sine-cosine methods for a reliable treatment of the modi.ed equal width equation and its variants, Commun Nonlinear Sci Numer Simul 11 (2006) 148–60.
- Wazwaz A. M. Travelling wave solutions of generalized forms of Burgers, Burgers-KdV and Burgers-Huxley equations, Applied Mathematics and Computation 169 (2005) 639–656.

- Zayed, E. M. E. A note on the modified simple equation method applied to Sharma-Tasso-Olver equation, Appl. Math. Comput. 218 (2011) 3962– 3964.
- 32. Zayed, E. M. E. The modified simple equation method for two nonlinear PDEs with power law and Kerr law nonlinearity, Pan American Mathematical Journal, International Publications USA 24(1) (2014) 65–74.
- Zayed, E. M. E. The modified simple equation method applied to nonlinear two models of diffusion-reaction equations, Journal of Mathematical Research and Applications 2(2) (2014) 5–13.
- 34. Zayed, E. M. E. and Ibrahim, S. A. H. Exact solutions of nonlinear evolution equations in mathematical physics using the modified simple equation method, Chinese Phys. Lett. 29 (6) (2012) p. 060201.
- Zhang, G. J., Xu, J. X., Yao, H. and Wei, R. Mechanism of bifurcationdependent coherence resonance of an excitable neuron model, International Journal of Nonlinear Sciences and Numerical Simulation 7(4) (2006) 447–450.
- 36. Zhang, S. Application of Exp-function method to a KdV equation with variable coefficients, Phys. Lett. A 365(2007) 448-453.
- 37. Zhu, S. D. *Exp-function method for the discrete mKdV lattice*, Int. J. Nonlinear Sci. Numer. Simul. 8 (2007) 465–469.

کاربرد روش معادله ساده توسعه یافته برای معادلات برگرز ، هوکسلی و هوکسلی برگرز

زینب آیاتی، مجتبی مرادی و محمد میرزازاده

دانشگاه گیلان، دانشکده فنی و مهندسی شرق گیلان، گروه علوم مهندسی

چکیده : در این مقاله، روش معادله ساده توسعه یافته برای به دست آوردن جواب-های معادلات برگرز، هوکسلی و شکل ترکیب شده آنها به کار رفته است. جوابهای دقیق جدیدی از این معادلات به دست آمده است.نشان داده شده است که روش ارائه شده یک ابزار ریاضی قوی و بسیار موثر برای حل معادلات با مشتقات جزئی می باشد.

كلمات كليدى : روش معادله ساده توسعه يافته؛ معادله برگرز؛ معادله هوكسلى؛ معادله برگرز-هوكسلى.

www.SID.ir