

## The Effect of Lateral Wedge Insole on Inter-joint Coordination during Walking in People with Medial Knee Osteoarthritis

Shima Hokmi<sup>1</sup>, Elaheh Azadian<sup>1\*</sup>, Mohammad Hasan Bijarchian<sup>2</sup>

1. Department of Motor Behavior, Faculty of Humanities, Islamic Azad University, Hamedan Branch, Hamedan, Iran.

2. Department of Sport Biomechanics, Faculty of Humanities, Islamic Azad University, Hamedan Branch, Hamedan, Iran.

### ABSTRACT

The purpose of this study was to investigate the features of inter-joint coordination in lower joints of osteoarthritis patients and compare them with healthy subjects. 10 patients with knee osteoarthritis and 10 healthy individuals volunteered to participate in this study. A Vicon 3D Motion Analysis System was used to record kinematic variables during walking under three conditions: (a) bare feet walking, (b) walking with shoes with a 5-degree lateral wedge insole and (c) 0-degree insole. In this research, the vector coding method was used to estimate inter-joint coordination. Repeated measurement and ANOVA were used for within and between group comparisons, respectively. The significant level was set at  $p < 0.05$  for all analysis. Within group comparisons in the osteoarthritis group showed that the overall use of insole did not significantly change the inter joints coordination ( $p > 0.05$ ). The results showed that difference between the control group with the injured foot in the osteoarthritis group occurred only in the mid-stance phase, and in the barefoot walking condition and walking with the 5-degree lateral wedge insole. According to the results of the present study, in addition to the necessity of treatment for the injured foot, the healthy foot should also be used treatment in patients with knee osteoarthritis.

**Keywords:** Osteoarthritis, Inter-joint coordination, Gait

### Introduction

Coordination in human movement is essential for organizing complex and redundant degrees of freedom of the musculoskeletal system [1]. In Dynamical Systems Theory, movement patterns are results of a synergistic organization of the neuromuscular system based on the constraints of anatomical structures, environmental factors, and movement tasks [2]. Precise control of human walking requires multi-joint coordination [3]. This includes higher levels of neuromuscular controls to regulate the initiation, intensity, and adaptability of locomotion [4].

Osteoarthritis (OA) is one of the most common causes of chronic disability in the older adults, which greatly affects their lives [5]. The tibiofemoral joint is commonly involved and OA changes are more prevalent in the medial compartment than the lateral compartment (nearly 10 times) [6, 7]. Studies in this area have clearly shown that knee OA not only changes the mechanics at the knee, but also at the other lower limb joints during functional activities [8, 9]. It has been shown that cartilage damage progresses due to high joint load in the medial compartment and that pain increases as the deterioration increases [10]. Therefore, reducing the joint load in the medial compartment of the knee acts to improve the disease or to relieve pain. There are several treatments for preventing pain and improving the mobility of patients. In the moderate to severe knee OA, surgical treatment is performed [11]. However, the use of an insole is recommended as a non-invasive method to reduce knee loading [10, 12], and therefore reducing pain [13-15] and increasing mobility [12] in OA patients.

The biomechanical changes associated with knee OA are well documented [7, 16], and the influence of knee OA on the inter-joint coordination during movement was investigated by wang et al. (2009) [5]. Their results showed that despite significant changes of the joint kinematics knee OA did not change significantly the way the motions of the lower limb joints are coordinated during obstacle-crossing [5]. According to the available

reports, the purpose of this study was to investigate the impact of insole on inter-joint coordination and to compare it with healthy individuals. We therefore assume that the use of insole can lead to increased coordination in the lower limb joints of patients with medial knee OA.

**Material and Methods**

The study population consisted of patients with medial knee OA who referred to Hamadan medical clinics and healthy individuals who volunteered to participate in the study. To determine the sample size with G\*Power software with  $\alpha= 0.05$  and statistical power= 0.80, 10 individuals were considered for each group. 10 patients with medial knee OA (OA group) and 10 matched healthy men participated as a control group. Participants in the OA group completed the Koos Quality Assessment Questionnaire, which included pain, quality of life, symptoms of illness, and daily activity. People who have had injuries (except for medial knee OA in the OA group) over the past six months in lower extremities or neurological (muscle disease) and orthopedic diseases (bone fracture, tendonitis, sprain, strain, and joint surgery) were excluded from the study. Subjects also completed the consent to participate in the test. The protocol of this study was approved by the Medical Ethics Committee of Hamadan University of Medical Sciences under IR.UMSHA.REC.1396.368 dated 29 July 2017.

A Vicon 3D Motion Analysis System (Oxford Metrics, Oxford, UK) with four T20 series cameras with a frequency of 100 Hz was used to record kinematic variables during walking under different conditions. Kinematic data were recorded and analyzed using Vicon Nexus software (v1.8.2) 1,8,2 and Polygan software (v3.5.2). A total of 16 retro-reflective markers were positioned on the skin overlying specific bony landmarks or anatomical positions of the lower body based on the Plug-In Gait Marker Set (Vicon Peak, Oxford, UK)[17]. Each participant was walked with self-selected speed. The subjects' task in the laboratory consisted of: (a) bare feet walking, (b) walking with shoes with a 5-degree lateral wedge insole and (c) 0-degree insole. Each of these conditions was repeated 6 times and the average value was used for statistical calculations.

In this research, the vector coding method was used to estimate inter-joint coordination. The relative motion between the two segments can be quantified with the coupling angle (Formula.1), an angle subtended from a vector adjoining two successive time points relative to the right horizontal [18].

Formula 1: 
$$\gamma_{j,i} = \tan^{-1} \left( \frac{y_{j,i+1} - y_{j,i}}{x_{j,i+1} - x_{j,i}} \right)$$

Where  $0^\circ \leq \gamma \leq 360^\circ$ , represents consecutive data points in a cycle (here expressed as a percent stance), and j identifies the multiple gait cycles. Once the corresponding vector is specified, this angle is denoted as the coupling angle. The mean coupling angle across the different gait cycles is again defined for each percentage (i) of the cycle (Formula.2) [18]:

Formula 2: 
$$\bar{\gamma}_i = \begin{cases} \arctan(\bar{y}_i / \bar{x}_i) & \text{if } \bar{x}_i \geq 0 \\ 180 + \arctan(\bar{y}_i / \bar{x}_i) & \text{if } \bar{x}_i \leq 0 \end{cases}$$

Shapiro-Wilk test was used to evaluate the normality of the data and the possibility of using parametric tests. The experimental design of this study included two intra-group factors: (a) factor of walking with three-level (bare feet walking, walking with 0- degree insole and 5-degree lateral wedge insole), and (b) factor of foot with two levels (OA foot in OA group and the healthy foot for the OA group and the factor of gait phase with three-step (loading response, mid-stance, and push off); all data were analyzed in three sagittal, frontal, and horizontal planes. In this study, within group factor with two levels of OA (individuals with medial knee osteoarthritis) and control group (healthy individuals) was considered. According to the research design,

repeated measure statistical method was used to compare within groups and ANOVA was used for comparison between groups. The calculations were done by SPSS (16.0 version), the statistical significance was assessed at  $p < 0.05$ .

**Results**

The mean and standard deviation of participants' characteristics are presented in Table 1. As can be seen, the control group did not differ significantly in terms of demographic characteristics from the OA group.

**Table 1.** Mean and standard deviations of subjects' demographic variables in groups

Variables	Groups		
	OA	Control	Sig.
Age	51.70±5.70	50.75± 3.23	0.72
Height	1.70±0.72	1.73±0.31	0.18
Mass	74.10±14.66	82.34±5.79	0.12
BMI	26.72±5.21	27.34±1.69	0.72
<b>Quality evaluation of koos</b>			
Pain (0-100)	46.82±15.78	-	-
Symptoms of Disease (0-100)	53.20±16.35	-	-
Daily Living Activity (0-100)	47.78±17.19	-	-
Quality of life (0-100)	35.49±21.27	-	-

Note: abbreviations: SD: standard deviation, BMI (Body Mass Index, age in year, height in meters, mass in kilograms).

Tables 2-4 show a descriptive analysis of the variables. In this study, the inter joints coordination in three gait conditions was investigated in barefoot and using two insoles with 0° insole and 5° lateral wedge insole. Within group comparisons in the OA group showed that the overall use of insole did not significantly change the inter joints coordination ( $p > 0.05$ ). Also, the factor of foot had a significant effect on a sagittal and horizontal plane in the knee-hip joint and also on the horizontal plane in the ankle-hip joint ( $p < 0.05$ ). Paired sample t-test results showed that there was a significant difference between injured and healthy legs in the mid-stance phase and in the ankle-hip joint. There were no significant differences in the other planes; injured and healthy foot. Therefore, there was no significant difference between the joints in both injured and healthy foot ( $p > 0.05$ ).

A comparison of the within group results showed that the difference between the control group with the injured foot in the OA group occurred only in the mid-stance phase, as shown in Table 1-3 with \*. The comparison between the control group and the healthy foot in the OA group also showed that there was a significant difference at the sagittal and horizontal planes in the mid-stance phase and between the knee-ankle joints as indicated in Table 1-3 with \*\*. In all cases, there was a difference between groups in barefoot walking or walking with a 5-degree lateral wedge insole.

**Table 2.** Mean and standard deviations of inter joint coordination in frontal plane in groups

Variables			OA group								
Plane	Joint	phase	Control group			OA Foot			Healthy Foot		
			N	D5	D0	N	D5	D0	N	D5	D0
Frontal	Ankle-Knee	L	119.5 (95.8)	116.5 (115.1)	98.4 (107.9)	106.2 (93.9)	130.4 (113.1)	103.4 (82.4)	110.1 (96.9)	76.1 (49.0)	58.7 (19.9)
		M	204.8 (43.2)	213.5 (45.5)	182.6 (42.3)	218.1 (74.9)	229.0 (40.7)	210.1 (53.2)	135.1 (99.7)	200.7 (52.4)	208.2 (67.0)
		P	76.6 (18.5)	76.1 (20.7)	75.0 (22.8)	97.6 (63.3)	69.9 (23.7)	66.8 (25.5)	102.3 (62.2)	99.0 (59.3)	83.6 (9.6)
		L	67.39 (22.5)	66.9 (12.6)	71.2 (9.8)	67.6 (61.1)	133 (112.2)	158.1 (117.6)	115.4 (104.5)	73.6 (45.3)	76.5 (59.6)
		M	224.3 (51.6)	240.6 (12.3)	222.5 (52.2)	168.7* (59.3)	194.8* (40.3)	208.5 (96.6)	192.8 (92.5)	199.5 (59.4)	183.3 (67.4)
		P	269.2 (112.7)	255.6 (91.8)	314.8 (26.7)	254.6 (110)	252.5 (115.2)	233.0 (115.0)	157.9 (129.5)	242.6 (109.4)	264.6 (91.9)
	Knee-Hip	L	90.1 (96.8)	59.8 (31.3)	74.3 (25.1)	47.2 (29.5)	139.5 (132.8)	134.5 (151.4)	128.2 (115.9)	81.1 (99.5)	67.8 (65.7)
		M	227.0 (70.8)	249.3 (37.8)	232.4 (66.7)	149.6* (63.9)	190.1* (63.7)	173.4 (44.3)	203.2 (90.3)	203.2 (76.9)	172.6 (67.1)
		P	212.5 (170.9)	267.7 (140.2)	296.5 (106.4)	267.8 (130.1)	241.4 (154.1)	272.7 (128.8)	197.4 (165.2)	265.4 (133.4)	279.3 (138.4)

**Table 3.** Mean and standard deviations of inter joint coordination in sagittal plane in groups

Variables			OA group								
Plane	Joint	phase	Control group			OA Foot			Healthy Foot		
			N	D5	D0	N	D5	D0	N	D5	D0
Sagittal	Ankle-Knee	L	88.9 (89.9)	73.9 (11.0)	92.9 (24.7)	68.6 (88.5)	83.6 (58.9)	96.5 (70.2)	87.1 (89.7)	78.1 (59.1)	60.7 (37.0)
		M	327.3 (25.2)	332.4 (14.1)	332.5 (12.3)	296.5* (164.7)	232.5 (152.9)	228.8 (136.6)	248.3 (149.5)	195.6** (146.5)	263.2 (128.3)
		P	122.2 (50.9)	109.3 (10.1)	105.2 (10.2)	107.8 (9.6)	108.9 (7.6)	96.4 (26.6)	111.3 (21.2)	114.8 (31.3)	103.9 (11.7)
		L	264.6 (82.4)	274.5 (18.9)	244.6 (46.2)	239.4 (108.3)	280.3 (26.7)	257.1 (55.5)	263.1 (58.7)	280.0 (25.7)	277.8 (30.9)
		M	260.7 (61.1)	284.0 (9.3)	288.5 (17.4)	263.1 (83.9)	286.2 (12.2)	266.4 (55.4)	261.7 (73.3)	282.2 (14.8)	288.0 (15.0)
		P	118.8 (20.35)	127.5 (10.8)	136.9 (31.5)	133.4 (25.2)	127.2 (10.6)	174.1 (86.5)	134.6 (34.6)	127.1 (27.2)	129.3 (30.6)
	Knee-Hip	L	302.5 (65.7)	320.8 (12.3)	291.0 (87.6)	236.9 (121.1)	304.5 (21.8)	247.5 (112.4)	266.3 (86.4)	298.5** (17.7)	308.8 (25.1)
		M	262.5 (27.5)	269.1 (3.1)	262.7 (18.0)	255.9 (59.1)	270.4 (8.2)	228.9 (87.4)	255.5 (53.5)	270.8 (8.1)	268.4 (14.1)
		P	44.0 (38.0)	28.3 (5.8)	94.4 (136.7)	61.2 (103.9)	29.6 (4.9)	92.0 (140.2)	62.1 (99.2)	63.7 (95.4)	64.3 (101.2)

**Table 4.** Mean and standard deviations of inter joint coordination in horizontal plane in groups

Variables			OA group								
Plane	Joint	phase	Control group			OA Foot			Healthy Foot		
			N	D5	D0	N	D5	D0	N	D5	D0
Horizontal	Ankle-Knee	L	158.4 (70.9)	135.0 (18.7)	147.5 (61.0)	126.6 (50.9)	128.8 (27.5)	133.4 (32.1)	142.8 (73.1)	141.4 (64.4)	139.3 (30.2)
		M	302.5 (104.8)	290.8 (100.5)	329.5 (17.1)	219.3 (148.1)	224.9 (150.3)	168.1 (143.8)	183.8** (130.2)	221.5 (150.5)	314.5 (56.9)
		P	155.4 (62.4)	109.5 (29.4)	116.3 (38.7)	115.0 (41.2)	139.7 (47.9)	108.4 (35.6)	140.5 (45.9)	120.3 (31.1)	120.0 (38.6)
		L	149.8 (48.5)	163.4 (25.0)	167.6 (35.3)	183.1 (94.6)	185.4 (38.9)	181.6 (52.8)	164.4 (82.1)	160.2 (63.4)	161.7 (40.5)
		M	66.0 (97.9)	41.1 (20.9)	23.2 (10.8)	132.4 (142.6)	191.4* (161.9)	188.7 (142.0)	101.1 (101.4)	124.4 (151.7)	70.4 (103.0)
		P	132.8 (46.1)	135.8 (51.1)	141.0 (49.6)	143.4 (57.5)	141.6 (27.5)	181.8 (66.8)	125.4 (49.9)	169.7 (51.6)	119.6 (40.0)
	Knee-Hip	L	80.4 (110.5)	64.2 (106.1)	122.2 (155.9)	153.2 (150.2)	198.8 (159.5)	199.2 (147.2)	112.5 (105.1)	114.0 (124.7)	105.5 (129.1)
		M	137.2 (51.5)	127.7 (27.4)	149.8 (17.3)	134.1 (92.8)	212.6* (63.7)	196.2 (83.3)	155.9 (104.6)	167.6 (62.3)	161.5 (39.1)
		P	84.7 (98.1)	95.2 (138.3)	116.3 (148.5)	108.4 (125.4)	37.1 (30.5)	119.1 (144.0)	92.9 (88.6)	95.1 (118.7)	79.8 (97.5)

## Discussion

The purpose of this study was to investigate the effect of insole on the lower limbs inter joints coordination in three dimensions in patients with medial knee OA during walking. The results of within group comparisons in the OA group showed that overall insole use did not cause a significant change in inter-joint coordination. Many studies have investigated the effect of insole or anti-pronation shoes on biomechanical variables during walking [19-22]. Jafarnejadgero et al., 2020 showed that prolonged use of the insoles improves coordination and coordination variability between lower extremity joints in people with flat foot during walking [23].

One possible reason for the difference in the results of the present study with Jafarnejadgero et al., study is that the present study was investigated the immediate effect of insole, while in that research was evaluated the effect of long-term use of the insole.

The results showed that there was a significant difference between injured and healthy foot in the mid-stance phase at the horizontal plane and ankle-hip joints. The difference was that the coordination between the ankle-hip joint on the frontal plane at the injured foot was anti-phase whereas in the healthy foot this coordination was in-phased. The injured foot is the movement of the hip toward adduction and the ankle motion toward the eversion, whereas in the healthy foot the thigh movement is towards the abduction and the ankle joint is directed toward the eversion. Increasing the hip joint adduction angle in the closed kinetic chain (lateral shift of the pelvis) can reduce injury of ankle eversion, caused by the ankle inversion, because it moves the center of mass more towards the lateral edge of the foot [24].

The findings of the present study showed that the ankle and hip joints have compensatory movements in the injured foot which can reduce the load on the joints. However, due to the pattern of coordination observed in the frontal plane between the ankle and hip joints during the mid-stance phase, there may be a long-term injury in the healthy foot. According to the results of the present study, in addition to the necessity of treatment for the injured foot, the healthy foot should also be used treatment in patients with knee OA.

However, the insole used in the present study did not show a change in the pattern of coordination between the lower extremity joints in both healthy and injured foot during walking in subjects with knee OA.

The present study has some limitations such as the low statistical sample size of the present study. On the other hand, in this study, only the immediate effect of using the insole was evaluated, while long-term use of insoles may show different results.

## Conclusion

According to the results of the present study, in addition to the necessity of treatment for the injured foot, the healthy foot should also be used treatment in patients with knee osteoarthritis. However, the insole used in the present study did not show a change in the pattern of inter joints coordination between the lower extremity in both healthy and injured foot during walking in subjects with medial knee osteoarthritis.

## Acknowledgement

We thank each of the subjects who participated in the study.

## References

1. Hafer, J.F. and K.A. Boyer, *Variability of segment coordination using a vector coding technique: reliability analysis for treadmill walking and running*. Gait & posture, 2017; **51**: 222-227.
2. Chiu, S.-L., *Assessing Inter-joint Coordination during Walking*. 2012.
3. Winter, D.A., *Foot trajectory in human gait: a precise and multifactorial motor control task*. Physical therapy, 1992; **72**(1): 45-53.
4. Krasovsky, T. and M.F. Levin, *Toward a better understanding of coordination in healthy and poststroke gait*. Neurorehabilitation and Neural Repair, 2010; **24**(3): 213-224.
5. Wang, T.-M., et al., *Bilateral knee osteoarthritis does not affect inter-joint coordination in older adults with gait deviations during obstacle-crossing*. Journal of Biomechanics, 2009; **42**(14): 2349-2356.
6. Schmitt, L.C. and K.S. Rudolph, *Influences on knee movement strategies during walking in persons with medial knee osteoarthritis*. Arthritis Care & Research, 2007; **57**(6): 1018-1026.
7. Thomas, R.H., et al., *Compartmental evaluation of osteoarthritis of the knee: a comparative study of available diagnostic modalities*. Radiology, 1975; **116**(3): 585-594.
8. Mündermann, A., C.O. Dyrby, and T.P. Andriacchi, *Secondary gait changes in patients with medial compartment knee osteoarthritis: increased load at the ankle, knee, and hip during walking*. Arthritis & rheumatism, 2005; **52**(9): 2835-2844.
9. Lu, T.-W., H.-L. Chen, and T.-M. Wang, *Obstacle crossing in older adults with medial compartment knee osteoarthritis*. Gait & Posture, 2007; **26**(4): 553-559.
10. Kuroyanagi, Y., et al., *The lateral wedged insole with subtalar strapping significantly reduces dynamic knee load in the medial compartment: Gait analysis on patients with medial knee osteoarthritis*. Osteoarthritis and cartilage, 2007; **15**(8): 932-93.
11. Maquet, P., *The treatment of choice in osteoarthritis of the knee*. Clinical orthopaedics and related research, 1985(192): 108-112.
12. Hinman, R.S., et al., *Lateral wedge insoles for medial knee osteoarthritis: effects on lower limb frontal plane biomechanics*. Clinical Biomechanics, 2012; **27**(1): 27-33.
13. Keating, E., et al., *Use of lateral heel and sole wedges in the treatment of medial osteoarthritis of the knee*. Orthopaedic review, 1993; **22**(8): 921-924.
14. Wolfe, S. and F. Breuckman, *Conservative management of Genu Valgus and Varum with medial/lateral Heel Wedges. 1991*. Indiana Med; **84**: 614-5.
15. Parkes, M.J., et al., *Lateral wedge insoles as a conservative treatment for pain in patients with medial knee osteoarthritis: a meta-analysis*. Jama, 2013;21;310(7):722-30.
16. Chen, H.-L., et al., *Biomechanical strategies for successful obstacle crossing with the trailing limb in older adults with medial compartment knee osteoarthritis*. Journal of Biomechanics, 2008; **41**(4): 753-761.



## Archive of SID

17. Majlesi, M., et al., *The effect of interventional proprioceptive training on static balance and gait in deaf children*. Research in Developmental Disabilities, 2014; **35**(12): 3562-3567.
18. Robertson, G.E., et al., *Research methods in biomechanics*. 2013: Human kinetics.
19. Alfuth, M., *Textured and stimulating insoles for balance and gait impairments in patients with multiple sclerosis and Parkinson's disease: A systematic review and meta-analysis*. Gait & Posture, 2017; **51**: 132-141.
20. Branthwaite, H.R., C.J. Payton, and N. Chockalingam, *The effect of simple insoles on three-dimensional foot motion during normal walking*. Clinical Biomechanics, 2004; **19**(9): 972-977.
21. Kerrigan, D.C., et al., *Effectiveness of a lateral-wedge insole on knee varus torque in patients with knee osteoarthritis*. Archives of Physical Medicine and Rehabilitation, 2002; **83**(7): 889-893.
22. Abdi, E., et al., *Evaluation of Appropriate Metatarsal Pad Length on Mechanical Parameters of Diabetic Foot Ulcer During Walking*. Journal of Sport Biomechanics, 2018; **4**(3): 38-51.
23. Jafarnejhadgero, A., et al., *Quantifying lower limb inter-joint coordination and coordination variability after four-month wearing arch support foot orthoses in children with flexible flat feet*. Human Movement Science, 2020; **1**;70:102593.
24. Kosonen, J., et al., *Effects of medially posted insoles on foot and lower limb mechanics across walking and running in overpronating men*. Journal of Biomechanics, 2017; **54**: 58-63.

**Corresponding Author:** Elaheh Azadian, Department of Motor Behavior, Faculty of Humanities, Islamic Azad University, Hamedan Branch, Hamedan, Iran.

Email: azadian1@yahoo.com

Tel.: +98 813 449 4042, Fax: +98 813 449 4042

چکیده فارسی

تأثیر کفی کفش با لبه خارجی بر هماهنگی بین مفصلی طی راه رفتن در افراد دارای استئوآرتریت داخلی

زانو

شیما حکمی<sup>۱</sup>، الهه آزادیان<sup>۱\*</sup>، محمد حسن بیجارچیان<sup>۱</sup>

۱- دانشکده علوم انسانی، دانشگاه آزاد اسلامی، واحد همدان، همدان، ایران.

هدف از این مطالعه بررسی هماهنگی بین مفاصل اندام تحتانی بیماران مبتلا به استئوآرتریت و مقایسه آن‌ها با افراد سالم بود. در این مطالعه ۱۰ بیمار مبتلا به استئوآرتریت داخلی زانو و ۱۰ فرد سالم و مشابه که داوطلب شرکت در این مطالعه بودند، شرکت کردند. برای ثبت متغیرهای کینماتیکی گام-برداری تحت شرایط مختلف از یک سیستم تحلیل حرکت وایکان استفاده شد. وظیفه افراد در آزمایشگاه شامل: (الف) گام‌برداری با پای برهنه، (ب) گام-برداری با کفش دارای کفی با لبه خارجی ۵ درجه و (ج) گام‌برداری با کفش دارای کفی با لبه خارجی صفر درجه بود. در این پژوهش، از روش برنامه-نویسی برداری برای برآورد هماهنگی بین مفاصل استفاده شد. مقایسه درون‌گروهی در گروه استئوآرتریت نشان داد که استفاده از کفی تأثیر معنی‌داری بر هماهنگی بین مفاصل ندارد ( $P > 0.05$ ). همچنین نتایج نشان داد که تفاوت بین گروه کنترل با پای آسیب‌دیده در گروه استئوآرتریت فقط در مرحله میداستانس رخ داده است، که در وضعیت گام‌برداری پابرهنه و راه رفتن با یک کفی ۵ درجه اختلاف معنی‌دار بوده است. با توجه به نتایج مطالعه حاضر، علاوه بر ضرورت درمان پای آسیب‌دیده، مداخلات درمانی برای پای سالم نیز باید در بیماران مبتلا به استئوآرتریت زانو نیز استفاده شود.

واژه‌های کلیدی: استئوآرتریت زانو، هماهنگی درون مفصلی، گام‌برداری