



First report of *Rhytidhysteron hysterinum* from Iran

S. A. Mousavi

S. A. Khodaparast ✉

S. Mousanejad

A. Rouhibakhsh

Department of Plant Protection, Faculty of Agricultural Sciences, University of Guilan, Rasht, Iran

Rhytidhysteron Speg. 1881 is a small genus of Dothideomycetes (Patellariales, Patellariaceae) comprising about 19 species (Wijayawardena et al. 2017). This genus includes saprobic to weakly plant pathogenic fungi that are found on woody plants (Soto-Medina et al. 2017). Members of *Rhytidhysteron* are clearly distinguished by large, elongate and boat-shaped ascomata, produced on branches and fallen wood (Soto-Medina et al. 2017). The genus shows wide distribution throughout the world (Soto-Medina et al. 2017), although there is no record from Iran.

During a study on fungi causing blight disease on boxwood (*Buxus sempervirens* L.), samples of dead twigs and decaying wood with ascoma of an attractive fungus were collected from the city of Fuman (Ghalehroodkhan), in the north of Iran, (49° 9' 54.74981"S, 37° 18' 16.58509"W), on June 2017. Isolation and purification of the fungus were done based on conventional methods and single spore cultures were obtained on WA 2%. The specimens were microscopically examined. Microscopic slides were prepared from asci and ascospores in 25% lactic acid. Both morphological and molecular characteristics were used in order to identify the isolate. Morphological study for identification was performed based on Boehm et al. (2009), Soto-Medina (2017) and Thambugala et al. (2016). Genomic DNA was extracted from mycelia produced on PDA using Chelex 5% (Walsh et al. 1991). The DNA amplification was obtained by polymerase chain reaction (PCR). Parts of large subunit (LSU) of rDNA was amplified as described by Khodaparast et al. (2012), using the primers LR5/LROR (Rehner & Samuels 1994, Vilgalys & Hester 1990). The nucleotide sequences of the PCR products were obtained using direct sequencing in an ABI 3730xl

sequencer (Applied Biosystems, USA). The obtained sequences were initially inspected manually and visually using MEGA7.0 (Kumar et al. 2016). Sequences were compared with the sequences available in the NCBI GenBank nucleotide database using a BLASTN search method.

Based on a BLAST search using partial LSU-sequences, the sequence of examined isolate showed 100% similarity to *Eutrybliidiella* (*Rhytidhysteron*) *hysterina* strain CBS 316.71 (MH871912) and *Rhytidhysteron hysterinum* (Dufour) Samuels & E. Müll. strain EB 0351 (GU397350) from GenBank. This result was in concordance with morphological characters of *R. hysterinum*. Examined specimens in this study had hysteriform, gregarious ascomata, closed at first, later opening to become apothecioid with incurved, striate laterally margins, with KOH extractable pigment, 1.7–3 × 0.8–1.4 mm wide, 0.7–1.2 mm high. Asci were bitunicate, cylindrical, 8-spored, 108–136.8 × 12–16.8 µm. Ascospores were ellipsoid with a rounded end, smooth, 1-septate, slightly constricted at the septa, thick-walled, 19.2–28.8 × 7.2–12 µm (Fig. 1). To our knowledge, *Rhytidhysteron hysterinum* is reported for the first time from Iran.

Specimens of *R. hysterinum* are deposited at the Fungarium of the Department of Plant Protection, Faculty of Agricultural Science, University of Guilan, Guilan, Iran (GUM 1546).

REFERENCES

- Boehm EWA, Mugambi GK, Miller AN, Huhndorf SM, Marincowitz S, Spatafora JW, Schoch CL. 2009. A molecular phylogenetic reappraisal of the Hysteriaceae, Mytiliniaceae and Gloniaceae (Pleosporomycetidae, Dothideomycetes) with keys to world species. *Studies in Mycology* 64:49–83.
- Khodaparast SA, Takamatsu S, Harada M, Abbasi M, Samadi S. 2012. Additional rDNA ITS sequences and its phylogenetic consequences for the genus

Submitted 10 Oct. 2018, accepted for publication 1 Dec. 2018

✉ Corresponding Author E-mail: Khodaparast@guilan.ac.ir

© 2018, Published by the Iranian Mycological Society

<http://mij.areeo.ac.ir>

Leveillula with emphasis on conidium morphology. Mycological Progress 11: 741–752.

Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis,

version 7.0. for bigger datasets. Molecular Biology and Evolution 33: 1870–4.




Fig. 1. *Rhytidhysteron hysterinum*. a. ascoma on wood; b. ascospores; c. vertical section through an ascoma, — Scale bars a = 1000 μ m, b = 20 μ m, c = 100 μ m.

Rehner SA, Samuels GJ. 1994. Taxonomy and phylogeny of Gliocladium analysed from nuclear large subunit ribosomal DNA sequences. Mycological Research 98: 625–634.

Soto-Medina E, Lücking R. 2017. A new species of *Rhytidhysteron* (Ascomycota: Patellariaceae) from Colombia, with a provisional working key to known species in the world. Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales 41: 59–63.

Thambugala KM, Hyde KD, Eungwanichayapant PD, Romero AI, Liu ZY. 2016. Additions to the genus *Rhytidhysteron* in Hysteriaceae. Cryptogamie Mycologie 37: 99–116.

Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172:4238–4246.

Walsh PS, Metzger DA, Higuchi R. 1991. Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. Biotechniques 10: 506–513.

Wijayawardene NN., Hyde KD, Rajeshkumar KC, Hawksworth DL, Madrid H, Kirk PM, Braun U, Singh RV, Crous PW, Kukwa M, Luecking R. 2017. Notes for genera: Ascomycota. Fungal Diversity 86:1–594.