



Received:
20 July, 2016

Accepted:
8 August, 2016

Published:
9 August, 2016

Subject Editor:
Kambiz Minaei

Contributions to the leafhoppers (Auchenorrhyncha: Cicadellidae) of Khuzestan, southwest of Iran

Farzad Pakarpour Rayeni^{1*}, Ali Asghar Seraj¹ and Jamasb Nozari²

¹ Department of Plant Protection, Faculty of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Iran.

² Department of Plant Protection, Faculty of Agriculture, University of Tehran, Karaj, Iran.

ABSTRACT. A faunal study of leafhoppers (Hemiptera: Auchenorrhyncha: Cicadellidae) was conducted in some parts of Khuzestan province (southwest of Iran) in 2014 and 2015. Totally, 90 species belonging to 49 genera and seven subfamilies were collected and identified. Most of samples were collected by using sweeping net and Malaise trap and a few by aspirator and sticky yellow trap. As a result of this study 47 species are recorded from Khuzestan province for the first time and eight species are new records for the fauna of Iran including *Euscelidius schenckii* (Kirschbaum, 1868); *Hardya (Hardya) melanopsis* (Hardy, 1850); *Paradorydium (Paradorydium) lanceolatum* (Burmeister, 1839); *Hecalus prasinus* (Matsumura, 1905); *Psammotettix emarginatus* Singh, 1969; *Grammacephalus pallidus* (Linnauvori, 1978); *Helionidia punctulata* (Ramakrishnan & Menon, 1974) and *Eupteryx (Eupteryx) kaghanensis* Ahmed, 1969. For each species geographical distribution in Iran are completely mentioned. In addition morphological data and illustration of adults and their male genitalia are given for the eight newly recorded species.

Key words: Leafhoppers, Fauna, Khuzestan, Distribution, new records, Iran.

Citation: Pakarpour Rayeni, F., Seraj, A.A. and Nozari, J. 2016 Contributions to the leafhoppers (Auchenorrhyncha: Cicadellidae) of Khuzestan, southwest of Iran. *Journal of Insect Biodiversity and Systematics*, 2(2): 229–257.

Introduction

Based on the number of described species, Cicadellidae (commonly named leafhoppers) is the largest family of the suborder Auchenorrhyncha with more than 20,000 described species (Dietrich, 2008). This family may be distinguished from other Hemiptera by the following combination of characters: the presence of rows of spin-like setae on their hind tibiae; postclypeus enlarged; antennal pedicel small; flagellum

aristiform, tegulae absent; forewing anal veins usually separate from base to wing margin; middle coxa small and narrowly separated (Anufriev and Emeljanov 1988; Dietrich 2005). The most important character to identify and classify nearly all species and most genera is the male genitalia. Character of similar value are almost completely lacking in female (Dakhil *et al.* 2011). In fact, the aedeagus is the most

Corresponding author: Farzad Pakarpour Rayeni, E-mail: pakarpour@ut.ac.ir

Copyright © 2016, Pakarpour Rayeni *et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

consistently used character in leafhoppers species differentiation (Ribaut 1936; 1952). Relatively, few species are economically important but there are some major pest, such as potato leafhopper, *Empoasca fabae* (Harris, 1841) and sugar beet leafhopper, *Neoliturus (Circulifer) tenells* (Baker, 1896) (Dietrich, 2005). Leafhoppers injure plants either directly through feeding (Backus 1988; Backus *et al.* 2005) or indirectly through transmission of plant pathogen, such as phytoplasmas and viruses (Nielson 1968; Maramorosch and Harris 1979; Wientraub and Beenland 2006). The fauna of leafhoppers in Iran is very diverse, the famous Czechoslovakian entomologist Dr. Jiri Dlabola, who worked on the Iranian Auchenorrhyncha fauna in the 1960s and 1970s, and described more than 100 new leafhoppers species (Dlabola 1960, 1963, 1964, 1965, 1971, 1972, 1974, 1977, 1979, 1980, 1981, 1984, 1987, 1994; Dlabola and Heller, 1962; Dlabola and Jankovich, 1981). After Dlabola's publications, new data on the Iranian leafhoppers were scarce and consist a few faunistic studies concerning leafhoppers have been conducted that most of them published in Conference Proceedings (Karimzadeh *et al.* 1998; Haghshanas and Khajeali 2000; Nematollahi and Khajehali 2000; Khajeali *et al.* 2001; Darvish-Mojeni and Taghizadeh 2008; Mozaffarian and Taghizadeh 2010; Olyaei Torshiz *et al.* 2011; Pakarpour *et al.* 2011a, b, c; Pakarpour and Nozari 2011; Moosavi Mahvelati and Modarres Awal 2011; Taghizadeh 2012; Khormali 2013; Abdollahi *et al.* 2013; 2014a, b, c; 2015; Rezaei *et al.* 2014; Pakarpour and Nozari 2014a, b; Pakarpour *et al.* 2014a, b; Pakarpour Rayeni *et al.* 2015 a, b, c. A checklist was provided for the subfamily Deltocephalinae by Pakarpour Rayeni *et al.* (2015a) and another one was published for all recorded species for the family Cicadellidae by Mozaffarian and Wilson

(2016) which presented the occurrence of 390 species.

Khuzestan province is located in southwest of Iran and covers 64,057 km² (Mossadegh and Kocheili 2003). This province has border with Ilam and Lorestan province in north, Isafahan, Chaharmal and Bakhtiari and Kohgiluyeh and Boyerahmad provinces in east, Bushehr province and Persian Gulf in south and with Iraq in west (Fig. 1). Abundance of water and fertile soils in Khuzestan favored the condition for existence of the various vegetation (Mossadegh and Kocheili 2003). However, southwest of Iran contains high insect diversity (Mossadegh and Kocheili 2003; Ramezani *et al.* 2006) but there is inadequate knowledge relating to the Cicadellids from this region. The objective of the current study is to report new data about occurrence of leafhoppers in this region as well as new records for fauna of Iran.

Material and methods

Leafhopper specimens were mostly collected with standard sweeping net and Malaise trap and a few by using aspirator and sticky yellow trap from tree locations in Khuzestan province with different climate, during 2014-2015. Sampling localities were selected to cover different types of habitats for collecting maximum occurring species (Table 1). Materials were killed by dropping into standard killing jar containing ethyl acetate and then mounted on the points, labeled and put into collection boxes.

The identifications were done according to the keys of Ribaut (1936; 1952), Le Quesne (1965; 1969), Le Quesne and Payne (1981), Anufriev and Emeljanov (1988), Della Guistica (1989), Biedermann and Niedringhaus (2009) and for endemic species were identified by using original description from Dlabola (1957, 1960, 1963, 1964, 1974, 1979, 1984, 1994).

Table 1. Coordinates, altitudes, localities and date of sampling in Khuzestan province

Localities	Coordinates	Altitude	Date of Sampling
Ahvaz I (Shahid Chamran University)	31°18'13.5"N, 48°39'40.3"E	17 m	13.iii.2014
Ahvaz II (Om-el-Tamir)	31°15'18.2"N, 48°32'38.3"E	17 m	07.ix.2014
Ahvaz III (Kut-e-Abdollah)	31°13'00.2"N, 48°41'42.3"E	15 m	26.iv.2015
Ahvaz IV (Mollasani)	31°34'11.3"N, 48°53'26.2"E	18 m	25.vii.2015
Bagh-e Malek I (Manjanigh Park)	31°31'06.5"N, 49°53'29.1"E	920 m	16.iii.2014
Bagh-e Malek II (Ghale-Tal)	31°32'04.8"N, 49°51'58.7"E	950 m	09.ix.2014
Bagh-e Malek III (Laleb village)	31°32'35.8"N, 49°58'26.7"E	1000 m	15.v.2015
Bagh-e Malek IV (10km, road to Ramhormoz)	31°30'38.2"N, 49°53'34.5"E	910 m	28.vii.2015
Dezfoul I (Andimeshk, Dezful Road)	32°22'29.2"N, 48°27'13.0"E	140 m	14.iii.2014
Dezfoul II (Sanjar village)	32°23'22.7"N, 48°21'13.9"E	140 m	12.ix.2014
Dezfoul III (Zaviyeh Mashali village)	32°17'36.8"N, 48°18'13.3"E	120 m	27.iv.2015
Dezfoul IV (Zaviyeh Hamudi village)	32°17'38.0"N, 48°19'12.3"E	120 m	27.vii.2015

Final confirmation was done mostly by Dr. Michael R. Wilson (National Museum of Wales, Cardiff) and a few by Dr. Christopher H. Dietrich (Illinois Natural History Survey) and Dr. James N. Zahniser (USDA-APHIS-PPQ, San Diego, California). The specimens were deposited in Jalal Afshar Zoological Museum at University of Tehran (JAZMUT). The identified species are ordered based on subfamilies, tribes and genera alphabetically (Emeljanov 1999; Dmitriev 2016; Zahniser 2016). The species marked with an asterisk are recorded for the first time from Khuzestan province and those marked with two asterisks are newly recorded for Iran. Abbreviation for the collector as follow: F.P.R.: Farzad Pakarpour Rayeni.

Results

Family Cicadellidae

This family includes 390 valid species belonging to 13 subfamilies in Iran (Mozaffarian and Wilson 2016).

Subfamily Aphrodinae

This subfamily includes three species in three genera in Iran (Mozaffarian and Wilson 2016).

Aphrodes bicincta (Schrank, 1776)*

Material examined: Bagh-e Malek, 1♂, 1♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Azarbajian-e Sharghi, Golestan, Guilan, Khorasan-e Razavi and Tehran provinces; north and northwest of Iran (Dlabola 1981; Mirzayans 1995; Abdollahi et al. 2015; Mozaffarian and Wilson 2016).

Subfamily Deltcephalinae

Deltcephalinae is the largest subfamily of Cicadellidae comprising more than 6700 valid species (Zahniser 2016) of which 232 extant species were recognized in Iran (Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Tribe Athysanini

45 species belonging to 15 genera have been recorded from Iran (Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Euscelidius schenckii (Kirschbaum, 1868)**

(Fig. 2a)

Material examined: Dezful, 1♂, 3♀, swept on weeds in citrus orchards, 14.iii.2014, leg.:

F.P.R.; Bagh-e Malek, 1♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Bagh-e Malek, 1♂, swept on *Medicago sativa*, 09.ix.2014, leg.: F.P.R.; Dezful, 1♀, swept on gramineae, 12.ix.2014, leg.: F.P.R.; Ahvaz, 3♂, 7♀, swept on weeds in palm orchards, 26.iv.2015, leg.: F.P.R.

Distribution in Iran: This species is a new record for the Iranian leafhoppers fauna.

Remarks: Male: Length, 4.5 mm.; Robust; Base color greyish yellow, frontoclypeus with two longitudinal and series of transverse dark brown streaks, vertex with irregular spots, fore wings longer than abdomen, marked with fine irregular dark brown dots which can locally be more or less fused into small patches; in male shaft of aedeagus up to middle more or less straight and then curved, apically with appendages like outwards directed hook (Fig. 3a).

Euscelidius mundus (Haupt, 1927)

Material examined: Bagh-e Malek, 1♂, 4♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Bagh-e Malek, 5♂, 12♀, swept on grasslands, 09.ix.2014, leg.: F.P.R.; Ahvaz, 2♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Sharghi, Bushehr, Fars, Guilan, Qom, Hormozgan, Khorasan-e Razavi, Kerman, Khuzestan and Tehran provinces; northwest, north and northeast, south and southwest of Iran (Dlabola 1965, 1972; Nast 1972; Mirzayans 1995; Pakarpour *et al.* 2011a, 2011c; Moosavi Mahvelati and Modarres Awal 2011; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Euscelis alsius Ribaut, 1952*

Material examined: Dezful, 3♂, 3♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Bagh-e Malek, 1♂, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Bagh-e malek, 1♂, 2♀, swept on grasslands,

09.ix.2014, leg.: F.P.R.; Ahvaz, 3♂, 4♀, swept on *Medicago sativa*, 26.iv.2015, leg.: F.P.R.; Ahvaz, 1♂, 1♀, swept on *Medicago sativa* in palm orchards, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Fars, Kerman and Tehran provinces; north, northwest, west and southwest of Iran (Dlabola 1971, 1981; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Pakarpour *et al.* 2011a, 2011c; Pakarpour Rayeni *et al.* 2015a; Abdollahi *et al.* 2014b; 2015; Mozaffarian and Wilson 2016).

Euscelis incisus (Kirschbaum, 1858)*

Material examined: Dezful, 1♂, swept on gramineae, 12.ix.2014, leg.: F.P.R.; Ahvaz, 1♂, swept on weeds in palm orchards, 26.iv.2015, leg.: F.P.R.

Distribution in Iran: Qom, Hormozgan, Isfahan, Kerman and Tehran provinces; north, center and south of Iran (Dlabola 1960; Nast 1972; Modarres Awal 1994; Mirzayans 1995; Nematollahi and Khajehali 2000; Haghshenas and Khajehali 2000; Khajehali *et al.* 2001; Pakarpour *et al.* 2011a, 2011c; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Euscelis (Euscelis) lineolatus Brullé, 1832*

Material examined: Dezful, 1♂, 1♀, swept on gramineae, 12.ix.2014, leg.: F.P.R.; Ahvaz, 3♂, 7♀, swept on weeds in palm orchards, 26.iv.2015, leg.: F.P.R.

Distribution in Iran: Isfahan; Khorasan-e Razavi and Khorasan-e Shomali provinces; northeast and center of Iran (Dlabola 1981, 1984; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Hardya (Hardya) iranicola Zachvatkin, 1946*

Material examined: Bagh-e Malek, 1♂, 3♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Ahvaz, 1♂, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Dezful, 1♂,

swept on weeds in citrus orchards, 12.ix.2014, leg.: F.P.R.

Distribution in Iran: Azarbaijan-e Sharghi, Guilan, Kerman, Kordestan, Qom, Tehran provinces; north, northwest and south of Iran (Dlabola 1981; Mirzayans 1995; Pakarpour et al. 2011a, 2011c; Abdollahi et al. 2015; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Hardya (Hardya) melanopsis (Hardy, 1850)** (Fig. 2b)

Material examined: Dezful, 1♂, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.; Bagh-e Malek, 1♂, 1♀, swept on grasslands, 15.v.2015, leg.: F.P.R.

Distribution in Iran: This species is a new record for the Iranian leafhoppers fauna.

Remarks: Male: Length, 2.8-3 mm.; Base color dark brown, face black with interrupted transverse light brown bands on fronto-clypeus, vertex brownish with irregular black streak along anterior margin and back marking posteriorly in form of outline of clover-leaf with median lobe bisected; aedeagus hind margin of shaft only apically and basally curved (Fig. 3b).

Platymetopius (Platymetopius) coronatus Dlabola, 1974*

Material examined: Bagh-e Malek, 1♂, 1♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz Mountains and north of Iran (Dlabola 1974, 1981; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Platymetopius (Platymetopius) cruentatus Haupt, 1927

Material examined: Ahvaz, 2♂, 3♀, 2 nymph, swept on weeds in palm orchards in Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Ahvaz, 1♂, 2♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Ahvaz, 2♂, 1 nymph, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Fars, Hormozgan, Kerman, Khuzestan, Sistan-o Baluchestan provinces; north, southwest, south and southeast of Iran (Dlabola 1960, 1981; Nast 1972; Mirzayans et al. 1976; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Abaii 2000; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Platymetopius (Platymetopius) exalbescens Dlabola, 1974*

Material examined: Bagh-e Malek, 1♂, swept on *Medicago sativa*, 09.ix.2014, leg.: F.P.R.; Dezful, 1♂, 1♀, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.

Distribution in Iran: Fars, Kohgiluyeh and Boyerahmad; southwest of Iran (Dlabola 1974; Mirzayans 1995; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Platymetopius (Platymetopius) rostratus (Herrich-Schäffer, 1834)*

Material examined: Ahvaz, 3♀, swept on weeds in palm orchards in Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Dezful, 1♂, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Bagh-e Malek, 6♂, 5♀, swept on *Medicago sativa*, 09.ix.2014, leg.: F.P.R.; Ahvaz, 2♂, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.; Dezful, 1♂, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

Distribution in Iran: Chaharmahal-o Bakhtiari, Isfahan, Kerman and Tehran provinces; south of central region of Alborz mountains, southeast and south of Iran (Dlabola 1981; Mirzayans 1995; Haghshenas and Khajehali 2000; Khajehali et al. 2001; Pakarpour et al. 2011a, 2011c; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Platymetopius (Platymetopius) shirazicus Dlabola, 1974*

Material examined: Bagh-e Malek, 1♂, swept on grasslands, 09.ix.2014, leg.: F.P.R.;

Dezful, 1♂, 1♀, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Sharghi, Fars, Golestan, Guilan, Mazandaran and Tehran provinces; north, northwest, west, southwest and center (Dlabola 1974, 1981; Mirzayans *et al.* 1976; Rajabi 1991; Modarres Awal 1994; Mirzayans 1995; Abaii 2000; Abdollahi *et al.* 2015; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Tribe Chiasmini

15 species belonging to six genera have been recorded from Iran (Pakarpour Rayeni *et al.*, 2015a; Mozaffarian and Wilson 2016).

Aconurella nuristana (Dlabola, 1957)

Material examined: Dezful, 4♂, 3♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.

Distribution in Iran: Khuzestan province (Pakarpour Rayeni *et al.* 2015c).

Aconurella prolixa (Lethierry, 1855)

Material examined: Bagh-e Malek, 2♂, 11♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Gharbi, Azarbaijan-e Gharbi, Fars, Kerman, Khuzestan, Sistan-o Baluchestan and Tehran provinces; north, northwest, west, southwest, south and center (Dlabola 1963, 1965; 1971, 1981; Nast 1972; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Pakarpour *et al.* 2011a; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Chiasmus sp.

Material examined: Ahvaz, 2♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Exitianus capicola (Stål, 1855)

Material examined: Dezful, 1♂, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Dezful, 1♂, swept on gramineae,

12.ix.2014, leg.: F.P.R.; Dezful, 3♂, 6♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.; Bagh-e-Malek, 3♂, 2♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Fars, Hormozgan, Kerman, Khuzestan, Sistan-o Baluchestan provinces; southwest, south and southeast (Dlabola 1960, 1963, 1965, 1981; Nast 1972; Mirzayans 1995; Pakarpour *et al.* 2011a; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Exitianus indicus (Distant, 1908)*

Material examined: Ahvaz, 1♂, swept on weeds in palm orchards in Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Dezful, 2♂, 1♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Bagh-e Malek, 3♂, 4♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Ahvaz, 4♂, 5♀, swept on *Medicago sativa*, 26.iv.2015, leg.: F.P.R.; Ahvaz, 2♂, swept on *Medicago sativa* in palm orchards, 25.vii.2015, leg.: F.P.R.; Dezful, 8♂, 11♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

Distribution in Iran: Kerman province (Pakarpour and Nozari 2014a; Pakarpour *et al.* 2015a).

Exitianus nanus (Distant 1908)

Material examined: Dezful, 5♀, swept on weeds in palm orchards, 14.iii.2014, leg.: F.P.R.; Bagh-e Malek, 1♂, 3♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Ahvaz, 1♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Dezful, 1♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

Distribution in Iran: Bushehr, Fars, Guilan, Hormozgan, Kerman, Kordestan, Khuzestan, Lorestan, Mazandaran, Sistan-o Baluchestan and Tehran provinces; center, southwest, south and southeast (Dlabola 1960, 1963, 1964, 1972, 1981; Nast 1972; Mirzayans *et al.* 1976; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Pakarpour *et al.* 2011a; Pakarpour and Nozari 2014a; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Tribe Cicadulini

10 species belonging to six genera have been recorded from Iran (Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Cicadula (Cicadula) lineatopunctata (Matsumura, 1908)*

Material examined: Dezful, 1♂, 1♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Sharghi, Guilan, Hormozgan, Isfahan, Kerman, Sistan-o Baluchestan and Tehran provinces; north, center, southwest, south and southeast (Dlabola 1981; Mirzayans 1995; Karimzadeh et al. 1998; Khajehali et al. 2001; Pakarpour et al. 2011a, 2011c; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Elymana sulphurella (Zetterstedt, 1828)*

Material examined: Ahvaz, 1♂, swept on *Medicago sativa*, 26.iv.2015, leg.: F.P.R.

Distribution in Iran: Kerman and Tehran provinces; south of central region of Alborz mountains and south of Iran (Mirzayans 1995; Pakarpour et al. 2011a; 2011c; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Mocydiopsis parvicauda Ribaut, 1939*

Material examined: Dezful, 1♂, 1♀, swept on gramineae, 12.ix.2014, leg.: F.P.R.; Dezful, 2♀, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.; Ahvaz, 1♂, 1♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Kerman province; north and south (Dlabola 1981; Pakarpour et al. 2011a; 2011c; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Proceps acicularis Mulsant and Rey, 1855*

Material examined: Bagh-e Malek, 2♂, swept on grasslands, 09.ix.2014, leg.: F.P.R.

Distribution in Iran: Hormozgan and Kerman provinces; south of Iran (Dlabola 1984; Mirzayans 1995; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Tribe Deltocephalini

Six species belonging to two genera have been recorded from Iran (Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Maiestas horvathi (Then, 1896)

Material examined: Ahvaz, 1♂, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Bagh-e Malek, 1♂, 2♀, swept on grasslands, 15.v.2015, leg.: F.P.R.

Distribution in Iran: Guilan, Hormozgan, Kerman, Khorasan-e Razavi and Khuzestan provinces; south, southwest and north (Dlabola 1981; Mirzayans 1995; Pakarpour et al. 2011a; 2011c; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Maiestas trifasciata (Lindberg, 1954)*

Material examined: Dezful, 1♂, 1♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.

Distribution in Iran: Hormozgan, Sistan-o Baluchestan provinces; south and southeast of Iran (Dlabola 1994; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Tribe Eupelicini

Four species belonging two genera have been recorded from Iran (Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Eupelix cuspidata (Fabricius, 1775)

Material examined: Bagh-e Malek, 2♂, 1♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Bagh-e Malek, 1♂, swept on *Medicago sativa*, 09.ix.2014, leg.: F.P.R.; Dezful, 1♀, swept on gramineae, 12.ix.2014, leg.: F.P.R.; Bagh-e Malek, 1♂, 1♀, swept on grasslands, 15.v.2015, leg.: F.P.R.

Distribution in Iran: Azarbaijan-e Sharghi, Golestan, Kerman, Khuzestan, Lorestan, Mazandaran and Tehran provinces; north, northwest, southwest, south and center of Iran (Dlabola 1981; Mirzayans 1995; Pakarpour *et al.* 2011a; Abdollahi *et al.* 2015; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Paradorydium breviceps* (Melichar, 1902)

Material examined: Dezful, 1♂, 3♀, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.

Distribution in Iran: Isfahan, Kerman, Sistan-o Baluchestan province; center, south and southeast of Iran (Dlabola 1960, 1972, 1981; Khajehali *et al.* 2001; Pakarpour *et al.* 2011c; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

***Paradorydium* (*Paradorydium*) *lanceolatum* (Burmeister, 1839)** (Fig. 2c)**

Material examined: Bagh-e Malek, 1♂, 2♀, swept on gramineae, 09.ix.2014, leg.: F.P.R.

Distribution in Iran: This species is a new record for the Iranian leafhoppers fauna.

Remarks: Male: Length, 7.2 mm.; Base color brownish yellow, macropterous and forewings much longer than abdomen, the head extends forwards and is 3 time longer than wide, The vertex is strongly ridged and narrows gradually towards the apex which is flat, round, dorsally incurved and has a concave dorsal surface; in male the aedeagus has a pentagonal base which is long rounded near the origin of the shaft that stems from the ventral portion of the base. The shaft is swollen at the base and then narrows sharply becoming cylindrical (Fig. 3c).

Paradorydium* (*Paradorydium*) *paradoxum* (Herrich-Schäffer, 1837)

Material examined: Bagh-e Malek, 2♂, 1♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Kerman province; southeast of Iran (Pakarpour *et al.* 2011b; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Tribe Fieberiellini

Five species belonging three genera have been recorded from Iran (Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Fieberiella florii* (Stål, 1864)

Material examined: Dezful, 1♂, 1♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.; Bagh-e Malek, 1♂, 3♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Recorded from Iran without certain locality (Dlabola 1961; 1965; Nast 1972; Modarres Awal 1994; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Fieberiella macchiai* Linnauvori, 1962

Material examined: Dezful, 1♂, 1♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

Distribution in Iran: Qom and Tehran provinces; north and center of Iran (Dlabola 1981, 1984; Rajabi 1991; Modarres Awal 1994; Mirzayans 1995; Abaii 2000; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Tribe Goniagnathini

10 species belonging three genera have been recorded from Iran (Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Goniagnathus* (*Goniagnathus*) *brevis* (Herrich-Schäffer, 1835)

Material examined: Dezful, 1♂, swept on grasslands, 12.ix.2014, leg.: F.P.R.

Distribution in Iran: Azarbaijan-e Sharghi, Kohgiluyeh and Boyerahmad provinces; northwest, west, and center of Iran (Nast 1972; Mirzayans *et al.* 1976; Dlabola 1981; Modarres Awal 1994; Mirzayans 1995;

Abdollahi et al. 2013, 2015; Pakarpour et al. 2015a; Mozaffarian and Wilson 2016).

Goniagnathus (Epistagma) guttulinervis (Kirschbaum, 1868)

Material examined: Ahvaz, 9♂, 8♀, swept on weeds in Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Dezful, 1♂, 1♀, swept on grasslands, 12.ix.2014, leg.: F.P.R.

Distribution in Iran: Hormozgan, Kerman, Khuzestan, Sistan-o Baluchestan and Tehran provinces; northwest, southwest, south, southeast and center of Iran (Dlabola 1972, 1981, 1994; Nast 1972; Mirzayans 1995; Pakarpour et al. 2011a; 2011c; Abdollahi et al. 2015; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Tribe Hecalini

Two species from one genus have been recorded from Iran (Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Hecalus eximius* (Kirschbaum, 1868)

Material examined: Ahvaz, 5♂, 4♀, swept on weeds in palm orchards in Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Ahvaz, 8♂, 2♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Dezful, 1♂, 2♀, swept on gramineae, 12.ix.2014, leg.: F.P.R.

Distribution in Iran: Bushehr, Fars, Hormozgan, Kerman, Khorasan-e Jonubi, Mazandaran, Sistan-o Baluchestan provinces; north, southwest, south, southeast and east of Iran (Dlabola 1972, 1981; Nast 1972; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Pakarpouret al. 2011a; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

***Hecalus glaucescens* (Fieber, 1866)**

Material examined: Bagh-e Malek, 2♂, 1 nymph, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Bagh-e Malek, 1♀, swept on

Medicago sativa, 09.ix.2014, leg.: F.P.R.; Dezful, 3♂, 3♀, swept on gramineae, 12.ix.2014, leg.: F.P.R.; Bagh-e Malek, 2♂, 1♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.; Ahvaz, 1♂, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Sharqi, Bushehr, Fars, Hormozgan, Kerman, Khorasan-e Jonubi, Khuzestan, Sistan-o Baluchestan provinces; northwest, southwest, south, southeast, east and center of Iran (Dlabola 1959, 1960, 1961, 1965, 1972, 1981; Mirzayans et al. 1976; Modarres Awal 1994; Mirzayans 1995; Abdollahi et al. 2015; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Hecalus prasinus* (Matsumura, 1905)*

(Fig. 2d)

Material examined: Ahvaz, 1♂, 5♀, swept on weeds in Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Ahvaz, 3♂, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.

Distribution in Iran: This species is a new record for the Iranian leafhoppers fauna.

Remarks: Male: Length, 7.2 mm.; Base color yellowish brown, vertex parabolic with acute margin at apex, macropterous and forewings longer than abdomen; in male, aedeagus symmetrical, tubular, with one pair of processes at apex (Fig. 3d).

Tribe Macrostelini

24 species from six genera have been recorded from Iran (Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Balclutharhenana* Wagner, 1939

Material examined: Ahvaz, 2♂, 1♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.

Distribution in Iran: North and northwest of Iran (Dlabola 1981; Abdollahi et al. 2015; Mozaffarian and Wilson 2016).

***Balclutharosea* (Scott 1876)**

Material examined: Bagh-e Malek, 4♂, 5♀, swept on grasslands, 15.v.2015, leg.: F.P.R.

Distribution in Iran: Kerman, Khuzestan and Tehran provinces; center, southwest and southeast of Iran (Mirzayans 1995; Pakarpour *et al.* 2011b; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Cicadulina (Cicadulina) mbila* (Naudé, 1924)

Material examined: Ahvaz, 3♂, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Ahvaz, 1♂, 1♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Kerman, Sistan-o Baluchestan provinces; south and southeast of Iran (Dlabola 1960, 1981; Pakarpour *et al.* 2011a; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

***Macrosteles* sp.**

Material examined: Dezful, 1♂, swept on grasslands, 12.ix.2014, leg.: F.P.R.; Ahvaz, 1♂, 1♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

***Macrosteleslaevis* (Ribaut, 1927)**

Material examined: Dezful, 1♂, 2♀, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.

Distribution in Iran: Ardebil, Azarbaijan-e Sharghi, Fars, Guilan, Golestan, Isfahan, Kerman, Khorasan-e Jonubi, Kordestan, Khuzestan and Tehran provinces; north, northwest, west, southwest, south and center of Iran (Kheyri and Alimoradi 1969; Kheyri 1989; Dlabola 1971, 1981; Mirzayans *et al.* 1976; Modarres Awal 1994; Mirzayans 1995; Karimzadeh *et al.* 1998; Haghshenas and Khajehali 2000; Nematollahi and Khajehali 2000; Khajehali *et al.* 2001; Saeidi and Azura Adam 2011; Rezaei *et al.* 2014; Abdollahi *et al.* 2015; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

***Macrosteles ossianilssonii* Lindberg, 1954**

Material examined: Ahvaz, 2♂, 2♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Dezful, 2♂, 1♀, swept on weeds in citrus orchards, 12.ix.2014, leg.: F.P.R.; Dezful, 2♂, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.

Distribution in Iran: Golestan, Khuzestan provinces; north and south of Iran (Dlabola 1971, 1981; Mirzayans 1995; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

***Macrosteles quadripunctulatus* (Kirschbaum, 1868)**

Material examined: Dezful, 1♀, swept on weeds in citrus orchards, 12.ix.2014, leg.: F.P.R.; Dezful, 1♂, 1♀, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.; Ahvaz, 1♂, 1♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Azarbaijan-e Gharbi, Hormozgan, Kerman, Kermanshah, Khuzestan and Tehran provinces; northeast, north, northwest, west, southwest, south and center of Iran (Dlabola 1981; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Moosavi Mahvelati and Modarres Awal 2011; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Tribe Opsiini

31 species belonging nine genera have been recorded from Iran (Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Hishimonus phycitis* (Distant, 1908)

Material examined: Ahvaz, 1♂, 3♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Bagh-e Malek, 3♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Hormozgan, Fars, Kerman, Sistan-o Baluchestan provinces; south of Iran (Salehi *et al.* 2000; Bagheri *et al.* 2008; 2009; Faghihi *et al.* 2011; Zohdi *et al.*

2012; Samavi *et al.* 2012; Mozaffarian and Wilson 2016).

Neoaliturus alboflavovittatus* (Lindberg, 1954)

Material examined: Ahvaz, 1♂, 2♀, swept on weeds in Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Dezful, 3♂, 1♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.

Distribution in Iran: Kerman province; south of Iran (Pakarpour *et al.* 2011a; 2014a; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Neoaliturus pulcher* (Haupt, 1927)

Material examined: Ahvaz, 4♂, 5♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Dezful, 3♂, 7♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

Distribution in Iran: Azarbaijan-e Sharghi, Hormozgan, Kerman, Sistan-o Baluchestan and Tehran provinces; northwest, centre, south and southeast of Iran (Dlabola 1960, 1961, 1972, 1981; Nast 1972; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Abdollahi *et al.* 2015; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Neoaliturus (Neoaliturus) fenestratus* (Herrich-Schäffer, 1834)

Material examined: Ahvaz, 1♂, swept on weeds in palm orchards at Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Dezful, 1♂, 1♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Bagh-e Malek, 2♂, swept on *Medicago sativa*, 09.Ix.2014, leg.: F.P.R.; Dezful, 1♂, swept on grasslands, 12.ix.2014, leg.: F.P.R.; Bagh-e Malek, 1♂, swept on grasslands, 15.v.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Sharghi, Kerman, Khorasan-e Razavi and Tehran provinces; northeast, north, northwest, center and south of Iran (Dlabola 1960; 1972; 1981; Kheyri 1989; Mirzayans *et al.* 1976; Behdad 1992; Modarres Awal 1994; Mirzayans 1995; Moosavi Mahvelati and Modarres Awal 2011; Pakarpour *et al.* 2011c; Saeidi and Azura Adam 2011; Dehghan *et al.* 2012; Pakarpour *et al.* 2014a; Abdollahi *et al.* 2015; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

al. 1976; Behdad 1992; Modarres Awal 1994; Mirzayans 1995; Moosavi Mahvelati and Modarres Awal 2011; Pakarpour *et al.* 2011c; Saeidi and Azura Adam 2011; Dehghan *et al.* 2012; Pakarpour *et al.* 2014a; Abdollahi *et al.* 2015; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Neoaliturus (Neoaliturus) guttulatus* (Kirschbaum, 1868)

Material examined: Dezful, 2♂, 5♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.

Distribution in Iran: Chaharmahal-o Bakhtiari, Isfahan, Kerman and Khorasan-e Razavi provinces; center, northeast and southeast of Iran (Dlabola 1972; Behdad 1992; Modarres Awal 1994; Haghshenas and Khajehali 2000; Khajehali *et al.* 2001; Moosavi Mahvelati and Modarres Awal 2011; Pakarpour *et al.* 2014a; Pakarpour Rayeni *et al.* 2015; Mozaffarian and Wilson 2016).

Neoaliturus (Circulifer) dubiosus* (Matsumura, 1908)

Material examined: Ahvaz, 1♂, 3♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Bagh-e Malek, 3♂, 5♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Isfahan and Tehran provinces; center of Iran (Mirzayans, 1995; Karimzadeh *et al.* 1998; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

***Neoaliturus (Circulifer) haematoceps* (Mulsant and Rey, 1855)**

Material examined: Dezful, 1♂, 6♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Ahvaz, 3♂, 11♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Bagh-e Malek, 5♂, 6♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Gharbi, Fars, Hormozgan, Isfahan, Kerman, Kermanshah, Khorasan-e Razavi, Khuzestan,

Kohgiluyeh and Boyerahmad, Kordestan, Qom, Sistan-o Baluchestan and Tehran provinces; nearly all of Iran (Nast, 1972; Mirzayans *et al.* 1976; Dlabola 1981; Kheyri 1989; Salehi and Izadpanah 1992; Modarres Awal 1994; Mirzayans 1995; Karimzadeh *et al.* 1998; Saeidi and Azura Adam 2011; Omidi *et al.* 2011; Pakarpour *et al.* 2011a; Pakarpour *et al.* 2014a; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

***Neoaliturus (Circulifer) opacipennis* (Lethierry, 1876)**

Material examined: Dezful, 4♂, 10♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.

Distribution in Iran: Azarbaijan-e Sharghi, Chaharmahal-o Bakhtiari, Golestan, Isfahan, Kerman, Khorasan-e Razavi, Khorasan-e Shomali, Khuzestan, Mazandaran, Qazvin, Sistan-o Baluchestan and Zanjan provinces; northwest, north, northeast, center, south and southeast of Iran (Farahbakhsh 1961; Dlabola 1960; 1961; 1965 1972; Kheyri and Alimoradi 1969; Modarres Awal 1994; Karimzadeh *et al.* 1998; Haghshenas and Khajehali 2000; Khajehali *et al.* 2001; Pakarpour *et al.* 2011c; 2014; Khormali 2013; Pakarpour *et al.* 2014a; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

***Neoaliturus (Circulifer) tenellus* (Baker, 1896)**

Material examined: Dezful, 14♂, 22♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Bagh-e Malek, 22♂♀, swept on grasslands, 16.Mar.2014, leg.: F.P.R.; Bagh-e Malek, 30♂♀, swept on grasslands, 09.ix.2014, leg.: F.P.R.; Dezful, 55♂♀, swept on weeds in citrus orchards, 12.ix.2014, leg.: F.P.R.; Ahvaz, 13♂, 18♀, swept on weeds in palm orchards, 26.iv.2015, leg.: F.P.R.; Dezful, 11♂, 30♀, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.; Bagh-e Malek, 8♂, 11♀, swept on grasslands, 15.v.2015, leg.: F.P.R.; Ahvaz, 45♂♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.; Dezful, 5♂, 12♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

F.P.R.; Bagh-e Malek, 24♂♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Fars, Isfahan, Kerman, Khuzestan, Lorestan, Sistan-o Baluchestan and Tehran provinces; center, south, southwest and southeast of Iran (Dlabola 1960; 1964; 1972; 1981; Nast 1972; Mirzayans *et al.* 1976; Kheyri 1989; Behdad 1992; Modarres Awal 1994; Mirzayans 1995; Karimzadeh *et al.* 1998; Khajehali *et al.* 2001; Pakarpour *et al.* 2011a; 2014a; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Opsiush sp.

Material examined: Dezful, 1♂, 3♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

***Orosius albicinctus* Distant, 1918**

Material examined: Ahvaz, 1♂, swept on weeds in palm orchards at Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Ahvaz, 7♂, 2♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Bagh-e Malek, 5♀, swept on grasslands, 15.v.2015, leg.: F.P.R.; Bagh-e Malek, 4♂, 2♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Bushehr, Hormozgan, Isfahan, Kerman and Khuzestan provinces; center, southwest, south and southeast of Iran (Farahbakhsh 1961; Dlabola 1960; 1981; Modarres Awal 1994; Mirzayans 1995; Karimzadeh *et al.* 1998; Shamszadeh 2002; Esmailzadeh-Hosseini *et al.* 2007; Omidi *et al.* 2011; Pakarpour *et al.* 2011a; Nourbakhsh *et al.* 2012; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

***Orosius minuicus* Dlabola, 1979**

Material examined: Ahvaz, 1♂, swept on *Medicago sativa* in palm orchards, 26.iv.2015, leg.: F.P.R.; Ahvaz, 5♂, 9♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Khuzestan province; southwest of Iran (Dlabola 1979; Pakarpour

Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Pseudophlepsius binotatus (Signoret, 1880)

Material examined: Ahvaz, 1♂, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Azarbaijan-e Sharghi, Kerman, Khorasan-e Razavi, Khuzestan, Kohgiluyeh and Boyerahmad, Qom and Tehran provinces; northeast, center, northwest, southwest and south of Iran (Dlabola 1961; 1972; 1981; Mirzayans 1995; Olyaei Torshiz et al. 2011; Abdollahi et al. 2015; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Tribe Paralimnini

30 species belonging to 19 genera have been recorded from Iran (Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Paralimnellas cingulatus (Dlabola, 1960)

Material examined: Dezful, 2♂, 10♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.; Bagh-e Malek, 5♂, 11♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Kerman and Khuzestan provinces; south and southwest of Iran (Dlabola 1960; 1981; Mirzayans 1995; Pakarpour et al. 2011a; 2011c; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Psammotettix alienus (Dahlbom, 1850)

Material examined: Ahvaz, 2♂, 7♀, swept on weeds in Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Bagh-e Malek, 73♂♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Bagh-e Malek, 2♂, 11♀, swept on grasslands, 15.v.2015, leg.: F.P.R.; Ahvaz, 7♂, 10♀, swept on weeds in palm orchards, 26.iv.2015, leg.: F.P.R.; Ahvaz, 30♂♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.; Dezful, 11♂, 6♀, collected with Malaise trap, 27.vii.2015, leg.:

F.P.R.; Bagh-e Malek, 13♂, 21♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Ardebil, Azarbaijan-e Gharbi, Azarbaijan-e Sharghi, Chaharmahal-o Bakhtiari, Guilan, Golestan, Isfahan, Kerman, Kermanshah, Khorasan-e Razavi, Khuzestan, Kohgiluyeh and Boyerahmad, Kordistan, Lorestan, Mazandaran and Tehran, provinces; northeast, north, northwest, southwest, center of Iran (Kheyri 1989; Dlabola 1971; 1981; Modarres Awal 1994; Mirzayans 1995; Karimzadeh et al. 1998; Nematollahi and Khajehali 2000; Haghshenas and Khajehali 2000; Khajehali et al. 2001; Darvish-Mojeni and Taghizadeh 2008; Pakarpour et al. 2011a; 2011c; Zohdi et al. 2012; Rezaei et al. 2014; Abdollahi et al. 2015; Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Psammotettix emarginatus Singh, 1969** (Fig. 2e)

Material examined: Ahvaz, 3♂, 7♀, swept on weeds in palm orchards, 26.iv.2015, leg.: F.P.R.; Dezful, 2♂, 4♀, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.

Distribution in Iran: This species is a new record for the Iranian leafhoppers fauna.

Remarks: Male: Length, 3.4 mm.; Base color light to dirty yellow, dark coloring partly distinct, vertex rectangular and as long as pronotum, fore wings with brownish spots and streaks; in male shaft of aedeagus mostly straight and evenly slender, aperture spoon-shaped, upper margin of aedagus rounded, aedeagus shaft has straight margins in lateral view (Fig. 3e).

Tribe Penthimiini

Two species belonging to one genus have been recorded from Iran (Pakarpour Rayeni et al. 2015a; Mozaffarian and Wilson 2016).

Penthimia nigra (Goeze, 1778)*

Material examined: Bagh-e Malek, 1♂, swept on grasslands, 15.v.2015, leg.: F.P.R.

Distribution in Iran: South and central region of Alborz mountains (Dlabola 1971; Mozaffarian and Wilson 2016).

Penthimia sp.

Material examined: Bagh-e Malek, 1♀, swept on grasslands, 09.ix.2014, leg.: F.P.R.

Tribe Phlepsiini

Seven species belonging to one genus have been recorded from Iran (Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Phlepsius intricatus (Herrich-Schäffer, 1838)*

Material examined: Ahvaz, 2♂, swept on weeds in Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Dezful, 3♂, 4♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Bagh-e Malek, 4♀, swept on grasslands, 09.ix.2014, leg.: F.P.R.; Dezful, 6♂, 4♀, swept on weeds in citrus orchards, 12.ix.2014, leg.: F.P.R.; Ahvaz, 1♂, 3♀, swept on *Medicago sativa* in palm orchards, 26.iv.2015, leg.: F.P.R.; Bagh-e Malek, 3♂, 3♀, swept on grasslands, 15.v.2015, leg.: F.P.R.; Bagh-e Malek, 2♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Azarbaijan-e Gharbi, Azarbaijan-e Sharqi, Chaharmahal-o Bakhtiari, Fars, Isfahan, Kerman, Kermanshah, Kordestan, Qom and Tehran provinces; northeast, north, northwest, west, south-west, south and center of Iran (Nast 1972; Dlabola 1974; 1981; Mirzayans *et al.* 1976; Modarres Awal 1994; Mirzayans 1995; Karimzadeh *et al.* 1998; Haghshenas and Khajehali 2000; Khajehali *et al.* 2001; Pakarpour *et al.* 2011a; 2011c; Moosavi Mahvelati and Modarres Awal 2011; Abdollahi *et al.* 2015; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Phlepsius kambysesi Dlabola, 1984*

Material examined: Bagh-e Malek, 1♂, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Fars and Kohgiluyeh and Boyerahmad provinces; southwest of Iran (Dlabola 1984; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Tribe Scaphoideini

21 species belonging to nine genera have been recorded from Iran (Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Grammacephalus genoicus Dlabola, 1984*

Material examined: Dezful, 1♂, 2♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

Distribution in Iran: Hormozgan province; south and southeast of Iran (Dlabola 1984; Mirzayans 1995; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Grammacephalus pallidus (Linnauvoori, 1978)** (Fig. 2f)

Material examined: Ahvaz, 1♂, 1♀ collected with sticky yellow trap in Shahid Chamran University of Ahvaz, 26.iv.2015, leg.: F.P.R.

Distribution in Iran: This species is a new record for the Iranian leafhoppers fauna.

Remarks: Male: Length, 6.1 mm.; Base color light to dirty yellow, head strongly produced, vertex brown with a stripe along midline, vertex markings are similar to *G. genoicus* and *G. minabicus*, face creamy white with narrow brown band along anterior margin; in male, aedeagus simple, shaft slender, curved dorsally and rounded apically (Fig. 3f).

Grammacephalus pugio (Noualhier, 1895)*

Material examined: Dezful, 4♂, 3♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Ahvaz, 3♂, 2♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.; Dezful, 1♂, 3♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

Distribution in Iran: Hormozgan and Kerman provinces; south and southeast of Iran (Dlabola 1960; 1965; 1977; 1981; Nast 1972; Mirzayans 1995; Pakarpour *et al.* 2011a; 2011c; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Neolimnus (Neolimnus) egyptiacus (Matsumura, 1908)*

Material examined: Dezful, 1♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Ahvaz, 5♂, 3♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Dezful, 2♂, swept on gramineae, 12.ix.2014, leg.: F.P.R.; Ahvaz, 4♂, 5♀, swept on *Medicago sativa* in palm orchards, 26.iv.2015, leg.: F.P.R.; Ahvaz, 3♂, 6♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Hormozgan, Kerman and Sistan-o Baluchestan provinces; south and southeast of Iran (Dlabola 1960; 1965; 1981; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Tribe Stenometopiini

Six species belonging two genera have been recorded from Iran (Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Doratulina lahorensis (Distant, 1918)*

Material examined: Ahvaz, 4♂, 4♀, swept on weeds in palm orchards, 13.iii.2014, leg.: F.P.R.; Dezful, 5♂, 7♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Bagh-e Malek, 1♂, 3♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Ahvaz, 6♀, swept on weeds in palm orchards, 26.iv.2015, leg.: F.P.R.; Ahvaz, 6♂, 13♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Sistan-o Baluchestan province; southeast of Iran (Pakarpour and Nozari 2011; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Doratulina (Doratulina) pallida Dlabola, 1994*

Material examined: Bagh-e Malek, 1♂, 2♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Dezful, 5♂, 1♀, swept on gramineae, 12.ix.2014, leg.: F.P.R.; Ahvaz, 1♂, 1♀, swept on weeds in palm orchards, 26.iv.2015, leg.: F.P.R.; Ahvaz, 1♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Hormozgan, Kerman and Sistan-o Baluchestan provinces; south of Iran (Dlabola 1994; Pakarpour *et al.* 2011c; Pakarpour Rayeni *et al.* 2015a; Mozaffarian and Wilson 2016).

Subfamily Lassinae

Four species belonging two genera have been recorded from Iran (Mozaffarian and Wilson 2016).

Batracomorphus irrortatus Lewis, 1834

Material examined: Ahvaz, 1♂, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Sharghi, Bushehr, Hormozgan, Khorasan-e Razavi, Khuzestan, Kohgiluyeh and Boyer-ahmad, Kordestan, Lorestan and Tehran provinces; northwest, west, southwest, south, southeast, east and center of Iran (Dlabola 1960; 1963; 1972; Mirzayans *et al.* 1976; Modarres Awal 1994; Moosavi Mahvelati and Modarres Awal 2011; Abdollahi *et al.* 2015; Mozaffarian and Wilson 2016).

Subfamily Idiocerinae

24 species belonging to 10 genera have been recorded from Iran (Mozaffarian and Wilson 2016).

Viridicerus sp.

Material examined: Dezful, 8♂, 16♀, swept on grasslands, 12.ix.2014, leg.: F.P.R.

Remark: These specimens have deep differences with other species of the genus *Viridicerus*, maybe a new species.

Subfamily Macropsinae

Nine species belonging to four genera have been recorded from Iran (Mozaffarian and Wilson 2016).

Macropsis graminea (Fabricius, 1798)*

Material examined: Dezful, 1♂, 1♀, swept on gramineae, 12.ix.2014, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Gharbi, Azarbaijan-e Sharghi, Hormozgan, Kerman, Kordestan, and Tehran provinces; northwest, west and center of Iran (Dlabola 1974; 1981; Mirzayans 1995; Mozaffarian and Wilson 2016).

Subfamily Megophthalminae

27 species belonging 10 genera have been recorded from Iran (Mozaffarian and Wilson 2016).

Tribe Agalliini

15 species belonging to four genera have been recorded from Iran (Mozaffarian and Wilson 2016).

Agallia harrarensis Melichar, 1911

Material examined: Ahvaz, 5♂, 8♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Ahvaz, 1♂, 3♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.; Bagh-e Malek, 3♂, 8♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Fars, Hormozgan, Kerman, Khuzestan and Tehran provinces; south of central region of Alborz Mountains, southwest, south and southeast of Iran (Dlabola 1960; 1972; 1981; Kheyri 1989; Nast 1972; Modarres Awal 1994; Mirzayans 1995; Mozaffarian and Wilson 2016).

Anaceratagallia aciculata (Horváth, 1894)*

Material examined: Ahvaz, 4♂, 4♀, swept on weeds in palm orchards, 26.iv.2015, leg.: F.P.R.; Ahvaz, 3♂, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Fars, Kerman and Tehran provinces; north, southwest and southeast (Dlabola 1981; Mirzayans 1995; Babaei and Izadpanah 2003; Pakarpour *et al.* 2011b; Mozaffarian and Wilson 2016).

Anaceratagallia laevis (Ribaut, 1935)*

Material examined: Bagh-e Malek, 2♂, 10♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Bagh-e-Malek, 3♂, 8♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Sharghi, Chaharmahal-o Bakhtiari, Guilan, Isfahan, Kordestan, Lorestan and Tehran provinces; north, northwest, west and center of Iran (Dlabola 1981; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Karimzadeh *et al.* 1998; Haghshenas and Khajehali 2000; Khajehali *et al.* 2001; Bagheri and Nasr Isfahani 2008; Abdollahi *et al.* 2015; Mozaffarian and Wilson 2016).

Anaceratagallia venosa (Fourcroy, 1785)*

Material examined: Ahvaz, 4♀, swept on weeds in palm orchards at Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Dezful, 1♂, 3♀, swept on weeds in citrus orchards, 14.iii.2014, leg.: F.P.R.; Ahvaz, 2♂, 5♀, swept on *Medicago sativa* in palm orchards, 26.iv.2015, leg.: F.P.R.

Distribution in Iran: North of Iran (Nast 1972; Dlabola 1981; Mozaffarian and Wilson 2016).

Austroagallia sinuata (Mulsant and Rey, 1855)

Material examined: Ahvaz, 3♂, 11♀, swept on weeds in palm orchards at Shahid Chamran University of Ahvaz, 13.iii.2014, leg.: F.P.R.; Bagh-e Malek, 1♂, 5♀, swept on grasslands, 16.iii.2014, leg.: F.P.R.; Ahvaz, 1♂, 3♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.; Bagh-e Malek, 1♂, 6♀, collected with Malaise trap, 28.Vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Sharghi, Bushehr, Chaharmahal-o Bakhtiari, Fars, Golestan, Hormozgan, Isfahan,

Kerman, Khorasan-e Jonubi, Khuzestan, Lorestan, and Tehran provinces; north, northwest, southwest, south, center and northeast of Iran (Farahbakhsh 1961; Dlabola 1963; 1964; 1971; 1972; 1981; Mirzayans et al. 1976; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Karimzadeh et al. 1998; Haghshenas and Khajehali 2000; Khajehali et al. 2001; Yarmand et al. 2006; Bagheri and Nasr Isfahani 2008; Pakarpour et al. 2011a; 2011c; Khormali 2013; Abdollahi et al. 2015; Mozaffarian and Wilson 2016).

Subfamily Typhlocybinae

Typhlocybinae is the second largest subfamily of Cicadellidae comprising 84 valid species in Iran (Mozaffarian and Wilson 2016).

Tribe Empoascini

16 species belonging seven genera have been recorded from Iran (Mozaffarian and Wilson 2016).

Asymmetrasca decedens (Paoli, 1932)*

Material examined: Ahvaz, 12♂, 18♀, swept on *Medicago sativa* in palm orchards, 07.ix.2014, leg.: F.P.R.; Ahvaz, 7♂, 15♀, swept on *Medicago sativa*, 26.iv.2015, leg.: F.P.R.; Bagh-e Malek, 5♂, 11♀, swept on *Medicago sativa*, 09.ix.2014, leg.: F.P.R.; Dezful, 15♂, 17♀, swept on grasslands, 12.ix.2014, leg.: F.P.R.; Ahvaz, 45♂♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.; Dezful, 55♂♀, collected with aspirator from *Vitis venifera*, 27.vii.2015, leg.: F.P.R.; Bagh-e-Malek, 50♂♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Chaharmahal-o Bakhtiari, Fars, Golestan, Kerman, Khorasan-e Razavi provinces; northeast, north, center and southwest of Iran (Haghhighian and Sadeghi 2001; Taghizadeh and Salehi 2002; Darvish-Mojeni and Taghizadeh 2008; Haghhighian et al. 2003; 2008; Olyaei Torshiz et al. 2001; Tahrir adabi et al. 2013; Mozaffarian and Wilson 2016).

Empoasca (Empoasca) decipiens Paoli, 1930

Material examined: This species has a widespread distribution in Khuzestan province and has been collected from all the sampling areas with high population.

Distribution in Iran: Alborz, Azarbaijan-e Gharbi, Azarbaijan-e Sharghi, Chaharmahal-o Bakhtiari, Fars, Golestan, Isfahan, Kerman, Kordestan, Khorasan-e Razavi, Khuzestan, Mazandaran, Semnan, Sistan-o Baluchestan and Tehran provinces; nearly all Iran (Nast 1972; Mirzayans 1976; Dlabola 1981; Kheyri 1989; Behdad 1992; Modarres Awal 1994; Mirzayans 1995; Karimzadeh et al. 1998; Noori and Javan Moghaddam 1998; Nematollahi and Khajehali 2000; Haghshenas and Khajehali 2000; Khajehali et al. 2001; Naseri et al. 2007a; 2007b; Kianpour et al. 2008; 2009; Izadpanah et al. 2008; Darvish-Mojeni and Taghizadeh 2008; Karimzadeh and Dworakowska 2011; Pakarpour et al. 2011a; 2011c; Sirjani 2012; Nourbakhsh et al. 2012; Rezaei et al. 2014; Mozaffarian and Wilson 2016).

Empoasca (Empoasca) punjabensis Singh-Pruthi, 1940

Material examined: Bagh-e Malek, 5♂, 3♀, swept on grasslands, 15.v.2015, leg.: F.P.R.; Ahvaz, 1♂, 2♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.; Bagh-e Malek, 7♂, 11♀, collected with Malaise trap, 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Gharbi, Azarbaijan-e Sharghi, Fars, Golestan, Isfahan, Kerman, Kermanshah, Khorasan-e Razavi, Khuzestan, Mazandaran, Qazvin, Tehran and Zanjan provinces; nearly all Iran (Zachvatkin 1945; Dlabola 1960; 1971; 1972; 1981; Kheyri and Alimoradi 1969; Kheyri 1989; Modarres Awal 1994; Mirzayans 1995; Karimzade and Dworakowska 2011; Moosavi Mahvelati and Modarres Awal 2011; Khormali 2013; Abdollahi et al. 2015; Mozaffarian and Wilson 2016).

Tribe Erythroneurini

33 species belonging to 14 genera have been recorded from Iran (Mozaffarian and Wilson 2016).

Arboridia (Arboridia) Kermanshah* (Dlabola, 1963)

Material examined: Dezful, 29♂♀, collected with aspirator from *Vitis venifera*, 27.vii.2015, leg.: F.P.R.

Distribution in Iran: Isfahan, Khorasan-e Razavi and Tehran provinces; northeast, center and west of Iran (Dlabola 1963; 1981; Nast 1972; Mirzayans 1995; Hesami *et al.* 2002; Latifian *et al.* 2008; Olyaei Torshiz *et al.* 2011; Mozaffarian and Wilson 2016).

Helionidia biplagiata* (Haupt, 1927)

Material examined: Ahvaz, 3♂, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Fars, Kerman and Tehran provinces; south of central region of Alborz Mountains, southwest and south of Iran (Dlabola 1960; 1981; Mirzayans 1995; Mozaffarian and Wilson 2016).

***Helionidia punctulata* (Ramakrishnan & Menon, 1974)** (Fig. 2g)**

Material examined: Bagh-e Malek, 6♂, 2♀, swept on grasslands, 09.ix.2014, leg.: F.P.R.

Distribution in Iran: This species is a new record for the Iranian leafhoppers fauna.

Remarks: Male: Length, 2.6 mm.; Base color yellowish green, vertex yellow with a pair of dark spots, pronotum greenish in anterior pale with 6 distinct dots, macropterous and forewings much longer than abdomen; in male, aedeagus distinct, apically branched with two pairs of short appendages (Fig. 3g).

***Negoneura* sp.**

Material examined: Dezful, 1♀, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.

Distribution in Iran: Hormozgan province; south of Iran (Dlabola 1994; Mozaffarian and Wilson 2016).

Remarks: Dlabola (1994) described this genus as *Genoneura* in 1994 but McKamey (2006) changed this genus name to *Negoneura* comb.nov.

***Tamaricella* sp.**

Material examined: Bagh-e Malek, 2♀, swept on grasslands, 09.ix.2014, leg.: F.P.R.

Zygina (Zygina) ordinaria* (Ribaut, 1936)

Material examined: Dezful, 2♂, 5♀, swept on grasslands and weeds, 27.iv.2015, leg.: F.P.R.

Distribution in Iran: North of Iran (Mozaffarian and Wilson 2016).

Tribe Typhlocybini

27 species belonging to nine genera have been recorded from Iran (Mozaffarian and Wilson 2016).

***Edwardsiana rosae* (Linnaeus, 1758)**

Material examined: Bagh-e Malek, 45♂♀, collected with aspirator from *Rosa* sp. 28.vii.2015, leg.: F.P.R.

Distribution in Iran: Alborz, Azarbaijan-e Sharghi, Fars, Kerman and Tehran provinces; northeast, north, northwest, center, southwest and south (Dlabola 1971; 1981; Esmaili 1984; Rajabi 1991; Behdad 1991; 1992; Mirzayans 1995; Abaii 2000; Pakarpour *et al.* 2011a; 2011c; Nourbakhsh *et al.* 2012; Abdollahi *et al.* 2015; Mozaffarian and Wilson 2016).

Eupteryx (Eupteryx) pavlovskii* Zachvatkin, 1947

Material examined: Ahvaz, 2♂, 3♀, swept on weeds in palm orchards, 07.ix.2014, leg.: F.P.R.; Ahvaz, 1♀, collected with Malaise trap, 25.vii.2015, leg.: F.P.R.

Distribution in Iran: Fars, Kohgiluyeh and Boyerahmad, Mazandaran and Tehran

provinces; north, northwest and southwest (Nast 1972; Dlabola 1974; 1981; Mirzayans 1995; Mozaffarian and Wilson 2016).

Remarks: Dlabola (1974) described a new species as *Eupteryx tochalica*; then he (1981) and Mirzayans (1995) reported *E. tochalica* from different parts of Iran but now *E. tochalica* is a synonym of *Eupteryx* (*Eupteryx*) *pavlovskii*.

Eupteryx (Eupteryx) kaghanensis Ahmed, 1969 ** (Fig. 2h)

Material examined: Dezful, 5♂, 8♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

Distribution in Iran: This species is a new record for the Iranian leafhoppers fauna.

Remarks: Male: Length, 3.2 mm.; Base color yellowish green, fore margin of vertex with two spots, hind margin with one variable

spot, pronotom and scutellum with numerous spots, fore wings with yellowish green and partly with brown pattern; in male shaft of aedeagus more or less straight in lateral view, apex with two pairs of recurrent appendages (Fig. 3h).

Zyginaella pulchra Löw, 1885*

Material examined: Bagh-e Malek, 1♂, 2♀, swept on grasslands, 09.ix.2014, leg.: F.P.R.; Dezful, 2♂, 2♀, collected with Malaise trap, 27.vii.2015, leg.: F.P.R.

Distribution in Iran: Fars, Khorasan-e Razavi, Markazi, Qazvin, Semnan and Tehran provinces; northeast, center and southwest of Iran (Rajabi and Mirzayans 1989; Rajabi 1991; Mirzayans 1995; Taghizadeh and Salehi 2002; Moosavi Mahvelati and Modarres Awal 2011; Mozaffarian and Wilson 2016).



Figure 1. Map of the study area; sampling sites consist of Ahvaz, Bagh-e Malek and Dezful in Khuzestan province (southwest of Iran)



Figure 2. Habitus figures of new records to Iranian fauna (scale bar = 1mm), **a.** *Euscelidius schenckii* (Kirschbaum); **b.** *Hardya (Hardya) melanopsis* (Hardy); **c.** *Paradorydium (Paradorydium) lanceolatum* (Burmeister); **d.** *Hecalus prasinus* (Matsumura); **e.** *Psammotettix emarginatus* Singh; **f.** *Grammacephalus pallidus* (Linnauvori); **g.** *Helionidia punctulata* (Ramakrishnan & Menon); **h.** *Eupteryx (Eupteryx) kaghanensis* Ahmed.




Figure 3. Aedeagus in lateral and dorsal view, **a.** *Euscelidius schenckii* (Kirschbaum); **b.** *Hardya (Hardya) melanopsis* (Hardy); **c.** *Paradorydium (Paradorydium) lanceolatum* (Burmeister); **d.** *Hecalus prasinus* (Matsumura); **e.** *Psammotettix emarginatus* Singh; **f.** *Grammacephalus pallidus* (Linnavuori); **g.** *Helionidia punctulata* (Ramakrishnan & Menon); **h.** *Eupteryx (Eupteryx) kaghanensis* Ahmed.

Discussion

Insect fauna of Iran are affected by elements from Oriental, Africa, Mediterranean area and Palaearctic, so remarkable difficulties have risen in studying any single group (Minaei 2013). However, many species of the family Cicadellidae have economic importance but the knowledge of the leafhoppers fauna and their distribution in Iran is far from complete and extensive field work is needed for a reasonable evaluation of the leafhoppers biodiversity in this region. Pakarpour Rayeni (2011) investigated leafhoppers fauna in Kerman province and reported 57 species from 42 genera in six subfamilies.

A faunistic study on the leafhoppers of northwestern Iran was carried out by Abdollahi *et al.* (2015) and they reported 69 species which collected by themselves or other specimens deposited in Hayk Mirzayans Museum. In the course of this study 90 species belonging to 49 genera and seven subfamilies were found and identified on the fauna of Khuzestan province, of which 33 species are those also collected by Pakarpour Rayeni (2011) in Kerman province and 20 species are those collected with Abdollahi *et al.* (2015) in Ardebil, Azarbaijan-e Gharbi and Azarbaijan-e Sharghi provinces. The results revealed are more similar between Khuzestan and Kerman provinces probably due to similar climatic conditions (especially with south of Kerman province).

In addition 47 species are reported from Khuzestan province for the first time and eight species are newly recorded for Iranian leafhoppers fauna. The result of this study can be helpful for completing the checklist of leafhoppers in Iran and the data will help other researcher in ongoing faunistic works. Finally it is expected that a large number of leafhoppers species remain to be found out in different parts of this region and other

parts of Iran, so future studies are highly recommended.

Acknowledgments

We express our sincere thanks to Dr. Michael R. Wilson (National Museum of Wales, Cardiff, UK) for his help in identification. The authors also warmly thanks to three reviewer for revising the manuscript and their valuable comments. This study is supported by Shahid Chamran University of Ahvaz.

References

- Abaii, M. 2000. *Pests of forest trees and shrubs of Iran*. Publication of Agricultural research, Ministry of Agriculture. Agricultural Research, Education and Extension Organization, Tehran, Iran, 178 pp.
- Abdollahi, T., Mozaffarian, F. and Jalali Zand, A.R. 2013. Key to the subfamilies of Cicadellidae (Hem: Auchenorrhyncha) in East Azarbaijan province, Iran and two new records for the region. *Applied Plant Protection*, 2 (2): 147–155.
- Abdollahi, T., Jalali Zand, A.R. and Mozaffarian, F. 2014a. First report of *Psammotettix seriphidii* Emeljanov, 1962 from Iran. Proceedings of 21th congress of Irnian Plant Protection 23–26 August 2014, University of Urmia, Vol. I. Pests, p. 489.
- Abdollahi, T., Jalali Zand, A.R. and Mozaffarian, F. 2014b. A faunistic study of the leafhoppers (Hemiptera: Auchenorrhyncha: Cicadellidae) of Ardabil province, Iran. Proceedings of 21th congress of Irnian Plant Protection 23–26 August 2014, University of Urmia, Vol. I. Pests, p. 570.
- Abdollahi, T., Jalali Zand, A.R. and Mozaffarian, F. 2014c. The first report of *Macrosteles chobauti* (Hem., Cicadellidae) from Iran. *Journal of Entomological Society of Iran*, 34 (2): 27–28.
- Abdollahi, T., Jalali Zand, A.R., Mozaffarian, F. and Wilson, M. 2015. A faunistic study on the leafhoppers of Northwestern Iran (Hemiptera, Cicadellidae). *Zookeys*, 496, 27–51.

- Anufriev, G.A. and Emeljanov, A.F. 1988. *Suborder Cicadinea (Auchenorrhyncha)- cicadas*. In: Lera, P. A. (Ed.), *Key to the Identification of Insects of the Far East USSR*. Vol. 2 Homoptera and Heteroptera. Leningrad, Nauka Publishing House, 495 pp.
- Babaie, Gh. and Izadpanah, K. 2003. Vector Transmission of Eggplant Mottled Dwarf Virus in Iran. *Journal of Phytopathology*, 151: 679-682.
- Bagheri, A., Faghihi, M.M. and Salehi, M. 2008. Study on the population fluctuations of *Hishimonus phycitis* hosting phytoplasma of witches' broom disease of lime trees in Hormozgan province. Proceedings of 18th congress of Irnian Plant Protection 24-27 August 2008, Bu-ali sina University of Hamedan, pp. 61-66.
- Bagheri, M.R. and Nasr Isfahani, M. 2008. The fauna of harmful and beneficial arthropods of medicinal and range plants in Isfahan. *Journal of Entomological Research*, 3 (2): 119-132.
- Bagheri, A.N., Salehi, M., Faghihi, M.M., Samavi, S. and Sadeghi, A. 2009. Transmission of *Candidatus Phytoplasma aurantifolia* to Mexican lime by the leafhopper *Hishimonus phycitis* in Iran. *Journal of Plant Pathology*, 91(4): 105.
- Behdad, E. 1991. *Pests of fruit crops in Iran*. Markaz-e Nashr-e Bahman, Iran, 822 pp.
- Behdad, E. 1992. *Pests of field crops in Iran*. Markaz-e Nashr-e Yadboud, Iran, 618 pp.
- Biedermann R. and Niedringhaus R. 2009. The plant and Leafhoppers of Germany, Identification key to all species. Wilhelm Brüggemann GmbH, Bremen, 409p.
- Backus, E.A. 1988. Sensory systems and behaviours which mediate hemipteran plant-feeding: a taxonomic overview. *Journal of Insect Physiology*. 34: 151-165.
- Backus, E.A., Serrano, M.S. and Ranger, C. M. 2005. Mechanisms of hopperburn: an overview of insect taxonomy, behavior, and physiology. *Annual Review of Entomology*, 50: 125-151.
- Dakhil, H.A., Abou-Fakhr, H.E., El-Mohtar, C. and Abou-Jawdah, Y. 2011. Survey of leafhopper species in almond orchards infected with almond witches'-broom phytoplasma in Lebanon. *Journal of Insect Science*, 11:60, available online: insectscience.org/11.60.
- Darvish-Mojeni, T. and Taghizadeh, M. 2008. Identification of leafhoppers species and study on biology of dominant species in the cotton fields of Golestan province. Proceedings of 18th congress of Irnian Plant Protection 24-27 August 2008, Bu-ali sina University of Hamedan, Vol. I. Pests, p. 475.
- Davoodi, Z. 1980. Some morphological and bio-ecological studies on *Edwardsiana rosae* L. *Applied Entomology and Plant Pathology*, 48 (1): 53-65
- Dehghan, S., Salehi, M., Khanchezar, A., Rastegari, N. and Salari, M. 2012. Transmission characteristics of lettuce phyllody phytoplasma by *Neoaliturus fenestratus* In Fars, Iran. *Journal of Plant Pathology*, 48 (1): 35-36.
- Della Giustina, W. 1989. *Homoptères Cicadellidae. III. Compléments aux ouvrages d'Henri Ribaut. Faune de France*. Paris, Lechevalier, 73: 350 pp.
- Dietrich, C.H. 2005. Keys to the families of Cicadomorpha and subfamilies and tribes of Cicadellidae (Hemiptera, Auchenorrhyncha). *Florida Entomologist*, 88 (4): 502-517.
- Dietrich, C.H. 2008. *Leafhoppers (Hemiptera: Cicadellidae)*. In: Capinera, J. (Eds.), Encyclopedia of entomology, Second edition, vol. 3. Springer-Verlag, Heidelberg, 2160-2162 pp.
- Dlabola, J. 1957. Die Zikaden Afghanistans nach den Ergebnissen der von Herrn Klapperich in den Jahren 1952-1953 nach Afghanistan untemommenen Expedition. *Mitteilungen der Miinchner Entomologischen Gesellschaft*, 1957: 265-303.
- Dlabola, J. 1960. Iranische Zikaden (Homoptera: Auchenorrhyncha). (Ergebnisse der entomologischen Reisen Willy Richter, Stuttgart, in Iran, 1954 und 1956, N: 31). *Stuttgarter Beiträge zur Naturkunde*, 41: 1-24.
- Dlabola, J. 1963. Typen und wenig bekannte Arten aus der Sammlung H. Haupt mit Beschreibungen einiger Zikadenarten aus Sibirien. *Acta Entomologica Musei Nationalis Pragae*, 35: 313-331.
- Dlabola, J. 1964. Die Zikaden Afghanistans (Homoptera, Auchenorrhyncha), II.

- Ergebnisse der Sammelreisen von Dr. H. G. Amsel, G. Ebert, Dr. Erichson, J. Klapperich und Dr. K. Lindberg. *Mitteilungen der Munchner Entomologischen Gesellschaft*, 54: 237–255.
- Dlabola, J. 1965. Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei. 54; Homoptera, Auchenorrhyncha. *Acta Faunistica Entomologica Musei Nationalis Pragae*, 11: 79–136.
- Dlabola, J. 1971. Taxonomische und chorologische Ergänzungen der Zikadenfauna von Anatolien, Iran, Afghanistan und Pakistan (Homoptera: Auchenorrhyncha). *Acta entomologica Bohemoslovaca*, 68: 377–396.
- Dlabola, J. 1972. Beiträge zur Kenntnis der Fauna Afghanistans. Homoptera Auchenorrhyncha. *Acta Musei Moraviae Scientiae Biologicae*, 16–17: 189–248.
- Dlabola, J. 1974. Ergebnisse der Tschechoslowakisch-Iranischen Entomologischen Expedition nach dem Iran, 1970, No. 3 (Homoptera: Auchenorrhyncha) (1 Teil). *Acta Entomologica Musei Nationalis Pragae*, Supplement 6: 29–73.
- Dlabola, J. 1977. Neue iranische Cicadelliden und Arten mit faunistischen Erstfunden (Homoptera: Auchenorrhyncha). *Acta Entomologica Bohemoslovaca*, 74 (4): 242–262.
- Dlabola, J. 1979. Neue Zikaden aus Anatolien, Iran und aus sudeuropäischen Ländern (Homoptera: Auchenorrhyncha). *Acta Zoologica Academiae Scientiarum Hungaricae*, 21 (3–4): 235–257.
- Dlabola, J. 1980. Drei neue Diplocolenus Arten und taxonomisch zoogeographische Übersicht der Gattung in der Palaarktis. *Acta Faunistica Entomologica Musei Nationalis Pragae*, 16: 73–82.
- Dlabola, J. 1981. Ergebnisse der tschechoslowakisch-iranischen entomologischen Expeditionen nach dem Iran (Mit Angaben über einige Sammelresultate in Anatolien) (1970 und 1973) (Homoptera, Auchenorrhyncha). II Teil. *Acta Musei Nationalis Pragae*, 40: 127–311.
- Dlabola, J. 1984. Neue Zikadenarten von dem Mediterraneum-Raum und Iran mit weiteren Beiträgen zur iranischen Fauna (Hom, Auchenorrhyncha). *Acta Musei Nationalis Pragae*, 40: 21–63.
- Dlabola, J. 1987. Neue ostmediterrane und iranische Zikadenarten. *Acta Entomologica Bohemoslovaca*, 84: 295–312.
- Dlabola, J. 1994. Ergänzungen zur iranischen, israelischen und benachbarten Zikadenfaunen mit Beschreibungen 30 neuer Taxone (Homoptera, Auchenorrhyncha). *Acta Musei Nationalis Pragae*, 49 (1–4): 41–110.
- Dlabola, J. and Heller, F. 1962. Iranische Zikaden II, (Ergebnisse der entomologischen Reisen Willi Richter, Stuttgart, im Iran 1954 und 1956, Nr. 42). *Stuttgarter Beiträge zur Naturkunde*, 90: 1–8.
- Dlabola, J. and Jankovic, L. 1981. Drei neue Erythria-Arten und einige Ergänzungen der jugoslawischen Zikadenfauna. *Bulletin de l'Academie Serbe des Sciences et des Arts Classe des Sciences Mathématiques et Naturelles Sciences Naturelles*, 75 (21): 67–79.
- Dmitriev, D.A. (2003-present) On ward. 3I Interactive Keys and Taxonomic Databases: Typhlocybinae. Available from: <http://dmitriev.speciesfile.org/> (accessed 20 June 2016).
- Esmaili, M. (1984) *Important Pests of Fruit Trees*. Markaz-e Nashr-e Sepehr publication, Tehran, 584 pp.
- Emeljanov A.F. 1999. A key to genera of the subfamily Deltoccephalinae s.l. (Homoptera, Cicadellidae) from Kazakhstan, Middle Asia, and Mongolia, with description of new genera and subgenera. *Entomological Review*, 79: 547–562.
- Esmailzadeh-Hosseini, S.A., Mirzaie, A., Jafari-Nodooshan, A. and Rahimian, H. 2007. The first report of transmission of a phytoplasma associated with sesame phyllody by *Orosiusa lobicinctus* in Iran. *Australasian Plant Disease Notes*, 2: 33–34.
- Faghihi, M.M., Bagheri, A.N., Bahrami, H.R., Hasanzadeh, H. and Rezazadeh, R. 2011. Witches'-Broom disease of lime affects seed germination and seedling growth but is not seed transmissible. *Plant Disease*, 95 (4): 419–422.
- Farahbakhsh, Gh. 1961. A checklist of economically important insects and other

- enemies of plants and agricultural products in Iran. *Plant organization Press*, Iran, 153pp.
- Haghigian, F., Sadeghi, S.A. and Talebi, M. 2003. *Empoasca decedens* (Hom: Cicadellidae) as a new pest of poplar and its abundance on colons in Chaharmahal and Bakhtiari province. *Iranian Journal of Forest and Protection Research*, 1 (1): 73–82.
- Haghigian, F., Sadeghi, S.A. and Talebi, M. 2008. Population fluctuation and natural enemies identify of *Empoasca decedens* (Homoptera: Cicadellidae) and control management of this pest on native poplar in Chaharmahal and Bakhtiari province. Proceedings of 18th congress of Irnian Plant Protection, 24–27 August 2008, Bu-ali sina University of Hamedan, Vol. I. Pests, p. 307.
- Haghshenas, A. and Khajehali, J. 2000. Leafhoppers (Homoptera: Auchenorrhyncha) fauna of potato fields in Chagar Mahal and Bakhtiari province. Proceedings of 13th congress of Irnian Plant Protection, 2–7 September 1995, Isfahan University of Technology, Vol. I. Pests, p. 249.
- Hesami, Sh., Seideslami, H. and Hatami, B. 2002. Impact of *Anagrus atomus* (Hym.: Mymaridae) overwintering refuge on its abundance and parasitism of grape leafhopper *Arboridia kermanshah* (Hom: Cicadellidae) eggs in Isfahan. Proceedings of 14th congress of Irnian Plant Protection, 7–11 September 2002, Razi University of Kermanshah, Vol. I. Pests, p. 113.
- Karimzadeh, J. and Dworakowska, I. 2011. Differentiation of two closely related species of the genus *Empoasca* (Hem.: Cicadellidae). *Journal of Entomological Society of Iran*, 30 (2): 89–91.
- Karimzadeh, J., Kharrazi Pakdel, A. and Kheyri, M. 1998. Leafhopper (Homoptera: Auchenorrhyncha) fauna of sugar beet fields in Isfahan province. Proceedings of 12th congress of Irnian Plant Protection, 23–27 August 1998, University of Tehran, Karaj, Vol. I. Pests, p. 69.
- Khajehali, J., Seyedoleslami, H. and Kamali, K. 2001. A contribution to study of hoppers (Hom.: Auchenorrhyncha) of potato fields in Isfahan and Daran. *Applied Entomology and Phytopathology*, 68 (1&2): 43–71.
- Kheyri, M. 1989. An inventory of pests attacking sugar-beet in Iran. *Applied Entomology and Phytopathology*, 56 (1&2): 75–91.
- Kheyri, M. and Alimoradi, I. 1969. *The leafhoppers of Sugarbeet in Iran and their role in curly-top virus disease*. Sugarbeet Seed Institute, Karaj, 54 pp.
- Khormali, S. 2013. Fauna study of Cicadellidae (Leaf hoppers Family) on cotton fields of Golestan province. *Iranian Journal of Cotton Researches*, 1 (1): 93–100.
- Kianpour, R., Fathipour, Y. and Kamali, K. 2008. Population fluctuation and spatial distribution patterns of *Bemisia tabaci* and *Bemisia argentifolii* (Homoptera: Aleyrodidae) and *Empoasca decipiens* (Homoptera: Cicadellidae) on eggplants in Varamin. Proceedings of 18th congress of Irnian Plant Protection, 24–27 August 2008, Bu-ali sina University of Hamedan, Vol. I. Pests, p. 339.
- Kianpour, R., Fathipour, Y. and Kamali, K. 2009. Population fluctuation and spatial distribution patterns of *Bemisia tabaci* and *Bemisia argentifolii* and *Empoasca decipiens* on eggplants in Varamin. *Applied Entomology and Phytopathology*, 77 (2): 71–94.
- Latifian, M., Seyedoleslami, H. and Ghajeali, G. 2008. A comparison on several sampling techniques to estimating relative population density of grape leafhopper (*Arboridia kermanshah*) in adult and nymph stages. Proceedings of 18th congress of Irnian Plant Protection, 24–27 August 2008, Bu-ali sina University of Hamedan, Vol. I. Pests, p. 335.
- Le Quesne, W.J. 1965. Hemiptera Cicadomorpha (excluding Deltocephalinae and Typhlocybinae). *Handbooks for the Identification of British insects II* (2a): 1–64.
- Le Quesne W.J., 1969. Hemiptera Cicadomorpha Deltocephalinae. *Handbooks for the Identification of British Insects II* (2b): 65–148.
- Le Quesne W.J., Payne K. R. 1981. Cicadellidae Typhlocybinae with a check list of the British Auchenorrhyncha (Hemiptera, Homoptera). *Handbooks for the Identification of British Insects II* (2c): 1–95.
- Maramorosch, K., and Harris, K.F. 1979. *Leafhopper Vectors and Plant Disease Agents*. Academic Press, New York. 368 pp.

- McKamey, S.H. 2006. Further new genus-group names in Cicadellidae (Homoptera). *Proceedings of Entomological Society of Washington*, 108 (3): 502–510.
- Minaei, K. 2013. Thrips (Insecta, Thysanoptera) of Iran: a revised and updated checklist. *Zookeys*, 330: 53–74.
- Mirzayans, A., Borumand, H., Zairi, M. and Rajabi, Gh. 1976. Insect fauna from province of Fars (Iran) (1). *Journal of Entomological Society of Iran*, 3 (1&2): 109–135.
- Mirzayans, H. 1995. *Insects of Iran, The list of Homoptera: Auchenorrhyncha in the Insect Collection of plant Pests and Diseases Research Institute*. Ministry of Agriculture, Agricultural Research, Iran, 63 pp.
- Modarres Awal, M. 1994. *List of Agricultural Pests and Their Natural Enemies in Iran*. Ferdowsi University of Mashhad publication, No 147, Mashhad, 364 pp.
- Moosavi Mahvelati N. and Modarres Awal, M. 2011. Faunistic study of Auchenorrhyncha in sugar beet fields of Mashhad and Chenaran. *Journal of Plant Protection*, 25 (3): 34.
- Mossadegh, M.S. and Kocheili, F. 2003. *A semi descriptive checklist of identified species of Arthropods (Agricultural, Medical) and other pests from Khuzestan, Iran*. ShahidChamran University Press, Ahvaz, 475 pp.
- Mozaffarian, F. and Taghizadeh, M. 2010. The first report of three leafhoppers and planthoppers from central parts (Tehran, Semnan, Ghom, Ghazvin and Makazi provinces) of Iran. Proceedings of 19th congress of Irnian Plant Protection, 31 July–03 August 2010, Tehran, Vol. I. Pests, p. 129.
- Mozaffarian, F. and Wilson, M.R. 2016. A checklist of the leafhoppers of Iran (Hemiptera: Auchenorrhyncha: Cicadellidae). *Zootaxa*, 4062 (1): 1–63.
- Nielson, M.W. 1968. *The Leafhopper Vectors of Phytopathogenic Viruses (Homoptera, Cicadellidae): Taxonomy, Biology, and Virus Transmission*. United States Department. Agriculture Technical Bulletin 1382, 386 pp.
- Naseri, B., Fathipour, Y. and Talebi, A.A. 2007a. Comparison of some biological aspects of Empoasca decipiens (Hom.: Cicadellidae) on four bean species. *Journal of Entomological Society of Iran*, 27 (1): 1–13.
- Naseri, B., Fathipour, Y. and Talebi, A.A. 2007b. Sexual parasitism of Empoasca decipiens by Anagrus atomus on four bean species in Tehran area. *Applied Entomology and Phytopathology*, 75 (1): 1–11.
- Nast, J. 1972. *Palaearctic Auchenorrhyncha (Homoptera) An annotated check list*. Polish Scientific Publishers, Warsaw, Poland, 550 pp.
- Nematollahi, M.R. and Khajehali, J. 2000. Major leafhoppers and planthoppers (Hom.: Auchenorrhyncha) of corn fields in Isfahan. Proceedings of 13th congress of Irnian Plant Protection, 2–7 September 1995, Isfahan University of Technology, Vol. I. Pests, p. 235.
- Noori, P. and Javan Moghaddam, H. 1998. Population fluctuation of *Empoasca decipiens* Paoli on cotton in Varamin. Proceedings of 12th congress of Irnian Plant Protection, 23–27 August 1998, University of Tehran, Karaj, Vol. I. Pests, p. 68.
- Nourbakhsh, S., Sahraian, H., Soroush, M.J., Rezaii, V. and Fotouhi, A.R. 2012. *List of important pests, diseases and weeds of major agricultural products, chemicals and recommended ways for their control*. Plant Protection organization, Ministry of Jihad-e Agriculture, 204 pp.
- Olyaei Torshiz, A., Karimzadeh Isfahani, G., Mohajeran, M. and Feizpur, R. (2011) Investigating fauna of Cicadellidae family leafhoppers in grape gardens Khalilabad town in Khorasan-e-Razavi province. Proceedings of 2nd Iranian Pest Management Conference, 14–15 September 2011, Shahid Bahonar University of Kerman, p. 80.
- Omidi, M., Hosseini-Pour, A., Rahimian, H., Massumi, H. and Saillard, C. 2011. Identification of *Circulifer haematoceps* (Hemiptera: Cicadellidae) as vector of *Spiroplasma citri* in the Kerman province of Iran. *Journal of Plant Pathology*, 93 (1): 167–172.
- Pakarpour, F. and Nozari, J. 2011. The first report of *Stirellus lahorensis* (Distant, 1918) (Hemiptera: Cicadellidae) from Iran, Sistan and Baluchestan province. Proceedings of 2nd Iranian Pest Management Conference,

- 14–15 September 2011, Shahid Bahonar University of Kerman, p. 64.
- Pakarpour, F., Nozari, J and Hosseininaveh, V. 2011a. A faunistic study of the Iranian leafhoppers (Hemiptera: Cicadellidae): Kerman province. Proceedings of 2nd Iranian Pest Management Conference, 14–15 September 2011, Shahid Bahonar University of Kerman, pp. 371–411.
- Pakarpour, F., Nozari J. and Hosseininaveh, V. 2011b. Identification and introduction of six new records of Cicadellidae (Hem: Auchenorrhyncha) from Iran. Proceedings of the Global Conference on Entomology (GCE), Chiang Mai, Thailand, p. 218.
- Pakarpour, F., Nozari, J. and Hosseininaveh, V. 2011c. A faunistic study of the Iranian leafhoppers (Hemiptera: Cicadellidae): Kerman province. Proceedings of the Global Conference on Entomology (GCE), Chiang Mai, Thailand, p. 226.
- Pakarpour, F. and Nozari, J. 2014a. Studies on taxonomy and distribution of genus *Exitianus ball* (Hemiptera: Cicadellidae) from Kerman and Sistan and Baluchestan provinces, with three new records for Iranian leafhoppers fauna. Proceedings of 3rd Iranian Pest Management Conference, 21–22 January 2014, Shahid Bahonar University of Kerman, pp. 383–393.
- Pakarpour, F. and Nozari, J. 2014b. The first record of genus, sub genus and species of *Diplocolenus (Diplocolenus) uniformis* Anufriev, 1970, for Iranian Cicadellids fauna from Kerman province. Proceedings of 3rd Iranian Pest Management Conference, 21–22 January 2014, Shahid Bahonar University of Kerman, p. 626.
- Pakarpour, F., Nozari, J., Hosseininaveh, V. and Almasi, A. 2014a. Morphological study on a leafhopper genus *Neosaliturus* Distant, 1918 (Hemiptera, Cicadellidae) by associating molecular methods to confirm identification, with the first record species of Cicadellidae for Iranians leafhoppers fauna. Proceedings of 21th congress of Irnian Plant Protection 23–26 August 2014, University of Urmia, Vol. I. Pests, p. 752.
- Pakarpour, F., Nozari, J. and Hosseininaveh, V. 2014b Genetic variation among populations of *Euscelis alsius* Ribout (Hemiptera: Cicadellidae) in south of Iran, Kerman province. Proceedings of 3rd Iranian Pest Management Conference, 21–22 January 2014, Shahid Bahonar University of Kerman, p. 508.
- Pakarpour Rayeni, F., Nozari, J., and Seraj, A.A. 2015a. A checklist of Iranian Deltocephalinae (Hemiptera: Cicadellidae). *Iranian Journal of Animal Biosystematics*, 11(2): 121–148.
- Pakarpour Rayeni, F., Seraj, A.A., and Nozari, J. 2015b. First report of Six species of leafhoppers (Hemiptera: Auchenorrhyncha: Cicadellidae) in Iranian fauna. Proceedings of 1st Iranian International Congress of Entomology, Tehran, p. 96.
- Pakarpour Rayeni, F., Seraj, A.A., Nozari, J. and Wilson, M.R. 2015c. First record of *Aconurella nuristana* Dlabola (Hemiptera: Cicadellidae) from Iran. *Journal of Crop Protection*, 4 (2): 225–229.
- Rajabi, Gh. 1991. *Insects attacking rosaceous fruit trees in Iran*. Vol. 3: Homoptera. Ministry of Agriculture, Iran, 256 pp.
- Rajabi, Gh. and Mirzayans, H. 1989. First report of *Zyginella pulchra* Low as a harmful insect on deciduous fruit trees in Iran. *Applied Entomology and Phytopathology*, 56 (1&2): 101–103.
- Ramezani L., Mosadegh M.S. and Shishehbor P. 2006. The fauna of super family Eriophyoidea of Khuzestan province. *The Scientific Journal of Agriculture*, 29 (3): 139–149.
- Rezaei, F., Hatami, B., Norbakhsh, S.H. and Jalalizand, A.R. 2014 Identifying the Dominant Leafhoppers in Four Field: Early Sowing Wheat, Late Sowing Wheat, Barley and Maize; and Their Frequency in Shahrekord City. *Agriculture Science Developments*, 3 (3): 85–87.
- Ribaut, H. 1936. *Homopteres Auchenorrhynques. I. (Typhlocybidae)*. Faune de France, Paris: Lechevalier, 31, 111 + 231 pp.
- Ribaut, H. 1952. *Homopteres Auchenorrhynques, II (Jassidae)*, Faune de France, 57. Paul Lechevalier, Paris, 1–474.
- Saeidi, K. and Azura Adam, N. 2011. A survey on pest insect fauna of safflower fields in the Iranian province of Kohkilouyeh and

- Boyerahmad. *African Journal of Agricultural Research*, 6 (19): 4441–4446.
- Salehi, M. and Izadpanah, K. 1992. Etiology and Transmission of Sesame, Phyllody in Iran. *Journal of Phytopathology*, 135 (1): 37–47.
- Salehi, M., Izadpanah, K. and Taghizadeh, M. 2000. A study on host range and possible vector of Lime witches' broom in Iran. Proceedings of 13th congress of Irnian Plant Protection, 2–7 September 1995, Isfahan University of Technology, p. 146.
- Samavi, S., Askar, M., Bagheri, A., Faghihi, M.M. and Salehi, M. 2012. Comparison of the effect of three insecticides on population reduction of leafhopper vector of witches' broom disease of lime. Proceedings of 20th congress of Iranian Plant Protection, 25–28 August 2012, Shiraz University, Vol. I. Pests, p. 290
- Shamszadeh, M. 2002. First record of *Achrusta ghizadehi* Dlab: and *Orosius albicinctus* Distant. (Hom. Cicadellidae) on *Haloxylon* in Iran. Proceedings of 14th congress of Irnian Plant Protection, 7–11 September 2002, Razi University of Kermanshah, Vol. I. Pests, p. 129.
- Sirjani, M. 2012. Determination of economic injury level of cotton leafhopper, *Empoasca decipiens* in Kashmar, Proceedings of 20th congress of Iranian Plant Protection, 25–28 August 2012, Shiraz University, Vol. I. Pests, p. 709.
- Taghizadeh, M. 2012. First report of *Amrascabiguttula biguttula* (Ishida, 1913) (= *Empoasca devastans* Distant, 1918) (Order Hemiptera: Cicadellidae: Typhlocybinae) from Zarrindasht/Fars/Iran cotton fields. Proceedings of 20th congress of Iranian Plant Protection, 25–28 August 2012, Shiraz University, Vol. I. Pests, p. 107.
- Taghizadeh, M. and Salehi, M. 2002. Leafhoppers of subfamily Typhlocybinae found on almond (*Prunus amygdalus*) in Fars province, Proceedings of 14th congress of Irnian Plant Protection, 7–11 September 2002, Razi University of Kermanshah, Vol. I. Pests, pp. 92–93.
- Tahrir Adabi, S., Sadeghi, S.E. and Bagheri, R. 2013. Influence of Irrigation Regimes and Poplar Clones on Densities of *Empoasca decedens* (Hom.: Cicadellidae), *Salicicola kermanensis* (Hom.: Diaspididae) and *Diaspidiotus slavoncus* (Hom.: Diaspididae) in Karaj, Iran. *Greener Journal of Agricultural Sciences*, 3 (8): 644–651.
- Weintraub, P.G. and Beanland, L. 2006. Insect vectors of phytoplasmas. *Annual Review of Entomology*, 51: 91–111.
- Yarmand, H., Sadeghi, S.E., Mohammadi, M., Ebrahimi, S.S. and Seif Allahi, A.R. 2006. Survey of arthropods associated with range medicinal plants in Tehran, Ghazvin and Isfahan provinces. *Iranian Journal of Forest and Range Protection Research*, 3 (2): 107–227.
- Zachvatkin, A.A. 1945. Homoptera-Cicadina from north-western Persia I. *Revue d'Entomologie de l'URSS*, 28 (3–4): 106–115.
- Zohdi, H., Taghizadeh, M. and Mardi, M. 2012. Determine the dominant species leafhoppers citrus orchards in Kerman province. Proceedings of 20th congress of Iranian Plant Protection, 25–28 August 2012, Shiraz University, Vol. I. Pests, pp. 197.
- Zahniser, J.N. (2007–present) An online interactive key and searchable database of Deltocephalinae (Hemiptera: Cicadellidae). Available from: <http://zahniser.speciesfile.org/> (accessed 20 June 2016).

مطالعه زنجرک‌های خانواده Cicadellidae (Auchenorrhyncha) استان خوزستان، جنوب غربی ایران

فرزاد پاکارپور راینی^{۱*}، علی اصغر سراج^۲، جاماسب نوذری^۲

۱ گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه شهید چمران اهواز، ایران

۲ گروه گیاه‌پزشکی، پردیس کشاورزی و منابع طبیعی، دانشگاه تهران، کرج، ایران

* پست الکترونیکی نویسنده مسئول مکاتبه: pakarpour@ut.ac.ir

تاریخ دریافت: ۳۰ تیر ۱۳۹۵، تاریخ پذیرش: ۱۸ مرداد ۱۳۹۵، تاریخ انتشار: ۱۹ مرداد ۱۳۹۵

چکیده: در پژوهش حاضر، فون زنجرک‌های خانواده Cicadellidae (Auchenorrhyncha) در مناطقی از استان خوزستان (جنوب غربی ایران)، طی سال‌های ۱۳۹۳ و ۱۳۹۴ مورد بررسی قرار گرفت. در مجموع ۹۰ گونه متعلق به ۴۹ جنس و ۷ زیر خانواده جمع‌آوری و شناسایی شد. اغلب نمونه‌برداری‌ها با استفاده از تور حشره‌گیری و تله مالايز انجام شد و در برخی موارد نمونه‌ها با استفاده از آسپیراتور و تله زرد چسبنده جمع‌آوری شدند. در نتیجه این تحقیق مشخص شد که از این تعداد نمونه ۴۷ گونه آن برای اولین بار از استان خوزستان گزارش می‌شوند و همچنین ۸ گونه نیز گزارش‌های جدیدی برای فون حشرات ایران هستند؛ که عبارتند از: *Euscelidius Hardya (Hardya) melanopsis* (Hardy, schenckii Kirschbaum, 1868) *Paradorydium (Paradorydium) lanceolatum* (Burmeister, 1850) *Psammotettix Hecalus prasinus* (Matsumura, 1905) ۱۸۳۹ *Grammacephalus pallidus* (Linnauvori, emarginatus Singh, 1969) *Helionidia punctulata* (Ramakrishnan & Menon, 1974) ۱۹۷۸ و *Eupteryx (Eupteryx) kaghanensis* Ahmed, 1969 پراکنش جغرافیایی تمامی گونه‌ها در ایران به تفصیل ارایه شده و همچنین برای هر یک از ۸ گزارش جدید، ویژگی‌های شکل‌شناسی به همراه تصاویر اصلی افراد بالغ و ترسیم‌های اصلی اندام نرینه ارایه شده است.

واژگان کلیدی: زنجرک‌ها، فون، خوزستان، پراکنش، گزارش‌های جدید، ایران.