



Dermestidae (Insecta: Coleoptera) of Niavaran Museum with a new record for Iran

Shima Shahrabi¹, Samin Seddigh^{2*}, Bita Sodaie¹ and Marcin Kadej³

¹ Department of Archaeology, College of human Science, Varamin-Pishva Branch, Islamic Azad University, Varamin, Iran.

² Department of Plant Protection, College of Agriculture, Varamin-Pishva Branch, Islamic Azad University, Varamin, Iran.

³ Department of Invertebrate Biology, Evolution and Conservation, University of Wrocław, Poland.

ABSTRACT. Museum collections are national assets of any country and also are very susceptible to pest damage. Moreover pests in a museum, library or archive environment can cause severe damage to highly valuable and irreplaceable materials. Hence, constantly monitoring of the collections against pest activity is essential in museums. Dermestidae family is known as one of serious pests in the museum collections which feed on various products and cause high infestation. In current study, we present the results of monitoring of all parts of two buildings of The Niavaran Palace Complex, including the private library and Ahmad Shahi Pavilion in Tehran, Iran. The following species of the dermestids have been collected and recorded for the first time from Tehran and Niavaran Museum: *Thylo-drias contractus* Motschulsky, 1839, *Attagenus lobatus* Rosenhauer, 1856, *Anthrenus (Anthrenops) coloratus* Reitter, 1881, *A. (Anthrenus) picturatus picturatus* Solsky, 1876. Moreover, *Thylo-drias contractus* Motschulsky, 1839 is recorded for the first time from Iran.

Key words: Dermestidae, Iran, New record, Niavaran museum

Received:
22 March, 2018

Accepted:
02 October, 2018

Published:
07 October, 2018

Subject Editor:
Hiva Naserzadeh

Citation: Shahrabi, Sh., Seddigh, S., Sodaie, B. & Kadej, M. (2018) Dermestidae (Insecta: Coleoptera) of Niavaran Museum with a new record for Iran. *Journal of Insect Biodiversity and Systematics*, 4 (2), 123–129.

Introduction

Dermestidae (Insecta: Coleoptera) is a diverse family with about 1500 cosmopolitan species worldwide (Háva, 2015b). Dermestids have a variable habitats; their larvae feed mainly on animal materials and dead insects, and few are able to live partly or entirely on vegetable material (i.e. *Trogoderma granarium* Everts, 1989, (Kadej, 2012a)). Sometimes the adults are found in the same habitats as the larvae and feed on the same materials like it is in case of *Dermestes* Linnaeus, 1758 and *Thylo-drias*

Motschulsky, 1839. But in most cases the adults do not feed or they may feed on pollen and nectar (Peacock, 1993). Dermestidae (mainly larvae) can feed on various on a variety of dead animals and animal products, such as skeletons, skins, animal hair, feathers, natural fibers, dead insects; occasionally they feed on stored plants' material such as certain spices and grains. For this reason some dermestids are classified as household and museum pests and are well-known all over the world as

Corresponding author: Samin Seddigh, E-mail: seddigh@iauvaramin.ac.ir

Copyright © 2018, Shahrabi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

pests of stored woolen fabrics and other materials (Hinton, 1945; Aitken, 1975; Veer et al., 1996; Black, 2004; Feroz et al., 2015). Some of the species are associated with carcasses and for this reason might be useful in criminal investigations or in forensic entomology (Catts & Goff, 1992; Peacock, 1993). Many dermestid species are also used in taxidermy and by natural history museums to clean animal skeletons (Ghahari & Háva, 2017).

Till date, one hundred twenty three species representing 12 genera (Table 1) belonging to the family Dermestidae have been reported from Iran.

Currently it comprises several buildings and a museum including: Niavaran Palace Museum (completed in 1968), Ahmad Shahi Pavilion (or Kushk-e Ahmad-Shahi), Sahebqaraniyeh Palace (from the time of Nasir al-Din Shah of Qajar dynasty), Jahan Nama museum and the private library, and other cultural, historical and natural attractions including the Blue Hall, Private Cinema, Jahan Nama Gallery, and Niavaran Garden. In the present paper we list the species of Dermestidae that have been found indoors of two buildings of Niavaran Place Complex (The library and The Ahmad Shahi Pavilion).

Table 1. The List of Dermestidae genera which are found in Iran.

Index	Genus	References
1	<i>Dermestes</i> Linnaeus, 1758	Háva & Kalík, 1999; Háva, 2007; Háva & Herrmann, 2011; Bulak et al., 2013
2	<i>Rhopalosilpha</i> Arrow, 1929	Háva, 2007, 2015b
3	<i>Thorictus</i> Germar, 1834	Háva, 2007, 2015a; Zhantiev, 2011
4	<i>Attagenus</i> Latreille, 1802	Mroczkowski, 1968; Modarres Awal, 1997; Tezcan et al., 2004; Zhantiev, 2005; Háva, 2007; Háva & Herrmann, 2011
5	<i>Paranovelsis</i> Casey, 1900	Háva & Herrmann, 2010; Háva et al., 2013a; Háva et al., 2014
6	<i>Anthrenus</i> Geoffroy, 1762	Zhantiev, 1976; Háva, 2007, 2009; Herrmann et al., 2010; Kadej & Háva, 2011; Háva et al., 2015
7	<i>Ctesias</i> Stephens, 1830	Háva, 2005; Háva et al., 2014
8	<i>Globicornis</i> Latreille [in Cuvier, 1829]	Mroczkowski, 1968; Zhantiev, 1976; Háva, 2007
9	<i>Hirtomegatoma</i> Pic, 1931	Háva, 2007
10	<i>Megatoma</i> Herbst, 1791	Mroczkowski, 1962
11	<i>Phradonoma</i> Jacquelin du Val, 1859	Tezcan et al., 2004; Háva, 2006, 2007; Herrmann & Háva, 2008; Bulak et al., 2013
12	<i>Trogoderma</i> Dejean, 1821	Modarres Awal, 1997; Abivardi, 2001; Khormali et al., 2002; Háva, 2007; Háva & Herrmann, 2011; Mehrabadi et al., 2011; Bulak et al., 2013; Forghani & Marouf, 2015

Material and methods

Study area

This investigation was carried out in two buildings of Niavaran palace complex including private library and Ahmad Shahi pavilion. Ahmad Shahi covers an area of 800 square meters in two stories with ornamental items made of different materials.

The private library is located in the eastern part of the Palace, in two floors and one underground floor in a 770 square meter area. This library consists of about 23,000 books, and more than 350 artifacts and paintings. All the samples were collected from all parts of these two buildings.

Data collection

Insect samples were collected from all the items in the hall including decorative items, furniture, carpets, signs, curtains, clothes, and also from the surface of the earth precisely in June and July 2017. The specimens were collected by hand or by using the aspirator. Trapping was carried out simultaneously during sampling. Yellow and blue sticky cards and light traps were used for trapping. Insects collected from museum palaces were kept in separate glasses of 96% ethanol and were coded. All the specimens were transferred to Entomology Laboratory of Varamin-Pishva Branch, Islamic Azad University. Identification down to species level has been done by Dr. Marcin Kadej (University of Wrocław, Poland).

Results

In total, four species of Dermestidae, the representatives of three genera have been collected from Niavaran Museum, Tehran. *Thylo-drias contractus* Motschulsky, 1839 is newly recorded for Iran. The list of species is given below.

Subfamily: Trinodinae Casey, 1900

Tribe: Thylo-driini Semenov-Tian-Shanskiy, 1909

Genus: *Thylo-drias* Motschulsky, 1839

***Thylo-drias contractus* Motschulsky, 1839**

Diagnosis. 2.0-3.0 mm in length. Thinly covered with short, pale hairs-clubbed, 9-segmented antennae uniform in thickness along length. Median ocellus between compound eyes. In Male: Resting elytra meet at along top third of length, then diverge to reveal abdomen. Long narrow body with thin appendages. Females are wingless with larviform body. Larvae are about 3.0 mm in length with single transverse row of hairs across the dorsum at each segment (Mertins, 1981).

Chorotype: worldwide (Nardi & Háva, 2013).

Remark: First record for Iran.

Subfamily Attageninae Laporte de Castelnau, 1840

Tribe: Attagenini Laporte de Castelnau, 1840

Genus: *Attagenus* Latreille, 1802

***Attagenus lobatus* Rosenhauer, 1856**

Distribution in Iran: Khuzestan (Modarres Awal, 1997), references with no specific locality given (Mroczkowski, 1968, Zhantiev, 1976; Háva, 2003a, 2007, 2015b).

Distribution outside Iran: Holarctic (Háva, 2015b). Centralasiatic-Europeo Mediterranean; this species was introduced to the USA (Háva et al., 2011; Háva et al., 2013b).

Remark: This species feeds on stored products (Modarres Awal, 1997). First record for Tehran province.

Subfamily: Megatominae Leach, 1815

Tribe: Anthrenini Gistel, 1848

Genus: *Anthrenus* Geoffroy, 1762

***Anthrenus (Anthrenops) coloratus* Reitter, 1881**

Distribution in Iran: Fars (Jahrom, Shiraz) (Karami et al., 2004), Sistan and Baluchestan (Nikshahr) (Herrmann et al., 2010), references with no specific locality given (Herrmann et al., 2010; Bulak et al., 2013; Háva, 2015b).

Distribution outside Iran: Holarctic (Háva, 2015b), Tajikistan; Turano-Europeo-Mediterranean species with extension to Canary Islands, USA (Beal, 1998; Háva, 2003b, 2007).

Remark: First record for Tehran province.

***Anthrenus (Anthrenus) picturatus picturatus* Solsky, 1876**

Distribution in Iran: Northern Iran (Mroczkowski, 1962), references with no specific locality given (Tezcan et al., 2004; Háva, 2007; 2015b; Herrmann et al., 2010; Bulak et al., 2013).

Distribution outside Iran: Afghanistan, Azerbaijan, Corsica, Georgia, Kazakhstan, Kyrgyzstan, Poland, Russia, Slovakia, South Africa (Pretoria), Tajikistan, Turkey, Turkmenistan, Uzbekistan, (Háva, 2015b).

Remark: First record for Tehran province.

Discussion

Some synanthropic species of Dermestidae are classified as serious pests of stored products and museum's collections (Peacock, 1993; Kadej, 2012a, 2012b; Kadej et al., 2013). Some of them are representatives of such genera like *Dermestes* Linnaeus, 1758, *Anthrenus* Geoffroy, 1762, *Attagenus* Latreille, 1802, *Trogoderma* Dejean, 1821 or *Reesa* Beal, 1967. Due to the worldwide distribution the most common and well known species are *Anthrenus* (*Nathrenus*) *verbasci* (Linnaeus, 1767) or *Trogoderma glabrum* (Herbst, 1783). Among species that were found during this survey inside the Niavaran Museum (Tehran, Iran) there were only four species, represented three genera. Only one species, *Thylocladius contractus* Motschulsky, 1839, was recorded as larva while three other species (*Attagenus lobatus* Rosenhauer, 1856, *Anthrenus* (*Anthrenops*) *coloratus* Reitter, 1881 and *A.* (s. str.) *picturatus picturatus* Solsky, 1876) were found as adults. All of above mentioned species have been previously recorded as pests from the private houses, storages, shops, and different collections as well (Peacock, 1993; Modarres Awal, 1997; Beal, 1998). It is suspected that new species will be found in the Niavaran Museum in the nearest future because of rich and variable set of the habitats that can be inhabited by dermestids. Based on these findings providing systematic and detailed inspections of the places where the skin and larder beetles can develop is recommended.

Acknowledgments

This research was supported by the Varamin-Pishva Branch of the Islamic Azad University. We express our sincere

appreciation. We also thank Niavaran palace complex staff for their kindly support and cooperation.

Conflict of Interests

The authors assert that there is no conflict of interest regarding the publication of this paper.

References

- Abivardi, C. (2001) *Iranian entomology - An introduction: Volume 2: Applied Entomology* (Schriftenreihe der Stiftung Franz Xaver Schnyder von Wartensee, Zentralbibliothek Zürich). Germany: Springer.
- Aitken, A.D. (1975) *Insect travellers. Volume I. Coleoptera*. London: Ministry of Agriculture, Fisheries (U. K.). *Technical Bulletin 31*, 31, xvi + 191 pp.
- Beal Jr, R. (1998) Taxonomy and biology of Nearctic species of *Anthrenus* (Coleoptera: Dermestidae). *Transactions of the American Entomological Society*, 124, 271–332.
- Black, J. (2004) Fabric and museum pests. In: Morland, D. (ed.) *Handbook of Pest Control (Mallis A), Ninth Edition*. GIE Media, Inc., pp. 581–633.
- Bulak, Y., Yildirim, E., Kadej, M. & Háva, J. (2013) Contribution to the knowledge of the Dermestidae (Coleoptera) fauna of Turkey. *Turkish Journal of Zoology*, 37, 621–626. <http://doi.org/10.3906/zoo-1212-8>
- Catts, E.P. & Goff, M.L. (1992) Forensic entomology in criminal investigations. *Annual Review of Entomology*, 37, 253–272. <http://doi.org/10.1146/annurev.en.37.0101.92.001345>
- Feroz, M., Tara, J., Háva, J. & Azam, M. (2015) New Faunistics Records of Dermestidae (Coleoptera) from Kargil, India. *International Journal of Pure & Applied Bioscience*, 3, 104–108. <http://doi.org/10.18782/2320-7051.2141>
- Forghani, S.H. & Marouf, A. (2015) An introductory study of storage insect pests in Iran. *Biharean Biologist*, 9, 59–62.
- Ghahari, H. & Háva, J. (2017) A Checklist of the Dermestidae (Coleoptera: Bostrichoidea) of Iran. *The Coleopterists Bulletin*, 71, 612–626. <https://doi.org/10.1649/0010-065X-71.3.612>

- Háva, J. (2003a) Notes on Dermestidae (Coleoptera) with description of eight new species. *Annales Historico-Naturales Musei Nationalis Hungarici*, 95, 19–35.
- Háva, J. (2003b) World catalogue of the Dermestidae (Coleoptera), Studie a Zprávy Oblastního Muzea Praha-východ v Brandýse nad Labem a Staré Boleslavi. *Brandýs nad Labem, District Museum Prague-East*, 1, 1–196.
- Háva, J. (2005) Dermestid beetles of the genus *Ctesias* (Coleoptera: Dermestidae) of Iran. *Studies and reports of District Museum Prague-East, Taxonomical Series*, 1, 69–76.
- Háva, J. (2006) Notes on the genus *Phradonoma* from the Middle East with description of nine new species (Coleoptera: Dermestidae: Megatominae). *Genus*, 17, 79–94.
- Háva, J. (2007) Dermestidae. In: LŐBL, I. & SMETANA, A. (eds.) *Catalogue of Palaearctic Coleoptera. Vol. 4, Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea, Cucujoidea* Denmark.: Apollo Books, Stenstrup, Denmark, pp. 299–320.
- Háva, J. (2009) Two species of the genus *Anthrenus* from Iran (Coleoptera: Dermestidae: Anthrenini). *Vernate*, 28, 359–362.
- Háva, J. (2015a) Contribution to the knowledge of *Thorictus* Germar, 1834 from the Middle East (Coleoptera: Dermestidae: Thorictinae). *Studies and Reports of District Museum Prague-East Taxonomical Series*, 11, 267–275.
- Háva, J. (2015b) *World Catalogue of Insects. Volume 13. Dermestidae (Coleoptera)* Leiden/Boston, Brill. xxvi + 419 pp.
- Háva, J. & Herrmann, A. (2010) New faunistic records of Dermestidae (Coleoptera) - Part 4. *Latvijas Entomologs*, 48, 76–79.
- Háva, J. & Herrmann, A. (2011) New faunistic records of Dermestidae (Coleoptera) - Part 6. *Boletín de la Sociedad Entomológica Aragonesa*, 49, 227–228.
- Háva, J., Herrmann, A. & Kadej, M. (2013a) New Faunistic Records of Dermestidae (Coleoptera) - Part 8. *Arquivos Entomoloxicos*, 8, 309–314.
- Háva, J., Herrmann, A. & Kadej, M. (2013b) New faunistic records of Dermestidae (Coleoptera)-Part 7. *Arquivos Entomoloxicos*, 8, 135–140.
- Háva, J., Herrmann, A. & Kadej, M. (2014) New Faunistic Records of Dermestidae (Coleoptera) - Part 11. *International Journal of Fauna and Biological Studies*, 1, 10–13.
- Háva, J., Herrmann, A. & Kadej, M. (2015) New Faunistic Records of Dermestidae (Coleoptera) - Part 14. *Euroasian Entomological Journal*, 14, 568–570.
- Háva, J. & Kalík, V. (1999) The dermestid beetles (Coleoptera: Dermestidae) of Iran. Part 1: genus *Dermestes*. *Acta Societatis Zoologicae Bohemicae*, 63, 443–450.
- Háva, J., Nardi, G., Nardi, G., Whitmore, D., Bardiani, M., Birtele, D., Mason, F., Spada, L. & Cerretti, P. (2011) First contribution to the knowledge of the Dermestidae of Sardinia (Coleoptera). *Biodiversity of Marganai and Montimannu (Sardinia). Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati*, 5, 407–446.
- Herrmann, A. & Háva, J. (2008) Two new *Phradonoma* species (Coleoptera: Dermestidae) from Iran. *Journal of Entomological Society of Iran*, 28, 87–91.
- Herrmann, A., Háva, J. & Fallahzadeh, M. (2010) Contribution to the knowledge of *Anthrenini* from Iran (Coleoptera: Dermestidae). *Journal of the Entomological Research Society*, 12, 113–115.
- Hinton, H. (1945) *A monograph of the beetles associated with stored products. Volume 1.* London: Order of the Trustees of the British Museum. viii + 443 pp.
- Kadej, M. (2012a) Detailed Description of the Morphology of the Last Instar Larva of *Trogoderma megatomoides* Reitter, 1881 (Dermestidae: Megatominae: Megatomini) with Comparison to Related Species. *Journal of the Kansas Entomological Society*, 85, 5–13. <https://doi.org/10.2317/JKES110707.1>
- Kadej, M. (2012b) Detailed morphological description of the mature larva of *Anthrenus latefasciatus* Reitter, 1892 (Dermestidae: Megatominae: Anthrenini) with comparisons to related species. *Zootaxa*, 3270, 31–40.

- Kadej, M. & Háva, J. (2011) Three new species of *Anthrenus pimpinellae* species group from Palaearctic region (Coleoptera: Dermestidae: Megatominae: Anthrenini). *Studies and Reports of District Museum Prague-East Taxonomical Series*, 7, 241–248.
- Kadej M., Jaroszewicz, S. & Tarnawski, D. (2013) Morphology of Mature Larvae of Three Species of the Genus *Anthrenus* (Dermestidae: Megatominae: Anthrenini) With Comparisons To Related Species. *Annals of the Entomological Society of America*, 13, 706–718.
<https://doi.org/10.1603/AN13032>
- Karami, Z., Bagheri Zenooz, E. & Zhantiev, R. D. (2004) Notes on the *Anthrenus* species (Col.: Dermestidae) in Fars province. *Proceedings of 16th Iranian Plant Protection Congress, 2004, 27 August - 1 September, Tabriz, Iran. Ministry of Agriculture, Tehran.*
- Khormali, S., Taheri, M.S. & Borumand, H. (2002) Faunal investigation on stored pests in Gonbad and Minoodasht. *Journal of Entomological and Phytopathological Appliqata*, 70, 13–23. [in Persian, English abstract].
- Mehrabadi, M., Bandani, A., Saadati, F. & Mahmudvand, M. (2011) α -Amylase activity of stored products insects and its inhibition by medicinal plant extracts. *Journal of Agricultural Science and Technology*, 13, 1173–1182.
- Mertins, J. W. (1981) Life History and Morphology of the Odd Beetle, *Thylocladius contractus*. *Annals of the Entomological Society of America*, 74, 576–581.
<https://doi.org/10.1093/aesa/74.6.576>
- Modarres Awal, M. (1997) Dermestidae. In: Modarres Awal, M. (ed.) *List of Agricultural Pests and their Natural Enemies in Iran*. Mashhad, Iran: Ferdowsi University Press.
- Mroczkowski, M. (1962) Dermestidae (Coleoptera) from the Kazakh SSR [Dermestidae (Coleoptera) Kazachskiej SRR]. *Annales Zoologici*, 20, 229–259.
- Mroczkowski, M. (1968) Rozmieszczenie Dermestidae (Coleoptera) na świecie wraz z katalogiem wszystkich znanych gatunków [Distribution of the Dermestidae (Coleoptera) of the world with a catalogue of all known species]. *Annales Zoologici*, 26, 15–19. [in Polish].
- Nardi, G. & Háva, J. (2013) Italian Dermestidae: notes on some species and an updated checklist (Coleoptera). *ZooKeys*, 360, 45–81.
<https://doi.org/10.3897/zookeys.360.6023>
- Peacock, E.R. (1993) *Adults and larvae of hide, larder and carpet beetles and their relatives (Coleoptera: Dermestidae) and of derodontid beetles (Coleoptera: Derodontidae)*. Handbooks for the Identification of British Insects, Vol.5 (3). 144 pp.
- Tezcan, S., Karsavuran, Y., Pehlivan, E. & Háva, J. (2004) Contribution to the Dermestidae (Coleoptera) fauna of Turkey alongwith new records. *Turkish Journal of Entomology*, 28, 27–37.
- Veer, V., Negi, B. & Rao, K. (1996) Dermestid Beetles and some other Insect Pests Associated with Stored Silkworm Cocoons in India, Including a World List of Dermestid Species Found Attacking this Commodity. *Journal of Stored Products Research*, 32, 69–89.
[https://doi.org/10.1016/0022-474X\(95\)00032-3](https://doi.org/10.1016/0022-474X(95)00032-3)
- Zhantiev, R. (1976) Zhuki kozheedy fauny SSSR. [The skin eaters family Dermestidae of fauna of the USSR.]. *Moskva Izdatelstvo Moskovskogo Universiteta*. [in Russian]
- Zhantiev, R.D. (2005) Novye i maloizuchennyye vidy zhukov kozheedov roda *Attagenus* (Coleoptera, Dermestidae). [New and little-known species of the genus *Attagenus* (Coleoptera, Dermestidae).] *Zoologicheskij Zhurnal*, 84, 1147–1152. [in Russian, English summary]
- Zhantiev, R.D. (2011) Novye i maloizuchennyye vidy zhukov-kozheedov roda *Thorictus* (Coleoptera, Dermestidae). [New and little known dermestidbeetles of the genus *Thorictus* (Coleoptera, Dermestidae).] *Zoologicheskij Zhurnal*, 90, 706–711. [in Russian, English summary]

سوسک‌های خانواده (Insecta: Coleoptera) Dermestidae موزه نیاوران به همراه یک گزارش جدید برای ایران

شیمما شهرابی^۱، ثمین صدیق^{۲*}، بیتا سودایی^۱ و مارسین کادج^۳

۱ گروه باستان‌شناسی، دانشکده علوم انسانی، دانشگاه آزاد اسلامی واحد ورامین-پیشوا، ورامین، ایران.

۲ گروه گیاهپزشکی، دانشکده کشاورزی، دانشگاه آزاد اسلامی واحد ورامین-پیشوا، ورامین، ایران.

۳ گروه بیولوژی مهره‌داران، تکامل و حفاظت، دانشگاه وروکلاو، لهستان.

* پست الکترونیکی نویسنده مسئول مکاتبه: seddigh@iauwaramin.ac.ir

تاریخ دریافت: ۰۲ فروردین ۱۳۹۷، تاریخ پذیرش: ۱۰ مهر ۱۳۹۷، تاریخ انتشار: ۱۵ مهر ۱۳۹۷

چکیده: آثار موجود در موزه‌ها دارایی‌های ملی هر کشوری هستند که نسبت به بسیاری از آفات نیز آسیب پذیرند. علاوه بر این آفات می‌توانند در یک موزه، کتابخانه یا یک انبار خسارات جدی در اشیاء باارزش و غیرقابل جایگزین ایجاد کنند. از این رو، نظارت مداوم بر مجموعه‌ها جهت بررسی فعالیت آفات در موزه‌ها ضروری است. خانواده Dermestidae به عنوان یکی از آفات جدی در مجموعه موزه شناخته شده است که بر روی محصولات مختلف تغذیه کرده و باعث آلودگی زیاد می‌شوند. در این تحقیق، ما نتایج نظارت بر تمام بخش‌های دو ساختمان مجموعه کاخ نیاوران تهران، شامل کتابخانه سلطنتی و کوشک احمد شاهی را ارائه می‌کنیم. گونه‌های زیر از خانواده Dermestidae برای اولین بار از تهران و موزه نیاوران جمع‌آوری شده است: *Attagenus lobatus* Rosenhauer, *Thylophorus contractus* Motschulsky, 1839 A. (*Anthrenus*) و *Anthrenus (Anthrenops) coloratus* Reitter, 1881. 1856 *Thylophorus contractus picturatus picturatus* Solsky, 1876. علاوه بر این گونه *Thylophorus contractus* Motschulsky, 1839 برای اولین بار از ایران گزارش شد.

واژگان کلیدی: Dermestidae، ایران، گزارش جدید، موزه نیاوران