



ESI

Spatulaphorus geotruporum (Acari: Pygmephoridae) a new record for mites fauna of Asia, with some notes on the genus

Sarina Seyedein, Vahid Rahiminejad* and Ahmad Nadimi

Department of Plant Protection, Faculty of Plant Production, Gorgan University of Agricultural Science and Natural Resource, Golestan, Iran.

Received:
24 April, 2019

Accepted:
15 May, 2019

Published:
21 May, 2019

Subject Editor:
Hamidreza Hajiqanbar

ABSTRACT. During a comprehensive survey of the heterostigmatic mites associated with dung beetles in the hyrcanian forests, northern Iran, several colonies of the pygmephoroid mites including four species of the genus *Spatulaphorus* Rack, 1993, *S. copridis* Khaustov, 2007, *S. gorganica* Rahiminejad & Hajiqanbar, 2011, *S. vladimiri* Khaustov, 2005 and *S. geotruporum* Khaustov, 2005 phoretic on *Geotrupes spiniger* (Marsham, 1802) were found. This is the first record of *S. geotruporum* from Asia. Furthermore, this is the first record of phoresy of *S. geotruporum* on *G. spiniger*.

Key words: Heterostigmata, Dung beetles, Phoresy, Host preference, Iran

Citation: Seyedein, S., Rahiminejad, V. & Nadimi, A. (2019) *Spatulaphorus geotruporum* (Acari: Pygmephoridae) a new record for mites fauna of Asia, with some notes on the genus. *Journal of Insect Biodiversity and Systematics*, 5 (1), 79–85.

Introduction

Dung beetles are considered as a perceptible component of the diversity of insects in tropical rain forests (Halffter & Matthews, 1966) which have received proportionate attention from entomologists because of their remarkable adaptations in decomposing vertebrate dung pads (Bornemissza, 1960; Villalba et al., 2002). They are utilized by many phoretic mites for dispersal purposes (Peck & Forsyth, 1982). Mites of the family Pygmephoridae Cross, 1965 consist primarily of free-living and generally fungivorous species, which are described based only on adult females (Kaliszewski et al., 1995; Krantz & Walter, 2009); the pygmephorid family is the

second largest family in the superfamily Pygmephoroida and include 31 genera and more than 300 species (Khaustov & Trach, 2018). Some mites of all four families of Pygmephoroida, Pygmephoridae Cross, 1965, Neopygmephoridae Cross, 1965, Microdispidae Cross, 1965 and Scutacaridae Oudemans, 1916, can establish close interaction with geotrupid beetles (Khaustov & Frolov, 2018). Furthermore, representatives of five genera of this family utilize dung beetles for phoretic dispersal: *Pediculaster* Vitzthum, 1927; *Geotrupophorus* Mahunka, 1970; *Pygmephorellus* Cross and Moser, 1971; *Spatulaphorus* Rack, 1993 and *Pseudopygmephorellus* Khaustov, 2008

Corresponding author: Vahid Rahiminejad, E-mail: vahidrahiminejad@gau.ac.ir

Copyright © 2019, Seyedein et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

(Khaustov & Trach, 2012). The genus *Spatulaphorus* Rack, 1993 is comprised of 14 described species and known to have phoretic relationships with different dung beetles from the families Geotrupidae and Scarabaeidae (Rahiminejad et al., 2011; Khaustov & Trach, 2012; Khaustov & Forolov, 2018). Heretofore four species of the genus *Spatulaphorus* have been described and recorded from Iran: *S. gorganica* Rahiminejad and Hajiqaanbar, 2011, *S. copridis* Khaustov, 2007, *S. vladimiri* Khaustov, 2005 and *S. gymnopleuri* Sobhi and Hajiqaanbar, 2017, which were collected from geotrupid beetles, *Geotrupes spiniger* Marsham, 1802 (Rahiminejad et al., 2011, 2013; Katlav et al., 2015) except *S. gymnopleuri* that found on Scarabaeid beetle, *Gymnopleurus flagellates* (F.) (Sobhi et al., 2017).

Following a comprehensive study upon heterostigmatic mites associated with coprophagous beetles in hyrcanian forests of northern Iran, several colonies of aforementioned species as well as *S. geotrurorum* as a new record for acarine fauna of Asia were found. In the current work, we are aiming to report some species of *Spatulaphorus*, also some notes on leg chaetotaxy, distributions and host range of this genus.

Material and methods

The host beetles were collected as a result of several samples from Golestan province (Naharkhoran, Alangdareh, Toshan and Ghorogh Forests and Hezarpich Region), northern Iran in 2018. All the mite species were detached from *Geotrupes spiniger* (Col.: Geotrupidae). Host beetles were captured by a light trap or direct sampling from Oak forests.

The mites were cleared in lactophenol and mounted in Hoyer's medium. A light microscope with phase contrast (Olympus BX51) was used for morphological study of the mites. The taxonomic system of

Pygmephoroida follows Khaustov (2004, 2008); the terminology of idiosoma and legs follows that of Lindquist (1986); the nomenclature of subcapitular setae and the designation of cheliceral setae follow that of Grandjean (1944, 1947), respectively. With using a global positioning system (GPS smart phone device), details of geographical position have been recorded.

Results

In this study four species of *Spatulaphorus* Rack, 1993 identified.

Family Pygmephoridae Cross, 1965

Genus *Spatulaphorus* Rack, 1993

Type species: *Spatulaphorus camerikae* Dastych and Rack, 1993, by original designation.

Spatulaphorus geotrurorum Khaustov, 2005

Material examined. Six specimens of adult females were obtained in bottom of a vial containing twelve of *Geotrupes spiniger* (Col.: Scarabaeoidea: Geotrupidae). The host beetles were collected with direct sampling in Toshan Forest, 36.46° N, 54.25° E, 203 m a.s.l., 12.IX.2018, leg., V. Rahiminejad.

Diagnosis. *Spatulaphorus geotrurorum* can be identified by the reduced apodemes 4 and the subequal setae c_1 and c_2 .

Remarks. This species is recorded from Iran for the first time. Moreover, the association between this mite and *G. spiniger* is new (see Table 2). This species was described from Ukraine (Khaustov, 2005) associated with *G. stercorarius* L..

Spatulaphorus vladimiri Khaustov, 2005

Material examined. Five large colonies of adult females were obtained under the elytra of *G. spiniger*. The host beetles were collected by light trap in Naharkhoran forest, 36.46° N, 54.27° E, 450 m a.s.l., 12.IX.2018 and Hezarpich region, Golestan province, northern Iran, 36.46° N, 54.24° E,

360 m a.s.l., 9.VIII.2018, leg., V. Rahiminejad and S. Seyedein.

Diagnosis. The species can be simply identified by short and foliate setae of sc_2 , c_2 , d , f , h_2 .

***Spatulaphorus copridis* Khaustov, 2007**

Material examined. Two females obtained in bottom of two vials containing *G. spiniger*, submerged in ethyl alcohol 75%. The host beetle was collected with light trap from Oak trees in Alangdareh forest, 36.46° N, 54.26° E, 311 m a.s.l., 21.IX.2018, leg., V. Rahiminejad,.

Diagnosis. *Spatulaphorus copridis*, clearly distinguished from all known species of *Spatulaphorus* by having only one seta (l') on genu I.

***Spatulaphorus gorganica* Rahiminejad & Hajiqanbar, 2012**

Material examined. Nine large colonies of females obtained under the elytra of *G. spiniger*. The host beetles were collected with light trap from Oak trees in Naharkhoran forest, 36.46° N, 54.27° E, 409 m a.s.l., 5.VII.2018; Alangdareh forest, 36.46° N, 54.26° E, 317 m a.s.l., 18.VII.2018 and Toshan Forest, 36.46°N, 54.25°E, 206 m a.s.l., 21.VIII.2018, leg., V. Rahiminejad and S. Seyedein.

Diagnosis. *Spatulaphorus gorganica*, can be identified by having modified and spine-like setae pl'' on tarsi II–IV; setiform setae sc_2 , c_2 , d , f and h_1 ; setae c_1 much longer than setae e and h_2 ; seta tc' on tarsus II setiform and very short and thin setae e and h_2 , versus long setae f and h_1 on the EF and H tergites.

Discussion

It is worth to be noted the state of possessing only one seta on genu I (absence of l'') is a unique autapomorphic character for *S. copridis* that may implies this to be considered as the most derivative species

within the genus; In the other hand, modified (Spine or Foliate -like) setae tc' , pl'' (on tarsus III) and pl''' (on tarsus IV) are another evidences that indicate various speciation paths in this genus (Table 1).

Among the pygmephorid mites associated to Coleoptera, females of *Geotrupophorus* and *Spatulaphorus* have most association with geotrupid beetles, and several species of *Elattoma* and *Pygmephorellus* are associated with a variety of subcortical and dung-inhabiting beetles (Krantz & Walter, 2009). All members of the genus *Spatulaphorus* have been recorded to be exclusively phoretic on coprophagous beetles belonging to the families Geotrupidae and Scarabaeidae (*Geotrupes* (Col.: Geotrupidae), *Copris*, *Onitis*, *Catharsius* (Col.: Scarabaeidae: Coprinae) and *Gymnopleurus* (Col.: Scarabaeidae: Gymnopleurini)) (Table 2). Heretofore the four species of the genus *Spatulaphorus*, *S. gorganica*; *S. copridis*, *S. vladimiri* and *S. geotruporum* have been described and recorded from Iran and Ukraine and all of them were collected from the dung beetle *Geotrupes spiniger*. This may corroborate the status of the beetles of the family Geotrupidae as a preferential host for these mites. However more investigations are required to ascertain whatever potential dung inhabiting host families they have phoretic relationship with.

Acknowledgments

The study was supported by the Gorgan University of Agricultural Science and Natural Resource.

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

Table 1. Leg setation difference among species of the genus *Spatulaphorus* (females).

	<i>S. geotrupi</i>	<i>S. foliatus</i>	<i>S. langi</i>	<i>S. camerikae</i>	<i>S. luriei</i>	<i>S. geotruporum</i>	<i>S. vladimiri</i>	<i>S. copridis</i>	<i>S. gorganica</i>	<i>S. venustus</i>	<i>S. enoplotruporum</i>	<i>S. altaicus</i>	<i>S. brasiliensis</i>	<i>S. gymnopileuri</i>
Seta <i>l''</i> on Genu I	+	+	+	+	+	+	+	-	+	+	+	+	+	+
Seta <i>u'</i> on Tarsus II	+	-	+	+	+	+	+	+	+	+	-	+	+	+
spine-like seta on Tarsus II	-	+	+	-	-	-	+	-	+	+	-	+	-	-
spine-like seta <i>pl'</i> on Tarsus III	-	+	+	-	-	-	+	-	+	+	-	+	-	-

Table 2. Information about hosts, locality and reference of all known species of the genus *Spatulaphorus*.

Species	Host	Locality	Reference
<i>S. geotrupi</i>	<i>Geotrupes</i> sp. (Geotrupidae)	Hungry Ukraine	Mahunka, 1970 Khaustov, 2005
<i>S. foliatus</i>	<i>Onitis</i> sp. (Scarabaeidae)	Botswana	Dastych & Rack, 1993
<i>S. langi</i>	<i>Catharsius</i> sp. (Scarabaeidae)	Vietnam	Dastych & Rack, 1993
<i>S. camerikae</i>	<i>Catharsius ulysses</i> (Boheman) (Scarabaeidae)	Botswana	Dastych & Rack, 1993
<i>S. luriei</i>	Scarab beetles (Scarabaeidae)	South Africa	Dastych et al., 1997
<i>S. geotruporum</i>	<i>Geotrupes stercorarius</i> L. (Geotrupidae) <i>Geotrupes spiniger</i> (Geotrupidae)	Ukraine Iran	Khaustov, 2005 Present study
<i>S. vladimiri</i>	<i>Geotrupes stercorarius</i> (Geotrupidae) <i>Geotrupes spiniger</i> (Geotrupidae)	Ukraine Iran	Khaustov, 2005 Katlav et al., 2015
<i>S. copridis</i>	<i>Copris lunaris</i> (Mueller) (Scarabaeidae) <i>Copris hispanus</i> (Scarabaeidae) <i>Geotrupes spiniger</i> (Geotrupidae)	Ukraine Iran	Khaustov, 2007 Rahiminejad et al., 2013

Table 2. Continued.

Species	Host	Locality	Reference
<i>S. gorganica</i>	<i>Geotrupes spiniger</i> (Geotrupidae) <i>Geotrupes stercorarius</i> (Geotrupidae)	Iran	Rahiminejad et al., 2011 Khaustov & Trach, 2012
<i>S. venustus</i>	<i>Geotrupes mutator</i> (Marsham, 1802) (Geotrupidae)	Ukraine	Khaustov & Trach, 2012
<i>S. enoplotruporum</i>	<i>Enoplotrupes</i> sp. (Geotrupidae)	Thailand	Khaustov & Frolov, 2018
<i>S. altaicus</i>	<i>Geotrupes baicalicus</i> Reitter (Geotrupidae)	Russia, Altay Republic	Khaustov & Trach, 2018
<i>S. brasiliensis</i>	<i>Dichotomius podalirius</i> Felsche, 1901 (Scarabaeidae)	Brazil, Amazonas	Khaustov & Frolov, 2017
<i>S. gymnopleuri</i>	<i>Gymnopleurus flagellatus</i> (Fabricius, 1787) (Scarabaeidae)	Iran	Sobhi et al., 2017

References

- Bornemissza, G.F. (1960) Could dung eating insects improve our pastures?. *Journal of the Australian Institute of Agricultural Science*, 26, 54–56.
- Dastych, H. & Rack, G. (1993) *Spatulaphorus* Rack gen. n. and three new species of pygmephorid mites associated with scarab beetles in Botswana and Vietnam (Acari: Heterostigmata). *Mitteilungen aus den Hamburgischen Zoologischen Museum und Institut*, 90, 265–84.
- Dastych, H., Rack, G. & Camerik, M. (1997) A new species of genus *Spatulaphorus* Rack (Acari: Heterostigmata, Pygmephoridae) associated with scarab beetles in South Africa. *Mitteilungen aus den Hamburgischen Zoologischen Museum und Institut*, 94, 145–151.
- Grandjean, F. (1944) Observations sur les Acariens de la famille des Stigmaeidae. *Archives des Sciences Physiques et Naturelles*, 26, 103–131.
- Grandjean, F. (1947) L'origine pileuse des mors et la chaetotaxie de la mandibule chez les Acariens actinochitineux. *Comptes rendus des Séances de l'Académie des Sciences*, 224, 1251–1254.
- Halfpeter, G. & Matthews, E.G. (1966) The natural history of dung beetles of the subfamily Scarabaeinae (Coleoptera: Scarabaeidae). *Folia Entomologica Mexicana*, 12–14, 1–312.
- Kaliszewski, M., Athias-Binche, F. & Lindquist, E.E. (1995) Parasitism and Parasitoidism in Tarsonemina (Acari: Heterostigmata) and Evolutionary Considerations. *Advances in Parasitology*, 35, 335–67.
[https://doi.org/10.1016/S0065-308X\(08\)60074-3](https://doi.org/10.1016/S0065-308X(08)60074-3)
- Katlav, A., Hajiqaanbar, H. & Talebi, A.A. (2015) A contribution to the knowledge of heterostigmatic mites (Acari: Prostigmata) in western Mazandaran Province, Northern Iran. *Acarologia*, 55, 311–320.
<http://dx.doi.org/10.1051/acarologia/20152175>
- Khaustov, A.A. (2004) Mites of the family Neopygmephoridae Cross, 1965 stat. n. and their position in Heterostigmata. In: Balashov, Y.S. (Ed.), *VIII Russian*

- Acarological Conference, St. Petersburg. Zoological Institute of RAS, St. Petersburg, p. 137. [in Russian]*
- Khaustov, A.A. (2005) Three new species of mite family Pygmephoridae (Acari: Heterostigmata) associated with scarab beetles (Coleoptera: Scarabaeidae) from Ukraine. *Acarina*, 13 (1), 35–42.
- Khaustov, A.A. (2007) a new species of the genus *Spatulaphorus* (Acari: Heterostigmata: Pygmephoridae) associated with *Copris lunaris* (Coleoptera: Acarabaeidae) from Crimea. *Acarina*, 15 (1), 155–157.
- Khaustov, A.A. (2008) *Mites of the family Scutacaridae of Eastern Palaearctic*. Akademperiodyka, Kiev, 291 pp.
- Khaustov, A.A. & Frolov, A.V. (2017) New species of heterostigmatic mites (Acari: Heterostigmata: Athyreacaridae, Dolicho-cybididae, Pygmephoridae) associated with scarab beetles (Coleoptera: Geotrupidae, Scarabaeidae) from Brazil. *Zootaxa*, 4294 (5), 501–521.
<http://dx.doi.org/10.11646/zootaxa.4294.5.1>
- Khaustov, A.A. & Frolov, A.V. (2018) New taxa of pygmephoroid mites (Acari: Pygmephoroida: Pygmephoridae, Neopygmephoridae) phoretic on *Enoplotrupes sharpi* (Coleoptera: Geotrupidae) from Thailand. *Zootaxa*, 4442 (2), 277–292.
<http://dx.doi.org/10.11646/zootaxa.4442.2.5>
- Khaustov, A.A. & Trach, V.A. (2012) A new species and new records of mites of the genus *Spatulaphorus* Rack (Acari: Heterostigmata: Pygmephoridae) from Ukraine. *International Journal of Acarology*, 38 (6), 480–485.
<http://dx.doi.org/10.1080/01647954.2012.677475>
- Khaustov, A.A. & Trach, V.A. (2018) New species of *Spatulaphorus* (Acari: Heterostigmata: Pygmephoridae) associated with *Geotrupes baicalicus* (Coleoptera: Geotrupidae) from Altai Republic, Russia. *Systematic & Applied Acarology*, 23, 123–131.
<https://doi.org/10.11158/saa.23.1.10>
- Krantz, G.W. & Walter, D.E. (2009) *A Manual of Acarology*. Texas tech university press, 807 pp.
- Lindquest, E.E. (1986) The world genera of Tarsonemidae (Acari: Heterostigmata): A morphological, phylogenetic and systematic revision, with a reclassification of family-group taxa in Heterostigmata. *Memoirs of the Entomological Society of Canada*, 136, 1–517.
<https://doi.org/10.4039/entm118136fv>
- Mahunka, S. (1970) Considerations on the systematics of the Tarsonemina and the description of new European taxa (Acari: Trombidiformes). *Acta Zoologica Hungarica*, 16 (2), 137–174.
- Peck, S.B. & Forsyth, A. (1982) Composition, structure, and competitive behaviour in a guild of Ecuadorian rain forest dung beetles (Coleoptera; Scarabaeidae). *Canadian Journal of Zoology*, 60, 1624–1634.
- Rahiminejad, V., Hajiqanbar, H. & Fathipour, Y. (2011) A new species of the genus *Spatulaphorus* (Acari: Heterostigmatina: Pygmephoridae) phoretic on *Geotrupes spiniger* (Coleoptera: Geotrupidae) from Iran. *Annales Zoologici*, 61 (3), 547–551.
<http://dx.doi.org/10.3161/000345411X603391>
- Rahiminejad, V., Hajiqanbar, H. & Fathipour, Y. (2013) First record of the genus *Spatulaphorus* (Acari: Pygmephoridae) associated with dung beetles (Coleoptera: Scarabaeidae) from Iran. *Entomological Society of Iran*, 33 (2), 81–82.
- Sobhi, M., Hajiqanbar, H. & Mortazavi, A. (2017) New species and records of heterostigmatic mites (Acari: Prostigmata: Heterostigmata) phoretic on scarabaeid dung beetles (Coleoptera: Scarabaeidae) from northwestern Iran. *Zootaxa*, 4276 (3), 427–434.
<https://doi.org/10.11646/zootaxa.4276.3.7>
- Villalba, S., Lobo, J.M., Martin-Piera, F. & Zardoya, R. (2002) Phylogenetic relationships of Iberian dung beetles (Coleoptera: Scarabaeinae): Insights on the evolution of nesting behavior. *Journal of Molecular Evolution*, 55, 116–126.
<https://doi.org/10.1007/s00239-002-2314-4>

اولین گزارش (*Spatulaphorus geotruperum*) (Acari: Pygmephoridae) برای فون کنه‌های آسیا، همراه با بیان برخی نکات در مورد جنس

سارینا سیدین، وحید رحیمی‌نژاد* و احمد ندیمی

گروه گیاهپزشکی، دانشکده تولید گیاهی، دانشگاه علوم کشاورزی و منابع طبیعی گرگان، گلستان، ایران.
* پست الکترونیکی نویسنده مسئول مکاتبه: vahidrahiminejad@gau.ac.ir
تاریخ دریافت: ۰۴ اردیبهشت ۱۳۹۸، تاریخ پذیرش: ۲۵ اردیبهشت ۱۳۹۸، تاریخ انتشار: ۳۱ اردیبهشت ۱۳۹۸

چکیده: در طول بررسی کنه‌های هترواستیگمای مرتبط با سوسک‌های سرگین-غلطان در جنگل‌های هیرکانی شمال ایران، چندین کلنی از کنه‌های خانواده Pygmephoridae شامل چهار گونه از جنس *Spatulaphorus* Rack, 1993، به نام‌های *S. gorganica* Rahiminejad & *S. copridis* Khaustov, 2007، *S. geotruperum* و *S. vladimiri* Khaustov, 2005، *Hajiqanbar*, 2011، *Khaustov*, 2005 بصورت هم‌سفر روی بدن سوسک *Geotrupes spiniger* (Marsham, 1802) پیدا شد. این اولین گزارش *S. geotruperum* از آسیا است. همچنین، این مقاله اولین گزارش هم‌سفری کنه *S. geotruperum* روی سوسک *G. spiniger* است.

واژگان کلیدی: هترواستیگماتینا، سرگین‌غلطان، هم‌سفری، ترجیح میزبانی، ایران