



Checklist of Iranian stored product beetles (Insecta: Coleoptera)

Najmeh Ebrahimi

Entomology Research Department, Iranian Research Institute of Plant Protection, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran.

ABSTRACT. An extensive literature review was performed to determine the list of stored product beetles reported from Iran. A total of 122 species of stored product beetles belonging to 59 genera and 45 tribes have been recorded in Iran. Species–richest families are Dermestidae (24 species) and Tenebrionidae (24 species), followed by Chrysomelidae (15 species), Cryptophagidae (13 species), Ptinidae (10 species), Curculionidae (6 species), Staphylinidae (4 species), Nitidulidae (4 species), Bostrichidae (3 species), Laemophloeidae (3 species), Silvanidae (3 species), Trogossitidae (2 species), Anthicidae (2 species), Cleridae (2 species) and seven families are monospecific: Carabidae, Cerambycidae, Anthribidae, Erotylidae, Latridiidae, Melyridae and Mycetophagidae. The present study is not only a faunistic report but also help in design and development of integrated pest management of stored product beetles.

Key words: Coleoptera, Checklist, Iran, Stored product beetles

Received:
27 January, 2020

Accepted:
16 June, 2020

Published:
29 June, 2020

Subject Editor:
Aref Marouf

Citation: Ebrahimi, N. (2020) Checklist of Iranian stored product beetles (Insecta: Coleoptera). *Journal of Insect Biodiversity and Systematics*, 6 (3), 261–305.

Introduction

Insects are considered as the significant and primary pests of stored products such as vegetable and animal origins, raw and processed cereals, pulses, seeds, spices, dried fruit and nuts, and other dry, durable commodities, raw animal products, dried or smoked fish and meat, skins, hides and museum specimens that present damage to the grains through direct feeding during some of their lifecycles, resulting large populations and considerable damages (Haines, 1991; Weidner & Rack, 1984; Rees, 1990; Shires, 1982). The pests cause significant quantitative and qualitative losses to the multibillion-dollar grain, food, and retail industries each year through their feeding, product adulteration, customer complaints, product rejection at the time of sale, and cost associated with their management (Campbell et al., 2004; Hagstrum et al., 2013; Thind & Dunn, 2002; Olsen, 1998; Stejskal & Verner, 1996; Sinha & Watters, 1985). About 40 percent of insects belonging to order Coleopteran. Some Species great damage too many seeds in grain stores and reduce in the weight and seed quality (Borrer et al., 1989).

Corresponding author: Najmeh Ebrahimi, E-mail: n.ebrahimi@iripp.ir

Copyright © 2020, Ebrahimi. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Coleoptera (Insecta) indicates one of the largest orders of diversity on world with four suborders, 24 superfamilies, 211 families, and over 350,000 described species (Bouchard et al., 2011). Iran forms a large part of the Iranian plateau, and covers an area of 1,623,779 km². It is bordered in the north by the Caucasus Mountains, Middle Asian natural regions and the Caspian Sea (-27 m below sea level), in the west by the Anatolian and Mesopotamian regions, in the east by the eastern part of the Iranian plateau (Afghanistan and adjacent west Pakistan) and the Baluch-Sindian region; and finally in the south by the Persian Gulf and Oman Sea, which are connected by the latter to the Indian Ocean (Zehzad et al., 2002).

Checklists of species are important for any kind of question or investigation related to biodiversity. With this paper, we present a checklist of the beetles (Coleoptera) of stored product pest from Iran. Mound (1989) estimates that the number of beetle species associated with stocks worldwide exceeds 600. However, it is not clear which storage pests are indigenous or intensive species. More than thirty years have passed since the publication of the *Checklist of insects, mites and rodents affecting stored products in Iran* by Shahhosseini & Kamali (1989). During that period, many changes have been made in the classification of Coleoptera. For example, the Scydmaenines, Pselaphines, Scaphidiines, and Micropeplines, which formerly regarded as distinct families, are now part of the Staphylinidae, the Lyctines are included in the Bostrichidae, the Anobiines in the Ptinidae, the Languriines in the Erotylidae, the Bruchines in the Chrysomelidae, the Apionines and Ithycerines in the Brentidae, and the Scolytines and Platypodines in the Curculionidae (Bousquet et al., 2013). Finally, considering the amount of new information available, we believe that a new edition of the checklist of beetles of stored product pest of Iran will help those interested in beetles of the Iran fauna. The aim of this paper is to provide a checklist of Iranian stored product beetles based on available literature data for this area.

Material and methods

The checklist is mainly based on published literature of Iranian researchers. The published data on distribution of the stored product beetles in Iran are summarized by province. Family, subfamily, tribe, genus and species are listed in alphabetical order. All the species and subspecies are listed with their valid names, correct spelling, author and year of description. When accurate data about local distribution in Iran are lacking in a quoted reference, the phrase "no locality cited" is used. Additional data were also provided for some species from Atlas of stored insects and mites (Hagstrum et al., 2013) and stored-product insect resource (Hagstrum & Subramanyam, 2009). Host data (where exist) and distribution data in Iran are taken from published literature especially Modarres Awal (1994, 1997), Shahhosseini & Kamali (1989), research articles, abstracts, books, congress proceeding, Ph. D. dissertations and M. Sc. theses.

Results

Data of 122 species of coleopteran are listed alphabetically below. The number of genera and species into families is (genus-species): from Anthicidae (1-2), Antribidae (1-1), Bostrichidae (2-3), Carabidae (1-1), Cerambycidae (1-1), Chrysomelidae (5-15), Cleridae (1-2), Cryptophagidae (2-13), Curculionidae (3-6), Dermestidae (6-24), Erotylidae (1-1), Laemophloeidae (1-3), Lathridiidae (1-1), Melyridae (1-1), Mycetophagidae (1-1), Nitidulidae (1-4), Ptinidae (7-10), Silvanidae (2-3), Staphylinidae (3-4), Tenebrionidae (16-

24) and Trogossitidae (2-2). The geographical scope of this checklist covers the Iranian provinces and information about each species are taken from the literature.

Checklist of the beetles of stored product pest from Iran

Family: Anthicidae Latreille, 1819

Subfamily: Anthicinae Latreille, 1819

Tribe: Anthicini Latreille, 1819

***Omonadus floralis* (Linnaeus, 1758)**

Distribution in Iran: Mazandaran ([Shahhosseini & Kamali, 1989](#)), Iran (no locality cited) ([Chandler et al., 2008](#)).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America ([Hagstrum et al., 2013](#)).

Host range: Beverage crops, dried fruit, fresh fruit or vegetable, grain, grain product, nut, oilseed, oilseed product, pulse, pulse product, root, other seed, spoiled other commodities and vegetable material ([Hagstrum et al., 2013](#); [Shahhosseini & Kamali, 1989](#)).

***Omonadus formicarius formicarius* (Goeze, 1777)**

Distribution in Iran: Hormozgan ([Bonadona, 1970](#)), Khorasan-e Razavi, Azarbayjan, Markazi, Lorestan ([Telnov & Ghahari, 2018](#)).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico and Oceania ([Hagstrum et al., 2013](#)).

Host range: Empty grain bins ([Hagstrum et al., 2013](#)).

Family: Anthribidae Billberg, 1820

Subfamily: Choraginae Kirby, 1819

Tribe: Araecerini Lacordaire, 1866

***Araecerus fasciculatus* (De Geer, 1775)**

Distribution in Iran: General distributed ([Shahhosseini & Kamali, 1989](#)), Mazandaran ([Salehi Ardakani & Nasserzadeh, 2014](#)).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America ([Hagstrum et al., 2013](#)).

Host range: Beverage crops (coffee bean), bake goods, dried fruit, fresh fruit or vegetable, grain (maize), grain product, medicine, nut, other commodities, oilseed, oilseed product, pulse, root (ginseng, yam) root product, other seed, seasoning, textile and vegetable material ([Hagstrum et al., 2013](#); [Shahhosseini & Kamali, 1989](#)).

Family: Bostrichidae Latreille, 1802

Subfamily: Dinoderinae Thomson, 1863

***Rhyzopertha dominica* (Fabricius, 1792)**

Distribution in Iran: Golestan ([Khormali et al., 2002](#); [Eyidozahi et al., 2013](#)), Tehran ([Mehrabadi et al., 2011](#); [Aref & Valizadegan, 2015](#)), West Azarbaijan ([Mahdneshin et al., 2009](#)) and generally distributed ([Modarres Awal, 1997](#); [Ziaee et al., 2006](#); [Ashouri & Shayesteh, 2009](#)).

General distribution: Probably Indian or Malaysian origin, Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013; Shahhosseini & Kamali, 1989).

Host range: Animal invertebrate and vertebrate, beverage crops, bake goods, dried fruit (apple), fresh fruit or vegetable, grain (barley, maize, millet, oat, paddy, rice, rye, sorghum, teff, wheat), grain product (broken barley, broken millet, broken sorghum, broken wheat, ground buckwheat, oat flake, polished rice, rice flour, wheat flour), medicine, nut (acorn), other commodities, oilseed (peanut, safflower), oilseed product, pulse (black gram, broad bean, chickpea, cowpea, kidney bean, lentil, moth bean, mung bean, pea, pigeon pea, soybean), pulse product (broken black gram, broken chickpea, broken cowpea, broken kidney bean, broken lentil, broken moth bean, broken mung bean, broken pea, broken pigeon pea, broken soybean, crushed broad bean, other commodities (ground dog biscuit), root (potato), residual commodities, root product, other seed, seasoning, vegetable material (dried parsle), wood and wood product (Hagstrum et al., 2013).

Subfamily: Lytinae Billberg, 1820

Tribe: Lyctini Billberg, 1820

Lyctus africanus Lesne, 1907

Distribution in Iran: Not exactly defined (Hagstrum et al., 2013)

General distribution: Africa, Asia and Middle East, Europe, Ethiopian origin, North America and Mexico, Oceania (Hagstrum et al., 2013).

Host range: Oilseed, root, root product, vegetable material and wood (Hagstrum et al., 2013; Shahhosseini & Kamali, 1989).

Lyctus brunneus (Stephens, 1830)

Distribution in Iran: Tehran and other northern and central provinces (Adeli, 1972; Hedjazi & Soleimani, 1976; Abai, 1984; Niloufari, 1984; Behdad, 1988; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Caspian forests (Niloufari, 1985).

General distribution: Probably Neotropical origin, Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host: Dried fruit, grain, medicine, root, seasoning, vegetable material and wood (Hagstrum et al., 2013; Shahhosseini & Kamali, 1989).

Family: Carabidae Latreille, 1802

Subfamily: Harpalinae Bonelli, 1810

Tribe: Sphodrini Laporte de Castelnau, 1834

Laemostenus terricola (Herbst, 1784)

Distribution in Iran: East Azarbaijan (Shahhosseini & Kamali, 1989).

General distribution: Asia and Middle East, Europe and North America (Hagstrum & Subramanyam, 2009; Shahhosseini & Kamali, 1989).

Host: Animal feed, cereal and currant (Hagstrum & Subramanyam, 2009; Shahhosseini & Kamali, 1989).

Family: Cerambycidae Latreille, 1802

Subfamily: Cerambycinae Latreille, 1802

Tribe: Hesperophanini Mulsant, 1839

Stromatium unicolor (Olivier, 1795)

Distribution in Iran: Gilan, Mazandaran, Gorgan and Tehran (Afshar, 1944; Picard, 1949; Mirzayans, 1950; Farahbakhsh, 1961; Abai, 1969, 1984; Bagheri Zenouz, 1986; Behdad, 1988), Caspian region up to 1800 m altitude and parts of Azarbaijan (Soleimani & Knopf, 1976).

General distribution: Caucasus, the Black Sea coast, Crimea and the Ukraine (Gussakovskii, 1949; Plavilstshikov, 1940).

Host range: Oriental plane, ash, elm, alder, teil, beech, willow, chestnut, poplar, pyrus, wych elm and other hard wood fruit trees and dried wood of these plants Tehran (Afshar, 1944; Picard, 1949; Mirzayans, 1950; Farahbakhsh, 1961; Abai, 1969, 1984; Bagheri-Zenouz, 1986; Behdad, 1988).

Family: Chrysomelidae Latreille, 1802

Subfamily: Bruchinea Latreille, 1802

Tribe: Acanthoscelidini Bridwell, 1946

Acanthoscelides obtectus (Say, 1831)

Distribution in Iran: Khorasan-e Razavi (Akbari Asl et al., 2009), generally distributed (Balachowsky, 1969; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989), Iran (no locality cited) (Bagheri-Zenous, 2013).

General distribution: South American (origin). Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Beverage crops, grain, grain product, pulse (broad bean, broad bean crushed, chickling vetch, chickpea, common vetch, field pea, garden pea, kidney bean, lentil, lupine, scarlet runner bean, soybean, tepary bean) and other seed (Shahhosseini & Kamali, 1989; Hagstrum et al., 2013).

Bruchidius cisti (Fabricius, 1775)

Distribution in Iran: Tehran (Borowiec, 1987 as *Bruchidius canus*) (Ghahari & Borowiec, 2017).

General distribution: Europe, Cyprus, Iran and Turkey (Ghahari & Borowiec, 2017).

Host range: Different cereals (Shahhosseini & Kamali, 1989).

Callosobruchus analis (Fabricius, 1781)

Distribution in Iran: Iran (no locality cited) (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, Oceania and South America (Ghahari & Borowiec, 2017).

Host range: Pulse (black gram, chickpea, cowpea, garden pea, moth bean, mung bean, pea, pigeonpea) and seasoning (Shahhosseini & Kamali, 1989; Hagstrum et al., 2013).

***Callosobruchus chinensis* (Linnaeus, 1758)**

Distribution in Iran: East Azarbaijan, Fars, Guilan, Isfahan, Razavi Khorasan, Khuzestan, Mazandaran, Tehran, Zanjan, central and southern provinces (Alavi, 1974; Frahbakhsh, 1961; Balachowsky, 1969; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Iran (no locality cited) (Decelle & Lodos, 1989; Anton, 1998).

General distribution: African origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Grain (wheat), nut, pulse (black gram, broad bean, catjang, chickling vetch, chickpea, cluster bean, cowpea, garden pea, gram, grass pea, hyacinth bean, kidney bean, lentil, moth bean, mung bean, pea, pigeonpea, ricebean, soybean, sword bean), pulse product, other seed (gourd, okra) and seasoning (Shahhosseini & Kamali, 1989; Hagstrum et al., 2013).

***Callosobruchus maculatus* (Fabricius, 1775)**

Distribution in Iran: Golestan (Khormali et al., 2002), Razavi Khorasan (Akbari Asl et al., 2009), Tehran (Modarres Awal, 1997 as *C. quadrimaculatus*; Anton, 1998, Kazemi et al., 2009), Iran (no locality cited) (Decelle & Lodos, 1989; Bagheri-Zenous, 2013), Generally distributed (Alavi, 1974; Balachowsky, 1969; Chokouhian, 1973; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Taheri, 1991; Hagstrum et al., 2013).

General distribution: African origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal invertebrate, dried fruit, grain, nut, oilseed (safflower), pulse (bambara groundnut, black gram, broad bean, chickpea, cowpea, grass pea, hyaciath bean, kidney bean, lentil, mung bean, pea, pigeonpea, soybean), pulse product, residual commodities, other seed, seasoning and vegetable material (Alavi, 1974; Balachowsky, 1969; Chokouhian, 1973; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Taheri, 1991; Hagstrum et al., 2013).

Tribe: Amblicerini Bridwell, 1932***Zabrotes subfasciatus* (Boheman, 1833)**

Distribution in Iran: Iran (no locality cited) (Anton, 1998, 2010)

General distribution: Central American, South American or West Indian origin, Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico and South America (Hagstrum et al., 2013).

Host range: Grain and pulse (cowpea, kidney bean) (Hagstrum et al., 2013).

Tribe: Bruchini Latreille, 1802***Bruchus atomarius* (Linnaeus, 1760)**

Distribution in Iran: southern provinces (Balachowsky, 1969; Modarres Awal, 1997), Iran (no locality cited) (Borowiec, 1983, 1987; Decelle & Lodos 1989; Anton, 1998).

General distribution: Africa, Asia and Middle East and Europe (Hagstrum et al., 2013).

Host range: Pulse (Hagstrum et al., 2013).

***Bruchus brachialis* Fahraeus, 1839**

Distribution in Iran: Iran (no locality cited) (Modarres Awal, 1997; Ghahari & Borowiec, 2017).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico (Hagstrum et al., 2013).

Host range: Pulse (Hagstrum et al., 2013).

***Bruchus dentipes* Baudi de Selve, 1886**

Distribution in Iran: Tehran (Balachowsky, 1969; Borowiec, 1987; Anton, 1998), East Azarbaijan (Farahbakhsh, 1961, Modarres Awal, 1997 as *B. dentipes ochreosignatus*).

General distribution: Eastern part of Mediterranean basin, Central Asia (Ghahari & Borowiec, 2017).

Host range: Bean, broad bean and other stored legumes, *Vicia* sp. (Fabaceae) (Farahbakhsh, 1961; Balachowsky, 1969; Modarres Awal, 1997).

***Bruchus emarginatus* Allard, 1868**

Distribution in Iran: East Azarbaijan (Nikdel et al., 2016), northwestern provinces (Modarres Awal, 1997; Shahhosseini & Kamali, 1989).

General distribution: Mediterranean basin, the Middle East and Central Asia (Ghahari & Borowiec, 2017).

Host range: Cowpea, pea and other legumes (Modarres Awal, 1997), *Vicia michauxii* Sprengel (Fabaceae) (Nikdel et al., 2016; Shahhosseini & Kamali, 1989).

***Bruchus ervi* Froelich, 1799**

Distribution in Iran: East Azarbaijan, Khuzestan (Hoffmann, 1945a; Alavi, 1974; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe and South America (Hagstrum et al., 2013).

Host range: Pulse (Hagstrum et al., 2013), lentil (Hoffmann, 1945a; Alavi, 1974; Modarres Awal, 1997).

***Bruchus lentis* Frolich, 1799**

Distribution in Iran: Kuhgiloyeh and Boyerahmad, Tehran (Anton, 1998; Khoobdel et al., 2011), Razavi Khorasan (Ghahari et al., 2010), Iran (no locality cited) (Decelle & Lodos 1989; Anton, 2010; Bagheri-Zenous, 2013), generalily distributed (Davatchi, 1946; Frahbakhsh, 1961; Balachowsky, 1969; Alavi, 1974; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico (Hagstrum et al., 2013).

Host range: Pulse (Hagstrum et al., 2013), lentil (Davatchi, 1946; Frahbakhsh, 1961; Balachowsky, 1969; Alavi, 1974; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989).

***Bruchus pisorum* (Linnaeus, 1758)**

Distribution in Iran: Golestan (Khormali et al., 2002), Tehran (Borowiec, 1987), generally distributed in Iran (Davatchi, 1946; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Behdad, 1982, 1988; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: West Asian origin, Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Grain, pulse and vegetable material (Hagstrum et al., 2013), Lentil (Davatchi, 1946; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Behdad, 1982, 1988; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Bruchus rufimanus Boheman, 1833

Distribution in Iran: Tehran (Borowiec, 1987; Borowiec & Anton, 1993; Anton 1998; Khoobdel et al., 2011), generally distributed (Davatchi, 1946; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Behdad, 1982; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Iran (no locality cited) (Bagheri-Zenous, 2013).

General distribution: West Asian origin, Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Grain, medicine, pulse, other seed and seasoning (Shahhosseini & Kamali, 1989; Hagstrum et al., 2013).

Bruchus tristis Boheman, 1833

Distribution in Iran: Kuhgiluyeh & Boyerahmad (Anton, 1998), Western provinces (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: South Europe, North Africa, Caucasian countries, Cyprus, Turkey and the Middle East (Ghahari & Borowiec, 2017).

Host range: Bean, pea (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Family: Cleridae Latreille, 1802

Subfamily: Korynetinae Laport de Castelnau, 1836

Necrobia rufipes (De Gree, 1775)

Distribution in Iran: Hormozgan, Kerman, Sistan & Baluchestan (Brodský & Winkler, 1988), generally distributed (Farahbakhsh, 1961; Modarres Awal, 1997), Golestan (Khormali et al., 2002) and In Iran (Sepasgosarian, 1978).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico and Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate (insect larva), animal vertebrate (dried fish), beverage crops (cocoa), bake goods, dried fruit, fresh fruit or vegetable, grain (maize), grain product, medicine, nut, other commodities, oilseed (palm kernel, peanut), oilseed product (copra), pulse, root, root product, other seed, seasoning, spoiled other commodities, textile and vegetable material (Sepasgosarian, 1978; Hagstrum et al., 2013).

Necrobia violacea (Linnaeus, 1758)

Distribution in Iran: Mazandaran (Brodský & Winkler, 1988), Kermanshah and central provinces (Farahbakhsh, 1961; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia, Middle East, Europe, North America and Mexico (Hagstrum et al., 2013).

Host range: Animal vertebrate and grain (Hagstrum et al., 2013), Animal (skin, fat, egg powder, dried, smoked, meat and fish), cheese, bone, dried fruits, nuts, cocoon of silkworm (in granary) and predator of *Calliptamus italicus*, *Dociostaurus maroccanus* and other grasshopper (Farahbakhsh, 1961; Shahhosseini & Kamali, 1989).

Family: Cryptophagidae Kirby, 1826

Subfamily: Atomariinae LeConte, 1861

Tribe: Atomariini LeConte, 1861

Atomaria (Atomaria) lewisi Reitter, 1877

Distribution in Iran: Iran (no locality cited) (Johnson et al., 2007).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico (Hagstrum et al., 2013).

Host range: Grain and medicine (Hagstrum et al., 2013).

Subfamily: Cryptophaginae Kirby, 1826

Tribe: Cryptophagini, Kirby, 1837

Cryptophagus araxicola Reitter, 1889

Distribution in Iran: East Azarbaijan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Armenia and Azarbaijan (Johnson et al., 2007).

Host range: Fungus and on dried fruits (in damp granaries) (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Cryptophagus cellaris (Scopoli, 1763)

Distribution in Iran: East Azarbaijan, Hamadan, Tehran (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, bake goods, dried fruit, fresh fruit or vegetable, grain, grain product, nut, other commodities, oilseed, oilseed product, pulse, pulse product, residual commodities, spoiled other commodities, vegetable material and wood product (Hagstrum et al., 2013; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Cryptophagus distinguendus Sturm, 1845

Distribution in Iran: East Azarbaijan, Mazandaran (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Europe and North America (Hagstrum & Subramanyam, 2009).

Host range: Dried fruit and residue (Hagstrum & Subramanyam, 2009), fungus and on dried fruits (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Cryptophagus hexagonalis Toumier, 1872

Distribution in Iran: Azarbaijan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Finland, Belarus, Russia, Turkey, Siberia, Mongolia, Kazakhstan, Turkmenistan (Johnson et al., 2007; Otero, 2013).

Host range: Fungus and on dried fruits (in damp granaries) (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Cryptophagus simplex Miller, 1858

Distribution in Iran: East Azarbaijan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Asia and Middle East, Europe (Hagstrum & Subramanyam, 2009).

Host range: Flourmills, granaries (Hagstrum & Subramanyam, 2009), Fungus and on dried fruits (in damp granaries) (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Cryptophagus laticollis P. H. Lucas, 1846

Distribution in Iran: East Azarbaijan, Mazandaran and Tehran (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Europe, North America and South America (Hagstrum & Subramanyam, 2009).

Host range: Almond, apricot (dried), bean, currant, dried fruit, esparto grass (moldy), fig, grain residue, illipe nut, lentil, lotus, maize, mushroom, oat, oilcake, paddy residue, palm kernel, pollard, prune, raisin, rye, seaweed (dried), soybean meal, sultana, valonia spillage, vegetable refuse (decaying), walnut, wheat and wheat flour (Hagstrum & Subramanyam, 2009), Fungus and on dried fruits (in damp granaries) (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Cryptophagus acutangulus Gyllenhal, 1827

Distribution in Iran: East Azarbaijan, Fars and Northern Khorasan (Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico and Oceania (Hagstrum et al., 2013).

Host range: Animal vertebrate, dried fruit, grain, grain product nut, other commodities, pulse product, spoiled other commodities and vegetable material (Hagstrum et al., 2013).

Cryptophagus dentatus (Herbst, 1793)

Distribution in Iran: Fars, Kuhgiluyeh-Boyerahmad and Mazandaran (Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico (Hagstrum et al., 2013).

Host range: Dried fruit, grain, grain product, medicine, oilseed, root product and vegetable material (Hagstrum et al., 2013).

Cryptophagus pilosus Gyllenhal, 1827

Distribution in Iran: Iran (No locality cited) (Hagstrum & Subramanyam, 2009).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico (Hagstrum et al., 2013).

Host range: Dried fruit, grain, grain product, grain spoiled, nut, pulse product and vegetable material (Hagstrum et al., 2013).

Cryptophagus scanicus (Linnaeus, 1758)

Distribution in Iran: Guilan, Mazandaran and Tehran (Modarres Awal, 1997) and Iran (no locality cited) (Majka & Langor, 2010).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico (Hagstrum et al., 2013).

Host range: Dried fruit, fresh fruit or vegetable, grain, grain product, nut and vegetable material (Hagstrum et al., 2013).

Cryptophagus scutellatus Newman, 1834

Distribution in Iran: Mazandaran (Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico (Hagstrum et al., 2013).

Host range: Grain (Hagstrum et al., 2013).

Cryptophagus subfumatus kraatz, 1856

Distribution in Iran: Iran (No locality cited) (Hagstrum & Subramanyam, 2009)

General distribution: Asia and Middle East, Europe, North America and Mexico (Hagstrum et al., 2013).

Host range: Animal invertebrate, dried fruit, grain, nut and seasoning (Hagstrum et al., 2013).

Family: Curculionidae Latreille, 1802

Subfamily: Cossoninae Schönherr, 1825

Tribe: Onycholipini Wollaston, 1837

Hexarthrum exiguum (Boheman, 1838)

Distribution in Iran: Tehran (Sakenin et al., 2018).

General distribution: Not exactly defined (Sakenin et al., 2018).

Host range: Not exactly defined (Sakenin et al., 2018).

Subfamily: Rhynchophorinae Schönherr, 1833

Tribe: Litosomini Lacordaire, 1866

Sitophilus granarius (Linnaeus, 1758)

Distribution in Iran: Generally distributed (Afshar, 1944; Hoffmann, 1945b; Dvatchi, 1947; Shayesteh, 1957; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Guilan, Khuzestan (Legalov et al., 2010).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal vertebrate, beverage crops (cocoa bean, coffee bean), dried fruit, fresh fruit or vegetable, grain (barley, buckwheat, maize, oat, paddy, rice, rye, sorghum, triticale, wheat), grain product (noodle, wheat bran, flour or meal), medicine, nut (acorn, almond,

chestnut), other commodities, oilseed (peanut, safflower, oilseed product, pulse (broad bean, lupine, pea, soybean), pulse product (crushed broad bean, soybean meal), root, residual commodities, root product, other seed, textile and vegetable material (alfalfa pellet, rye grass pellet) (Afshar, 1944; Hoffmann, 1945b; Dvatchi 1947; Shayesteh, 1957; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Modarres Awal, 1997; Hagstrum et al., 2013).

Sitophilus oryzae (Linnaeus, 1763)

Distribution in Iran: Generally distributed (Afshar, 1944; Hoffmann, 1945b; Dvatchi 1947; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Habibian, 1987; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Fars, Golestan, Guilan, Hamadan, Isfahan, Khorasan, Semnan (Legalov et al., 2010).

General distribution: Oriental origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, beverage crops, bake goods, dried fruit, fresh fruit or vegetable, grain (barley, buckwheat, maize, oat, rice, sorghum, teff, triticale, wheat), grain product, medicine, nut acorn, chestnut, other commodities, oilseed (safflower), oilseed product, pulse (broad bean), pulse product (crushed broad bean, split pea), root, residual commodities, root product, other seed, seasoning, textile and vegetable material (tamarind) (Afshar, 1944; Hoffmann, 1945b; Dvatchi, 1947; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Habibian, 1987; Shahhosseini & Kamali, 1989; Modarres Awal, 1997; Hagstrum et al., 2013).

Sitophilus zeamais Motschulsky, 1855

Distribution in Iran: Tehran and Balouchestan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Fars, Golestan, Isfahan, Mazandaran, Sistan & Baluchestan, Tehran (Legalov et al., 2010).

General distribution: Oriental origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, beverage crops 2.8%, bake goods, dried fruit, dried fruit product, fresh fruit or vegetable (melon), grain (barley, buckwheat, maize, millet, oat, rice, sorghum, triticale, wheat, grain product (ground sorghum), medicine, nut (acorn, chestnut), nut product, other commodities, oilseed (peanut), oilseed product, pulse (broad bean, bambara groundnut, cowpea, soybean), pulse product (crushed broad bean), root, root product (cassava chip, yam chip), other seed, seasoning (pepper) and vegetable material (Hagstrum et al., 2013), corn (in granary) (Modarres Awal, 1997; Shahhosseini & Kamali, 1989).

Subfamily: Scolytinae Latreille, 1804

Tribe: Cryphalini Lindemann, 1876

Hypothenemus eruditus Westwood, 1836

Distribution in Iran: Northern provinces (Abai, 1984; Behdad, 1988; Modarres Awal, 1997), Guilan, Tehran, and other northern provinces (Amini et al., 2013; Modarres Awal, 1997 (as

Hypothenemus lezhavai Pjatnitskii, 1929 and *Hypothenemus aspericollis* Wollaston, 1860 (see Ghahari & Colonelli, 2012)), Iran (no locality cited) (Balachowsky, 1949; Mifsud & Knížek, 2009; Knížek, 2011).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Beverage crops, grain, other commodities, pulse, root, root product and vegetable material (Hagstrum et al., 2013), tree of heaven, apricot, almond (Abai, 1984; Behdad, 1988; Modarres Awal, 1997).

Hypothenemus hampei (Ferrari, 1867)

Distribution in Iran: Iran (No locality cited) (Hagstrum & Subramanyam, 2009; Knížek, 2011).

General distribution: Central African origin. Africa, Asia and Middle East, Central America and Caribbean, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Beverage crops, grain, nut, other commodities, pulse, vegetable material (Hagstrum et al., 2013).

Family: Dermestidae Latreille, 1804

Subfamily: Attageninae Laporte, 1840

Tribe: Attagenini Laporte, 1840

Attagenus bifasciatus (Oliver, 1790)

Distribution in Iran: No locality cited (Mroczkowski, 1968; Háva, 2003a, 2007, 2015; Tezcan et al., 2004).

General distribution: Africa, Asia and Middle East and Europe (Hagstrum et al., 2013).

Host range: Shops, residences (Hagstrum et al., 2013)

Attagenus fasciatus (Thunberg, 1795)

Distribution in Iran: Tehran (Modarres Awal, 1997), no locality cited (Mroczkowski, 1968; Zhantiev, 1976; Háva, 2007, 2015).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate, animal vertebrate (dried milk), beverage crops, grain, grain product, grain spoiled, medicine, other commodities, oilseed, oilseed product, pulse, pulse product, other seed and seasoning, textile (Hagstrum et al., 2013).

Attagenus lobatus Rosenhauer, 1856

Distribution in Iran: Khuzestan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997), no locality cited (Mroczkowski, 1968; Zhantiev, 1976; Háva, 2003b, 2007, 2015).

General distribution: Africa, Asia and Middle East, Europe and North America and Mexico (Hagstrum et al., 2013).

Host range: Animal vertebrate and grain (Hagstrum et al., 2013), storage products (Shahhosseini & Kamali, 1989).

Attagenus pellio (Linnaeus, 1758)

Distribution in Iran: Alborz (Farahbakhsh, 1961), Tehran (Modarres Awal, 1997), no locality cited (Zhantiev, 1976; Abivardi, 2001; Háva, 2007, 2015).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, grain, grain product soybean meal, other commodities, oilseed, oilseed product, residual commodities, textile, vegetable material (Hagstrum et al., 2013; Farahbakhsh, 1961).

Attagenus unicolor (Brahm, 1791)

Distribution in Iran: Alborz (Farahbakhsh, 1961), Khorasan, Tehran and other areas (Modarres Awal, 1997), no locality cited (Abivardi, 2001; Háva, 2007, 2015).

General distribution: Oriental region origin. Africa, Asia and Middle East, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, beverage crops, bake goods, dried fruit, grain, grain product, medicine, nut, nut product, other commodities, oilseed, oilseed product, pulse, pulse product, residual commodities, other seed, seasoning, textile and vegetable material (Hagstrum et al., 2013).

Subfamily: Dermestinae Latreille, 1807**Tribe: Dermestini Latreille, 1807***Dermestes ater* De Gree, 1774

Distribution in Iran: Hormozgan (Bagh-e Tang), Sistan and Baluchestan (Chabahar) (Háva & Kalík, 1999), no locality cited (Mroczkowsky, 1968; Zhantiev, 1976; Háva, 2007).

General distribution: American origin. Africa, Central America and Caribbean, Europe, North America and Oceania (Hagstrum & Subramanyam, 2009).

Host range: Animal feed, animal product, bacon, basket ware, bone, Brazil nut, Chinese traditional medicine, cocoa, cotton seed, dried fish, fertilizer, fish air bladder (dried), fishmeal, grain residue, groundnut cake, hide, meat (dried), museum specimen, nut, oilcake, oyster (dried), sesame seed, shark fin (dried), tobacco and turtle shell (Hagstrum & Subramanyam, 2009).

Dermestes frischii Kugelann, 1792

Distribution in Iran: Ardabil (Khalkhal, Kivi Bala), East Azarbaijan (Mianeh, Tabriz), Fars (Háva & Kalík, 1999; Keshavarzi et al., 2015), Isfahan (Robot Tork), Kuhgiluyeh and Boyerahmad (Kooch-e-Dinar), Mazandaran (Alamdeh, Chalus, Kojur, Veresk), Sistan and Baluchestan (Bampur, Chabahar), Tehran (Darakeh, Evin, Golhak) (Háva & Kalík, 1999), generally distributed (Farahbakhsh, 1961; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), no locality cited (Abivardi, 2001; Háva, 2007) .

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate (fishmeal), beverage crops, bake goods, dried fruit, grain, grain product (wheat flour, wheat germ), medicine, nut, other commodities, oilseed (peanut), oilseed product, pulse, pulse product (soybean, meal),

seasoning, wood product, chocolate, cacao, bone and cheese (in the granary and ship) (Farahbakhsh, 1961; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Modarres Awal, 1997; Hagstrum et al., 2013).

Dermestes lardarius Linnaeus, 1758

Distribution in Iran: Tehran, West Azarbaijan (Khoy) (Háva & Kalík, 1999), no locality cited (Háva, 2007).

General distribution: Eurasian origin, Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate, animal vertebrate (fishmeal), beverage crops (cocoa), bake goods, dried fruit, grain, grain product (wheat flour, wheat germ), nut (almond), other commodities, oilseed (peanut), oilseed product copra, pulse, pulse product (soybean meal), other seed, seasoning, spoiled other commodities and vegetable material (Hagstrum et al., 2013).

Dermestes maculatus De Geer, 1774

Distribution in Iran: Tehran, West Azarbaijan (Khoy) (Háva & Kalík, 1999), no locality cited (Háva, 2007).

General distribution: Eurasian origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate, animal vertebrate (blood meal, bone meal, fishmeal, fish (dried), sardine meal, hide), beverage crops (cocoa), bake goods, fresh fruit or vegetable, grain, grain product (wheat flour, wheat germ, medicine, nut (almond)), other commodities (dog chow), oilseed (peanut), oilseed product (copra, palm kernel meal), pulse, pulse product (soybean meal), root, other seed, seasoning, textile, vegetable material, wood product and vegetable material (alfalfa pellet, ryegrass pellet) (Hagstrum et al., 2013).

Dermestes murinus Linnaeus, 1758

Distribution in Iran: Guilan (Assalem), Mazandaran (Háva & Kalík, 1999), no locality cited (Háva, 2007).

General distribution: Africa, Asia and Middle East and Europe (Hagstrum et al., 2013).

Host range: Animal invertebrate (silk cloth), animal vertebrate (hog bristle, woolen) (Hagstrum et al., 2013).

Dermestes mustellinus Erichson, 1846

Distribution in Iran: East Azarbaijan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Europe, Asia (Hagstrum & Subramanyam, 2009).

Host range: Strong product (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Subfamily: Megatominae Leach, 1815

Tribe: Anthrenini Gistel, 1856

Anthrenus flavipes LeConte, 1854

Distribution in Iran: No locality cited (Háva, 2007; Herrmann et al., 2010), East Azarbaijan (Alavi, 1974; Khorramshahi, 1982).

General distribution: Oriental Region origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico and Oceania (Hagstrum et al., 2013).

Host range: Animal invertebrate (silk cloth), animal vertebrate (hog bristle, woolen), fresh fruit or vegetable, grain, grain product, other commodities, oilseed product, other seed, textile (cotton cloth), vegetable material and wood (Hagstrum et al., 2013), Goat wool (Alavi, 1974; Khorramshahi, 1982).

Anthrenus museorum (Linnaeus, 1761)

Distribution in Iran: Generally distributed (Farahbakhsh, 1961; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), no locality cited (Háva, 2007; Herrmann et al., 2010).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania (Hagstrum & Subramanyam, 2009).

Host range: Apricot (dried), carpet, cereal, currant, feather, fig, fur, hide, insect (dried), museum specimen, rice, silk, sorghum, sultana, upholstery, wheat flour, wool and Predator gypsy moth eggs, herbarium (Farahbakhsh, 1961; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Modarres Awal, 1997; Hagstrum & Subramanyam, 2009).

Anthrenus pimpinellae (Fabricius, 1775)

Distribution in Iran: Generally distributed (Farahbakhsh, 1961; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), no locality cited (Herrmann et al., 2010).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, other commodities (Shahhosseini & Kamali, 1989; Hagstrum et al., 2013).

Anthrenus scrophulariae (Linnaeus, 1758)

Distribution in Iran: Generally distributed (Farahbakhsh, 1961; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), no locality cited (Háva, 2007; Herrmann et al., 2010).

General distribution: Eurasian origin. Africa, Asia and Middle East, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, grain, grain product, other commodities, other seed, textile and vegetable material (Shahhosseini & Kamali, 1989; Hagstrum et al., 2013).

Anthrenus verbasci (Linnaeus, 1767)

Distribution in Iran: Fars (Ardakan, Shiraz) (Karami et al., 2004), Isfahan (Bagheri & Nasr Isfahani, 2011), Tehran (Shahhosseini & Kamali, 1989; Modarres Awal, 1997), no locality cited (Háva, 2007; Herrmann et al., 2010).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, beverage crops, bake goods, dried fruit, grain, grain product, grain spoiled, medicine, nut, nut product, other commodities, oilseed, oilseed product, pulse, pulse product, root, other seed, seasoning, textile, vegetable material and wood product (Hagstrum et al., 2013), carpet and storage product (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Subfamily: Megatomininae Leach, 1815

Tribe: Megatomini Leach, 1815

Phradonoma nobile (Reitter, 1881)

Distribution in Iran: Hormozgan (Isin), Kerman (Nosrat-Abad, Rafsanjan), Khuzestan (Ahwaz) (Háva, 2006), Sistan and Baluchestan (Nikshahr) (Herrmann et al., 2010, on flowers of *C. aurantifolia*), no locality cited (Modarres Awal, 1997; Háva, 2003b, 2007, 2015; Herrmann & Háva, 2008), in Iran (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East and Europe (Hagstrum et al., 2013).

Host range: Animal vertebrate, beverage crops, dried fruit, grain, grain product, medicine, oilseed and oilseed product (Hagstrum et al., 2013), storage product (Shahhosseini & Kamali, 1989).

Trogoderma glabrum (Herbst, 1783)

Distribution in Iran: Iran (No locality cited) (Háva, 2007; Bulak et al., 2013).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico (Hagstrum et al., 2013).

Host range: Animal invertebrate (dried insect, pollen), animal vertebrate (fishmeal, dried milk), dried fruit (raisin), grain (barley, maize, oat, wheat), grain product (cracked maize, maize meal, noodle, oatmeal, polished rice, rolled oat, rolled barley, rye, sorghum, wheat flour, wheat flour with germ, wheat germ), nut (almond, walnut), other commodities (cat food, egg noodle, kibbled dog food, poultry laying mash), oilseed (peanut, safflower), oilseed product (broken safflower, safflower meal), pulse (cowpea, kidney bean, lima bean), pulse product and other seed (Hagstrum et al., 2013).

Trogoderma granarium Everts, 1898

Distribution in Iran: Golestan (Khormali et al., 2002), Semnan (Forghani & Marouf, 2015), Tehran (Mehrabadi et al., 2011), generally distributed (Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Taheri, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), no locality cited (Abivardi, 2001; Háva, 2007).

General distribution: Indian origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal vertebrate (beef blood, dried milk), beverage crops (coffee bean), bake goods (cracker), dried fruit (apricot, date, prune, raisin), fresh fruit or vegetable (peach), grain (barley, maize, oat, rice, rye, shelled maize, wheat), grain product (barley cereal, cracked oat, cracked rice, ground barley, ground buckwheat, ground maize, ground rice, instant white rice, maize cereal, maize meal, malt, noodle, oatmeal, polished rice, precooked oatmeal, rice flake, rolled barley, rolled oat, wheat feed, wheat flour, germ or semolina), medicine, nut, nut product (almond meat, kola nut, pecan meat, walnut meat), other

commodities (egg noodle, ground dog food, gelatin, halvah, pablum, oilseed (flax, peanut, sesame seed), oilseed product (copra meal, cottonseed, ground flax), pulse (black gram, chickpea, cowpea, hyacinth bean, lentil, lima bean, mung bean, pigeonpea, pinto bean, soybean), pulse product (cracked pea, lentil flour, soybean meal), root, root product (tapioca), other seed (alfalfa, grass), seasoning, textile, vegetable material (herb, dried mushroom, yeast and wood product (cork , paper) (Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Taheri, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997; Hagstrum et al., 2013).

Trogoderma inclusum Le Conte, 1854

Distribution in Iran: East Azarbaijan, Kerman, Kermanshah, Khuzestan, Semnan, Tehran (Shahhosseini & Kamali, 1989; Modarres Awal, 1997), no locality cited (Háva, 2004, 2007; Háva & Herrmann, 2011).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal invertebrate (dried insect, pollen), animal vertebrate (fishmeal, dried milk), bake goods, dried fruit (raisin), dried fruit product, grain (barley, maize, oat, rye, sorghum, wheat), grain product (maize meal, malt, noodle, oatmeal, polished rice, rolled barley, rolled oat, wheat flour or germ), nut (almond, walnut), other commodities (cat food, egg noodle, kibbled dog food, poultry laying mash), oilseed (peanut, safflower), oilseed product (broken safflower, safflower meal), pulse (cowpea, kidney bean, lima bean), pulse product, residual commodities, root product, other seed, seasoning, textile and vegetable material (Shahhosseini & Kamali, 1989; Hagstrum et al., 2013).

Trogoderma megatomoides Reitter, 1881

Distribution in Iran: East Azarbaijan, Fars, Khorasan, Mazandaran, Tehran (Modarres Awal, 1997, as *Entomotrogus megatomoides*), no locality cited (Abivardi, 2001; Háva, 2007, 2015).

General distribution: Africa, Central America, Europe, North America and Mexico (Hagstrum & Subramanyam, 2009).

Host range: Insect (dried) (Hagstrum & Subramanyam, 2009).

Trogoderma variabile Ballion, 1878

Distribution in Iran: Semnan (Forghani & Marouf, 2015), no locality cited (Somerfield, 1981; Stejskal et al., 2005; Háva, 2007; Bulak et al., 2013).

General distribution: Central Asia and Middle Eastian origin, Africa, Asia and Middle East, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal invertebrate (dried insect, pollen), animal vertebrate (dried milk, fishmeal), beverage crops, bake goods, dried fruit (raisin), grain (barley, oat, maize, rice, rye, shelled maize, sorghum, wheat), grain product (barley cereal, instant oatmeal, maize cereal, maize meal, noodle, oatmeal, polished rice, rolled barley, rolled oat, wheat feed, wheat flour, wheat germ, white rice), medicine, nut, nut product, other commodities (cat food, kibbled or ground dog food, egg noodle, flax, gelatin, pablum, poultry laying mash), oilseed (safflower), oilseed product (copra meal, safflower meal), pulse (cowpea, kidney bean, lima bean, pinto bean), root, root product, other seed, seasoning, textile and vegetable material (Hagstrum et al., 2013).

Trogoderma versicolor (Creutzer, 1799)

Distribution in Iran: Generally distributed (Farahbakhsh, 1961; Sepasgosarian, 1978; Modarres Awal, 1997; Shahhosseini & Kamali, 1989), no locality cited (Abivardi, 2001; Háva, 2007).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, dried fruit, fresh fruit or vegetable, grain (maize, wheat), grain product (oatmeal, wheat flour), nut, oilseed, oilseed product, pulse, root, other seed and seasoning (Hagstrum et al., 2013), entomological collections, legumes grains, dried fish (Farahbakhsh, 1961; Sepasgosarian, 1978; Modarres Awal, 1997; Shahhosseini & Kamali, 1989).

Subfamily: Thorictinae Agassiz, 1846

Tribe: Thaumaphrastini Anderson, 1949

Thorictodes heydeni Reitter, 1875

Distribution in Iran: Bandar Abbas (Shahhosseini & Kamali, 1989), not exactly defined (Hagstrum et al., 2013).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico and Oceania (Hagstrum et al., 2013).

Host range: Animal vertebrate, grain, grain product, medicine, other commodities, oilseed, oilseed product, pulse, pulse product, root, residual commodities, root product and other seed (Hagstrum et al., 2013).

Family: Erotylidae Latreille, 1802

Subfamily: Cryptophilinae Casey, 1900

Tribe: Cryptophilini Casey, 1900

Cryptophilus integer (Heer, 1841)

Distribution in Iran: Iran (No locality cited) (Hagstrum & Subramanyam, 2009).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Dried fruit, grain, grain product, medicine, pulse product, residual commodities, spoiled other commodities, vegetable material (Hagstrum et al., 2013).

Family: Laemophloeidae Ganglbauer, 1899

Subfamily: Laemophloeinae Ganglbauer, 1899

Cryptolestes ferrugineus (Stephens, 1831)

Distribution in Iran: Tehran (Farahbakhsh 1961 as *Laemophloeus ferrugineus* (Stephens, 1828)), generally distributed (Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Iran (no locality cited) (Wegrzynowicz, 2007).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal vertebrate, beverage crops, dried fruit, fresh fruit or vegetable, grain (barley, maize, millet, oat, rice, rye, wheat), grain product (cracked barley, ground barley,

ground maize, ground millet, ground oat, ground rice, ground rye, ground wheat), medicine, nut, other commodities, oilseed (canola, flax, rape, sunflower, oilseed) product (ground flax, ground rape, ground sunflower), pulse (broad bean, soybean), pulse product, crushed broad bean, soybean meal, root, residual commodities, root product, other seed (clover, ground clover, mustard), seasoning, vegetable material (alfalfa pellet, rye grass pellet) (Hagstrum et al., 2013), Adults and larvae feed on cake and seeds (Farahbakhsh, 1961); on cotton, wheat, in granary (Farahbakhsh 1961; Modarres Awal, 1997; Shahhosseini & Kamali, 1989).

Cryptolestes pusillus (Schönherr, 1817)

Distribution in Iran: Isfahan, Tehran, Tabriz, Rasht (Farahbakhsh 1961 as *Laemophloeus pusillus* (Schönherr, 1817)); (Shahhosseini & Kamali, 1989; Davatchi, 1950; Modarres Awal, 1997 as *Laemophloeus minutus* (Olivier, 1791)).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Beverage crops, bake goods, dried fruit, dried fruit product, fresh fruit or vegetable, grain (maize, rice, teff, wheat), grain product (maize meal, rice meal, wheat feed, wheat flour, wheat meal), medicine, nut, other commodities, oilseed (peanut, safflower), oilseed product (peanut meal, safflower meal), pulse (cowpea, pulse product (cowpea meal, lentil flour, soybean meal), root, root product, other seed, seasoning and vegetable material (Hagstrum et al., 2013).

Cryptolestes turcicus (Grouvelle, 1876)

Distribution in Iran: Tehran (Farahbakhsh, 1961 as *Laemophloeus turcicus* Grouvelle, 1876), generally distributed (Davatchi, 1950; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Iran (no locality cited) (Wegrzynowicz, 2007).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, South America (Hagstrum et al., 2013).

Host range: Beverage crops, dried fruit, grain (maize, rice, wheat), grain product (cracked barley, cracked maize, cracked oat, cracked rice, cracked wheat, maize meal, rice meal, wheat meal), medicine, oilseed (peanut), oilseed product (cracked peanut, peanut meal), pulse (broad bean, cowpea), pulse product (cowpea meal, cracked cowpea, crushed broad bean, soybean meal), residual commodities, root product, seasoning, vegetable material (Hagstrum et al., 2013), Cacao, cotton, rice, wheat and other cereals and their products (in granary) (Davatchi, 1950; Sepasgosarian, 1978; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Family: Lathridiidae Erichson, 1842

Subfamily: Corticariinae, Erichson, 1842

Corticaria serrata (Paykull, 1798)

Distribution in Iran: In Iran (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia, Europe, North America and Oceania (Hagstrum & Subramanyam, 2009).

Host range: Beverage crops, dried fruit, dried fruit product, grain, grain product, nut, oilseed product, pulse and wood product (Hagstrum et al., 2013), storage products (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Family: Melyridae Leach, 1815

Subfamily: Melyrinae Leach, 1815

Tribe: Melyrini Leach, 1815

Melyris oblonga (Fabricius, 1775)

Distribution in Iran: Tehran (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East and Europe (Hagstrum et al., 2013).

Host range: Animal invertebrate, dried fruit, dried fruit product, grain, nut, pulse and root (Hagstrum et al., 2013), dry wood (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Family: Mycetophagidae Leach, 1815

Subfamily: Mycetophaginae Leach, 1815

Tribe: Typhaeini Nikitsky, 1993

Typhaea stercorea (Linnaeus, 1758)

Distribution in Iran: Mazandaran, Gilan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal vertebrate, beverage crops, dried fruit, fresh fruit or vegetable, grain, grain product (kibbled wheat, medicine), nut, nut product, oilseed, oilseed product, pulse, pulse product (soybean meal), root, residual commodities, root product, other seed, seasoning, spoiled other commodities, vegetable material (Hagstrum et al., 2013), fungus on raisin, date and dried apricot (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Family: Nitidulidae Latreille, 1802

Subfamily: Carpophilinae Erichson, 1842

Tribe: Carpophilini Erichson, 1842

Carpophilus dimidiatus (Fabricius, 1792)

Distribution in Iran: Khuzestan (Azimi et al., 1993), generally distributed in Iran (Esmaili, 1983; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal vertebrate, beverage crops, dried fruit, dried fruit product, fresh fruit or vegetable, grain, grain product (cracked rough rice, wheat bran), medicine, nut, nut product, other commodities, oilseed, oilseed product, pulse, pulse product, root, residual commodities, root product, other seed, seasoning, spoiled other commodities and vegetable material (Hagstrum et al., 2013), citrus and also rice and fruits (Esmaili, 1983; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Carpophilus hemipterus (Linnaeus, 1758)

Distribution in Iran: generally distributed in Iran (Farahbakhsh, 1961; Alavi, 1974; Khorramshahi, 1982; Esmaili, 1983; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Golestan (Khormali et al., 2002).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, beverage crops, bake goods, dried fruit, dried fruit product, fresh fruit or vegetable, grain, grain product, grain spoiled, medicine, nut, nut product, other commodities, oilseed, oilseed product, pulse, pulse product, root, residual commodities, root product, other seed, seasoning, spoiled other commodities and vegetable material (Hagstrum et al., 2013), date-palm, dried apricot, almond, citrus and other dried fruits and agricultural products (in granary) (Farahbakhsh, 1961; Alavi, 1974; Khorramshahi, 1982; Esmaili, 1983; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Carpophilus mutilatus Erichson, 1843

Distribution in Iran: Golestan (Khormali et al., 2002).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate, beverage crops, dried fruit, fresh fruit or vegetable, grain, grain product, grain spoiled, nut, nut product, oilseed, oilseed product, pulse, root, residual commodities, root product, seasoning and vegetable material, flour and spaghetti (Hagstrum et al., 2013; Khormali et al., 2002).

Carpophilus obsoletus Erichson, 1843

Distribution in Iran: Khuzestan, Markazi (Shahhosseini & Kamali, 1989; Naeem & Akhyani, 1988; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Beverage crops, dried fruit, fresh fruit or vegetable, grain, grain product, medicine, nut, oilseed, oilseed product, pulse, root, residual commodities, root product, seasoning, and vegetable material (Hagstrum et al., 2013), fig and also date, dried apricot and raising (Shahhosseini & Kamali, 1989; Naeem & Akhyani, 1988; Modarres Awal, 1997).

Family: Ptinidae Latreille, 1802

Subfamily: Anobiinae Fleming, 1821

Tribe: Anobiini Fleming, 1821

Anobium punctatum (De Geer, 1774)

Distribution in Iran: Tehran, Gilan, central and northern provinces and probably other areas (Afshar, 1944; Farahbakhsh, 1961; Niloufari, 1984; Abai, 1984; Behdad, 1988; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Golestan (Abd-Rabou et al., 2005), Guilan, northern and central provinces (Modarres Awal, 1997), Tehran (Modarres Awal, 1997 as *Anobium domesticum* (Geoffroy, 1785)), Iran (no locality cited) (Zahradnik, 2007).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico and Oceania (Hagstrum et al., 2013).

Host range: Other commodities, root, other seed, wood (Hagstrum et al., 2013), pine, poplar, alder, oak, ash, walnut and other hard wood (Afshar, 1944; Farahbakhsh, 1961; Niloufari, 1984; Abai, 1984; Behdad, 1988; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Tribe: Stegobiini White, 1982

Stegobium paniceum (Linnaeus, 1758)

Distribution in Iran: Largely distributed, no locality cited (Davatchi, 1947; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Niloufari, 1984; Shahhosseini & Kamali, 1989; Modarres Awal, 1997, Zahradvik, 2007).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate, animal vertebrate (meat meal, dried milk), beverage crops, bake goods, dried fruit, fresh fruit or vegetable, grain (barley, maize), grain product (oatmeal flour, polished rice, wheatfeed, wheat flour), medicine, nut, nut product, other commodities, oilseed (safflower), oilseed product (safflower meal), pulse, pulse product (ground pea, lentil flour, pea flour, soybean meal), root, root product, other seed (pomegranate, poppy), seasoning (aniseed, black mustard, caraway, cardamom, chili pepper powder, coriander, cumin, fennel powder, garlic, ginger, turmeric), textile, vegetable material (alfalfa pellets, onion, rye grass pellet) wood, wood product (Davatchi, 1947; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Niloufari, 1984; Shahhosseini & Kamali, 1989; Modarres Awal, 1997; Hagstrum et al., 2013).

Subfamily: Ptininae Latreille, 1802

Tribe: Ptinini Latreille, 1802

Niptus hololeucus (Faldermann, 1835)

Distribution in Iran: General distributed (Davatchi, 1947; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Bagheri Zenouz, 1986; Khorramshahi, 1982; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Fars (Dashan et al., 2014), generally distributed (Farahbakhsh, 1961; Modarres Awal, 1997), Iran (no locality cited) (Borowski, 2007).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania (Hagstrum et al., 2013).

Host range: Animal invertebrate, animal vertebrate (fishmeal), beverage crops, bake goods, fresh fruit or vegetable, grain, grain product (wheat flour, wheatfeed), medicine, other commodities, oilseed product, pulse product, root, other seed, seasoning, textile, vegetable material and wood product (Hagstrum et al., 2013), wool, skin, silk, linen, paper, flour, carpet, cotton and related products, tobacco, tea, medicinal plants, rice and other cereals, feather (Davatchi, 1947; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Bagheri Zenouz, 1986; Khorramshahi, 1982; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

***Ptinus (Ptinus) fur* (Linnaeus, 1758)**

Distribution in Iran: Northern and central provinces (Afshar, 1944; Farahbakhsh, 1961; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Iran (no locality cited) (Borowski, 2007).

General distribution: Africa, Asia and Middle East, Europe, North America and Mexico, Oceania (Hagstrum et al., 2013).

Host range: Animal invertebrate, animal vertebrate (fishmeal), beverage crops, bake goods, dried fruit, fresh fruit or vegetable, grain, grain product (wheatfeed), medicine, nut, other commodities, oilseed, root, other seed, seasoning, textile, vegetable material, wood and wood product (Hagstrum et al., 2013), insect and other animal collections, wool, skin (Afshar, 1944; Farahbakhsh, 1961; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

***Ptinus (Ptinus) latro* Fabricius, 1775**

Distribution in Iran: Widely distributed (Farahbakhsh, 1961, Modarres Awal, 1997 as *Ptinus brunneus* Duftschmid, 1825), Iran (no locality cited) (Modarres Awal, 1997 as *Ptinus clavipes* Panzer, 1805; Borowski, 2007).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico and Oceania (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, beverage crops, dried fruit, grain, grain product (wheatfeed), medicine, nut, other commodities, oilseed, oilseed product, pulse, pulse product, root, residual commodities, other seed, seasoning, spoiled other commodities, textile and vegetable material (Hagstrum et al., 2013).

***Ptinus (Tectoptynus) tectus* Boieldieu, 1856**

Distribution in Iran: Not exactly defined (Borowski, 2007; Hagstrum et al., 2013).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate (dried insect), animal vertebrate (beef, casein, dried milk, egg, fishmeal, herring meal, meat meal), beverage crops, bake goods, dried fruit, fresh fruit or vegetable (cabbage, carrot, ground cabbage, ground carrot), grain (wheat), grain product (rice flour, wheat bran, flour, germ or meal, wheatfeed, medicine, nut, other commodities, oilseed, oilseed product (copra meal, cottonseed meal), pulse, pulse product (pea flour, soybean meal, root (beet, ground beet, ground potato, potato), root product, other seed, seasoning, textile, vegetable material (alfalfa pellet, grass meal, ground onion, ground swede, onion, rye grass pellet, swede, yeast) and wood (Hagstrum et al., 2013).

Subfamily: Gibbiinae Mulsant & Rey, 1868**Tribe: Gibbiini Jacquelin du Val, 1860*****Gibbium aequinoctiale* Boieldieu, 1854**

Distribution in Iran: General distributed (Modarres Awal, 1997), Northern Iran (no locality cited) (Belles, 1985), Iran (no locality cited) (Borowski, 2007).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Grain, medicine, seasoning, spoiled other commodities, textile and vegetable material (Hagstrum et al., 2013).

***Gibbium psylloides* (de Czeppinski, 1778)**

Distribution in Iran: General distributed (Afshar, 1944; Farahbakhsh, 1961; Habibian, 1987; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Iran (no locality cited) (Belles, 1985; Borowski, 2007).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate, animal vertebrate (fishmeal), beverage crops, bake goods, grain, grain product (wheat flour, wheatfeed, medicine, other commodities, oilseed, oilseed product, pulse, pulse product, residual commodities, other seed, seasoning, textile, vegetable material and wood product (Hagstrum et al., 2013), wool, paper, animal and plant collections, cereals, seeds (in granary) (Afshar, 1944; Farahbakhsh, 1961; Habibian, 1987; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Subfamily: Xyletininae Gistel, 1856

Tribe: Lasiodermi White, 1982

***Lasioderma serricorne* (Fabricius, 1792)**

Distribution in Iran: General distributed (Modarres Awal, 1997; Afshar, 1944; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Niloufari, 1984; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989), Ardabil (Ghahari & Lim, 2012), Golestan (Khormali et al., 2002; Marouf et al., 2009), Kordestan (Allahvaisi, 2013), Mazandaran (Marouf et al., 2009), largely distributed (Modarres awal, 1997).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate, animal vertebrate (dried sardine), beverage crops (cocoa bean, cocoa powder, crushed cocoa bean, roasted coffee bean), bake goods, dried fruit (carob), dried fruit product (crushed carob) fresh fruit or vegetable, grain (maize, wheat), grain product (maize flour, wheat bran, wheat flour, wheat feed), medicine, nut, nut product, other commodities, oilseed (coconut meal, cotton seed, peanut, safflower), oilseed product (cotton seed meal, safflower meal), pulse (bean, cowpea, horse bean, pea), pulse product (lentil flour, soybean meal), root (beet, carrot, licorice), root product, other seed, pomegranate, poppy, seasoning (allspice, anise, black pepper, caraway, caraway flour, cardamom, cayenne pepper, celery salt, chili pepper, cinnamon, clove, coriander, cumin, curry powder, fennel, fennel flour, garlic, ginger, mustard, nutmeg, oregano, paprika, thyme, turmeric), textile, vegetable material (chamomile, cigar tobacco, onion, roselle, rosemary leaf, tobacco, yeast) and wood product (Hagstrum et al., 2013), tobacco products especially cigarette and dried leaves (Modarres Awal, 1997; Afshar, 1944; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Niloufari, 1984; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989).

Tribe: Xyletinini Gistel, 1848

***Xyletinus (Xeronthobius) pallens* Germar, 1824**

Distribution in Iran: Most of areas (Farahbakhsh, 1961; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Iran (no locality cited) (Zahradnik, 2007).

General distribution: Not exactly defined (Farahbakhsh, 1961; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Host range: Tobacco and related products (Farahbakhsh, 1961; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Family: Silvanidae Kirby, 1837

Subfamily: Silvaninae Kirby, 1837

Ahasverus advena (Waltl, 1834)

Distribution in Iran: Guilan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: American, Ethiopian or Oriental origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate (dried insect), animal vertebrate, beverage crops (cocoa), bake goods, dried fruit, fresh fruit or vegetable, grain (wheat), grain product (kibbled wheat, rolled oat, wheat flour, wheat germ), grain spoiled, medicine, nut, nut product, other commodities, oilseed (copra, palm kernel, peanut), oilseed product, pulse, pulse product (soybean meal), root, root product, other seed, seasoning and vegetable material (Hagstrum et al., 2013), grain and other storage products (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Oryzaephilus mercator (Fauvel, 1889)

Distribution in Iran: Generally distributed (Alavi, 1974; Sepasgosarian, 1978; Khorramshahi, 1982; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal vertebrate, beverage crops, bake goods, dried fruit (apricot, date, fig, mulberry, prune, raisin), dried fruit product (apricot meal, date meal, fig meal, raisin meal), fresh fruit or vegetable, grain (maize, millet, oat, wheat), grain product (cracked rice, cracked wheat, ground millet, maize meal, polished rice, rice meal, rolled oat, wheatfeed, wheat meal), medicine, nut (almond, cashew, hazelnut, pine nut, pistachio, walnut), nut product (almond meal, cashew meal), other commodities, oilseed (flax, peanut, rape), oilseed product (coconut meal, cottonseed meal, ground flax, peanut meal, soybean meal), pulse (broad bean, cowpea, soybean), pulse product (crushed broad bean), root, root product, other seed (clover, milled African mango seed, mustard, sunflower), seasoning, vegetable material (grass meal, sugar) (Hagstrum et al., 2013), dried apricot, raisin, date-palm and other storage products (Alavi, 1974; Sepasgosarian, 1978; Khorramshahi, 1982; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Oryzaephilus surinamensis (Linnaeus, 1758)

Distribution in Iran: Generally distributed (Afshar, 1944; Farahbakhsh, 1961; Alavi, 1974; Sepasgosarian, 1978; Habibian, 1987; Shahhosseini & Kamali, 1989; Modarres Awal, 1997), Golestan (Khormali et al., 2002), Razavi Khorasan (Akbari Asl et al., 2009), Tehran (Alipour et al., 2020), Iran (no locality cited) (Abivardi, 2001; Khaleghizaden & Sehhatiasabet, 2006; Valizadegan et al., 2009; Bagheri-Zenouz, 2010; Hosseinzadeh et al., 2010; Ratti & Nardi, 2011).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal invertebrate, animal vertebrate (dried milk), beverage crops, bake goods 1, dried fruit (damaged prune, prune, raisin), dried fruit product, fresh fruit or vegetable, grain (maize, millet, oat, polished rice, teff, wheat), grain product (ground millet, maize meal, cracked barley, cracked oat, cracked rice, rice meal, rolled oat, wheat cracked, kibbled, meal or semolina, wheatfeed), medicine, nut (almond), nut product, other commodities, oilseed (damaged peanut, flax, peanut, rape, safflower, sunflower), oilseed product (broken safflower, coconut meal, cottonseed meal, cracked peanut, ground flax, ground rapeseed, ground sunflower, peanut meal, safflower meal), pulse (broad bean, soybean, pulse product (cracked cowpea, crushed broad bean, soybean meal), root, residual commodities, root product, other seed (clover, mustard), seasoning, ground mustard, vegetable material (alfalfa pellet, grass meal, ground clover, ryegrass pellet, sugar) (Hagstrum et al., 2013).

Family: Staphylinidae Latreille, 1802

Subfamily: Paederinae Fleming, 1821

Tribe: Paederini Fleming, 1821

Paederus fuscipes Curtis, 1826

Distribution in Iran: Fars, Gilan (Assalem, Bandar Pahlavi, Rasht), Golestan (Gorgan), Hormozgan (Minab), Kerman (Anbar-Abad, Djiroft), Khuzestan Mazandaran, Zanzan (Scheerpeltz, 1961; Jarrige, 1971; Bohac, 1981; Nikbakhtzadeh & Sadeghiani, 1999; Tirgari & Nikbakhtzadeh, 2002; Zargari et al., 2003; Smetana, 2004; Nikbakhtzadeh & Tirgari, 1999, 2008).

General distribution: Not exactly defined (Scheerpeltz, 1961; Jarrige, 1971; Bohac, 1981; Nikbakhtzadeh & Sadeghiani, 1999; Tirgari & Nikbakhtzadeh, 2002; Zargari et al., 2003; Smetana, 2004; Nikbakhtzadeh & Tirgari, 1999, 2008).

Host range: Not exactly defined (Scheerpeltz, 1961; Jarrige, 1971; Bohac, 1981; Nikbakhtzadeh & Sadeghiani, 1999; Tirgari & Nikbakhtzadeh, 2002; Zargari et al., 2003; Smetana, 2004; Nikbakhtzadeh & Tirgari, 1999, 2008), Predator on strong products (Shahhosseini & Kamali, 1989).

Paederus littoralis subsp. *ilsae* Bernhauer, 1932

Distribution in Iran: Ardabil (Zargari et al., 2003; Nikbakhtzadeh & Tirgari, 2008); Northern provinces (Afshar, 1944; Modarres Awal, 1997), the presence of this subspecies in Iran is doubtful. It should be confirmed.

General distribution: Not exactly defined (Afshar, 1944; Modarres Awal, 1997).

Host range: Predator on strong products (Afshar, 1944; Modarres Awal, 1997; Shahhosseini & Kamali, 1989).

Subfamily: Staphilininae Latreille, 1802

Tribe: Staphilinini Latreille, 1802

Creophilus maxillosus (Linnaeus, 1758)

Distribution in Iran: Northwest Iran (Talesh) (Fauvel, 1874; Herman, 2001; Smetana, 2004).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal vertebrate (Hagstrum et al., 2013).

Subfamily: Tachyporinae MacLeay, 1825

Tribe: Tachyporini MacLeay, 1825

Tachyporus hypnorum (Fabricius, 1775)

Distribution in Iran: Golestan, Sistan Baluchestan, Tehran (Fauvel, 1875; Scheerpeltz, 1961; Jarrige, 1971; Herman, 2001; Smetana, 2004; Anlas & Newton, 2010).

General distribution: Africa, Asia and Middle East, Europe (Hagstrum et al., 2013).

Host range: Flour milling areas (Hagstrum et al., 2013).

Family: Tenebrionidae Latreille, 1802

Subfamily: Diaperinae Latreille, 1802

Tribe: Diaperini Latreille, 1802

Alphitophagus bifasciatus (Say, 1823)

Distribution in Iran: Gilan and Markazi (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Mediterranean origin. Africa, Asia and Middle East, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Cellars, empty grain bins, feed mills, feed stores, flour mills, grain bins, grain elevators, growing floor at maltings, peanut shelling plants, peanut warehouses, pet stores, rabbit cages and residences (Hagstrum et al., 2013), fungus on stored cereals (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Sitophagus hololeptoides (Laporte de Castelnau, 1840)

Distribution in Iran: Boushehr (Sepasgosarian, 1968; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Europe, North America and South America (Hagstrum & Subramanyam, 2009).

Host range: Avocado, cereal, cocoa, copra, maize, nut, nutmeg, peanut and yam (Hagstrum & Subramanyam, 2009), wheat, barley, groat, flour, bran (in granary) (Sepasgosarian, 1968; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Subfamily: Pimeliinae Latreille, 1802

Tribe: Akidini Billberg, 1820

Cyphogenia gibba Fischer de Waldheim, 1821

Distribution in Iran: Tehran, Gorgan (Afshar, 1944; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Not exactly defined (Afshar, 1944; Shahhosseini & Kamali, 1989; Modarres Awal, 1997)

Host range: Cigarette, dried fruit and bread (in granary) (Afshar, 1944; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Cyphogenia lucifuga (Adams, 1817)

Distribution in Iran: East Azarbaijan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Not exactly defined (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Host range: Different storage product (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Tribe: Tentyriini Eschscholtz, 1831

Mesostena puncticollis Solier, 1835

Distribution in Iran: Khozestan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East and Europe (Hagstrum & Subramanyam, 2009).

Host range: Barley, bone, cotton seed, date, linseed, oilcake and rice bran (Hagstrum & Subramanyam, 2009), different storage products (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Tentyria tessulata Tauscher, 1812

Distribution in Iran: East Azarbaijan, Tehran (Afshar, 1944; Baroughi, 1977; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, bake goods, dried fruit, grain, grain product, grain spoiled, medicine, nut, other commodities, oilseed, oilseed product, pulse, pulse product (soybean, meal), root, residual commodities, root product, other seed, seasoning, textile (Hagstrum et al., 2013), cigarette and bread (in granary) (Afshar, 1944; Baroughi, 1977; Modarres Awal, 1997).

Subfamily: Tenebrioninae Latreille, 1802

Tribe: Alphitobiini Reitter, 1917

Alphitobius laevigatus (Fabricius, 1781)

Distribution in Iran: West Azarbaijan, Kozestan, Gilan, Zanjan, Fars, Kermanshah and Khorasan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Tropical origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal vertebrate, beverage crops, bake goods, dried fruit, fresh fruit or vegetable, grain, grain product, grain spoiled, medicine, nut, other commodities, oilseed, oilseed product, pulse, root, other seed, seasoning and vegetable material (Hagstrum et al., 2013), flour, barley, wheat and other storage products (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Tribe: Blaptini Leach, 1815*Blaps mortisaga* (Linnaeus, 1758)

Distribution in Iran: Markazi and Semnan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997; Zahradnik, 1977).

General distribution: Asia and Middle East and Europe (Hagstrum et al., 2013).

Host range: Grain (Hagstrum et al., 2013), vegetable materials (Shahhosseini & Kamali, 1989; Modarres Awal, 1997; Zahradnik, 1977).

Blaps mucronata Latreille, 1804

Distribution in Iran: West Azarbaijan, Gilan and Fars (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Europe (Hagstrum et al., 2013).

Host range: Grain (damaged) and wheat (Karachi) (Hagstrum et al., 2013), flour and barely (in granary) (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Tribe: Opatrini Brulle, 1832*Gonocephalum rusticum* (Olivier, 1811)

Distribution in Iran: Mazandaran (Farahbakhsh, 1961; Modarres Awal, 1997).

General distribution: Not exactly defined (Farahbakhsh, 1961; Modarres Awal, 1997).

Host range: Potato, tomato, melon and other truck crops (Farahbakhsh, 1961; Modarres Awal, 1997).

Gonocephalum setulosum (Faldermann, 1837)

Distribution in Iran: Gilan and Khozestan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Not exactly defined (Farahbakhsh, 1961; Modarres Awal, 1997).

Host range: Different storage product (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Gonocephalum prolixum (Erichson, 1843)

Distribution in Iran: Not exactly defined (Hagstrum & Subramanyam, 2009).

General distribution: Africa, Europe (Hagstrum & Subramanyam, 2009).

Host range: Bone, cocoa bean, maize, palm kernel, peanut (Hagstrum & Subramanyam, 2009).

Opatroides punctulatus Brulle, 1832

Distribution in Iran: Mazandaran, Gorgan and other northern provinces (Salavatian, 1959; Farahbakhsh, 1961; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East and Europe (Hagstrum & Subramanyam, 2009).

Host range: Almond, barley, residue, valonia beard and wheat (Hagstrum & Subramanyam, 2009), cotton, tobacco and storage products (Salavatian, 1959; Farahbakhsh, 1961; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Tribe: Palorini Matthews, 2003***Palorus depressus* Fabricius, 1790**

Distribution in Iran: Mazandaran, Gorgan and other northern provinces (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Oceania, South America (Hagstrum et al., 2013).

Host range: Root (Hagstrum et al., 2013), flour, wheat (in granary) (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

***Palorus ratzeburgi* (Wissmann, 1848)**

Distribution in Iran: Gilan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: North African origin. Africa, Asia and Middle East, Central America, Europe, North America, Oceania and South America (Hagstrum & Subramanyam, 2009)

Host range: Barley, carob, cassava root, cassava root flour, cassava root meal, Chinese traditional medicine, cocoa, cotton seed, cowpea, dried fruit, grain, grain product, illipenut, lentil, maize, maize meal, malt, millet, oilseed, palm kernel, peanut, peanut (shelled), pulse, rice, sago, sago flour, sesame seed, sorghum, spice, wheat, wheat flour, wheat meal and yam (Hagstrum & Subramanyam, 2009), grain and flour (in granary) (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Tribe: Tenebrionini Latreille, 1802***Alphitobius diaperinus* (Panzer, 1797)**

Distribution in Iran: Khuzestan, Azarbaijan, Tehran, Gilan, Mazandaran, Gorgan, Kordestan, Zanzan and Khorasan (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, beverage crops, bake goods, grain (barley, barley flour, maize, maize flour, rice, rice flour, wheat, wheat broken, wheat flour, wheat ground), grain product, grain spoiled, medicine, nut, other commodities, oilseed (peanut), oilseed product, pulse (black gram, cowpea, pulse product, root, root product, other seed, seasoning and vegetable material (Hagstrum et al., 2013), peanuts, goat, cotton, seed, biscuit, skin, cocoon of silkworm, barley, wheat, bran, potato seed of flax (in granary) (Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

***Tenebrio molitor* Linnaeus, 1758**

Distribution in Iran: Azarbaijan, Tehran, Gilan, Mazandaran, Gorgan, Markazi, Khordestan, Semnan, Zanzan, Esfahan and Hormozgan (Afshar, 1944; Salavatian, 1959; Farahbakhsh, 1961; Alavi, 1974; Baroughi, 1977; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, bake goods, dried fruit, grain (barley, wheat), grain product (ground barley, ground buckwheat, ground millet, oat flake, wheat bran, flour or semolina), nut, other commodities, oilseed (safflower), oilseed product (safflower meal), pulse, pulse product (soybean meal, root, residual commodities, root

product, other seed, seasoning, textile and vegetable material (Hagstrum et al., 2013), entomological collections, flour and related products, dried fruits, tobacco products, meat, vegetable seeds, goat, wool (Afshar, 1944; Salavatian, 1959; Farahbakhsh, 1961; Alavi, 1974; Baroughi, 1977; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

***Tenebrio obscurus* Fabricius, 1792**

Distribution in Iran: Tehran, west Azarbaijan, Khorasan and other northern provinces (Salavatian, 1959; Farahbakhsh, 1961; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia, Europe, Oceania, South America, Central America (Hagstrum & Subramanyam, 2009).

Host range: Entomological collections, flour and related products, dried fruits, tobacco products, meat and vegetable seeds (in granary) provinces (Salavatian, 1959; Farahbakhsh, 1961; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

***Prosodes (Dilopersina) cribella* Baudi, 1874**

Distribution in Iran: Gorgan (Afshar, 1944; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Not exactly defined (Afshar, 1944; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Host range: Cigarette and dried fruits and bread (in granary) (Afshar, 1944; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Tribe: Triboliini Mulsant, 1854

***Latheticus oryzae* Waterhouse, 1880**

Distribution in Iran: Generally distributed (Davatchi, 1950; Salavatian, 1959; Farahbakhsh, 1961; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Africa, Asia and Middle East and Europe (Hagstrum et al., 2013).

Host range: Beverage crops, grain and oilseed (Hagstrum et al., 2013), cotton, tobacco and storage products) (Davatchi, 1950; Salavatian, 1959; Farahbakhsh, 1961; Bagheri Zenouz, 1986; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

***Tribolium castaneum* (Herbst, 1797)**

Distribution in Iran: Generally distributed (Davatchi, 1947; Salavatian, 1959; Farahbakhsh, 1961; Balachowsky, 1969; Alavi, 1974; Bagheri Zenouz, 1986; Habibian, 1987; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Oriental origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania and South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, beverage crops, bake goods, dried fruit, dried fruit product, fresh fruit or vegetable, grain (hungary rice, maize, millet, rice, sorghum, teff, wheat), grain product (barley flour, cracked or ground barley, ground buckwheat, maize cracked, flour, grit or meal, millet flour, meal or semolina, oat cracked, flour or rolled, polished rice, rice flour, rice meal, sorghum flour or meal, wheat bran, flour,

meal or semolina, wheatfeed), grain spoiled, medicine, nut (walnut), nut product, other commodities, oilseed cottonseed, flax, peanut, rape, safflower, sesame, sunflower), oilseed product (broken peanut, broken safflower, cottonseed flour, ground flax, ground rape, ground sunflower, peanut meal, safflower meal), pulse, broad bean, soybean), pulse product (chickpea flour, cowpea flour, cracked cowpea, crushed broad bean, lentil flour, moth bean flour, mung bean flour, pea flour, pigeonpea flour, soybean cracked or flour, soybean meal), root (cassava flour), residual commodities, root product cassava meal, yam flour or meal, other seed (clover, ground clover, ground flax, ground mustard, mustard), seasoning, spoiled, other commodities, textile, vegetable material (cellulose, dried mushroom, dried parsley, hemp), wood and wood product (Hagstrum et al., 2013).

***Tribolium confusum* Jacquelin du Val, 1861**

Distribution in Iran: Generally distributed (Davatchi, 1947; Salavatian, 1959; Farahbakhsh, 1961; Balachowsky, 1969; Alavi, 1974; Bagheri Zenouz, 1986; Habibian, 1987; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: Ethiopian origin. Africa, Asia and Middle East, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal invertebrate and vertebrate, beverage crops (tea), bake goods (biscuit), dried fruit, dried fruit product, fresh fruit or vegetable, grain (hungary rice, maize, millet, rice, wheat), grain product (cracked barley, cracked maize, cracked oat, cracked soybean, crack wheat, graham flour, ground barley, ground buckwheat, ground millet, maize meal, millet flour or semolina, polished rice, rice flake, rice meal, rolled oat, wheat bran, flour or meal, wheat semolina , medicine, nut, nut product, other commodities, oilseed (flax, peanut, rape, safflower, sunflower), oilseed product (broken safflower, cracked peanut, ground flax, ground rape, ground sunflower, safflower meal, peanut meal), pulse (broad bean, soybean), pulse product (cracked cowpea, crushed broad bean, soybean meal), root, residual commodities, root product, other seed (clover, ground clover, ground mustard, mustard), seasoning, textile, vegetable material (dried mushroom, dried parsley, hemp, herb, malt coffee (Hagstrum et al., 2013).

Tribe: Ulomini Blanchard, 1845

***Uloma culinaris* Linnaeus, 1758**

Distribution in Iran: Mazandaran Province, northern Iran (Katlav et al., 2014).

General distribution: Asia and Middle East, Europe (Hagstrum et al., 2013).

Host range: Granaries, hotels, meat shops (Hagstrum et al., 2013).

Family: Trogossitidae Latreille, 1802

Subfamily: Trogossitinae Latreille, 1802

Tribe: Trogossitini Latreille, 1802

***Tenebroides mauritanicus* (Linnaeus, 1758)**

Distribution in Iran: Generally distributed (Davatchi, 1947; Salavatian, 1959; Alavi, 1974; Behdad, 1984; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: African origin. Africa, Asia and Middle East, Central America and Caribbean, Europe, North America and Mexico, Oceania, South America (Hagstrum et al., 2013).

Host range: Animal vertebrate, beverage crops, bake goods, dried fruit, fresh fruit or vegetable, grain (rice, wheat), grain product (wheat flour), grain spoiled, medicine, nut, nut product, other commodities, oilseed, oilseed product, pulse, pulse product (soybean meal), root, residual commodities, root product, other seed, seasoning, vegetable material (Hagstrum et al., 2013).

Temnoscheila caerulea (Olivier, 1790)

Distribution in Iran: Northern provinces (Afshar, 1944; Farahbakhsh, 1961; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

General distribution: America, palaeartic region (Kolibáč, 2013).

Host range: Wheat, bread, dried fruits and other products (in granary) (Afshar, 1944; Farahbakhsh, 1961; Shahhosseini & Kamali, 1989; Modarres Awal, 1997).

Discussion

The majority of stored product pests in Iran include the families Anthicidae, Anthribidae, Bostrichidae, Carabidae, Cerambycidae, Chrysomelidae, Cleridae, Cryptophagidae, Curculionidae, Dermestidae, Erotylidae, Laemophloeidae, Latridiidae, Melyridae, Mycetophagidae, Nitidulidae, Ptinidae, Silvanidae, Staphylinidae, Tenebrionidae and Trogossitidae. One hundred twenty two species from 59 genera and 45 tribes are listed from Iran. Among the 60 genera, *Cryptophagus* (Cryptophagidae) with 12 species is the most species-reich genus. Iran is divided into 31 provinces of which stored-product beetles have been recorded from 22 provinces: Alborz, Boushehr, East Azarbaijan, Esfahan, Fars, Gilan, Golestan, Hamadan, Hormozgan, Kerman, Kermanshah, Khorasan, Khuzestan, Kordestan, Kuhgiluyeh and Boyerahmad, Lorestan, Markazi, Mazandaran, Sistan and Baluchestan, Tehran, West Azarbaijan, Zanjan (Fig.1).

There is a broad range of natural enemies, including parasitoids, predators and pathogens that attack various developmental stages of stored product pests. For example *Xylocoris flavipes* (Reuter) (Hemiptera: Anthocoridae) (Lattin, 2000) and *Anisopteromalus calandrae* (Howard) (Hymenoptera: Pteromalidae) have been reported as parasitoid of the following pests: *Sitophilus granarius*, *Sitophilus zeamais*, *Rhyzopertha dominica*, *Stegobium paniceum*, *Lasioderma serricorne*, *Acanthoscelides obtectus* and *Callosobruchus maculatus* (Williams & Floyd, 1971; Arbogast & Mullen, 1990; Reichmuth, 2000; Reichmuth et al., 2007; Ngamo et al., 2007; Kazemi et al., 2008). *Theocolax elegans* (Westwood) (Hymenoptera: Pteromalidae) has been reported as a parasitoid of *S. granarius*, *S. oryzae*, *S. zeamais*, *R. dominica*, *S. paniceum*, *Callosobruchus spp.* and of the grain moth *Sitotroga cerealella* (Reichmuth et al., 2007), *Lariophagus distinguendus* (Förster) (Hymenoptera: Pteromalidae) has been reported as potential agent for biological control for a wide number of beetles that infest stored agricultural products (Steidle & Schöller, 1997): *S. oryzae* (Lucas & Riudavest, 2002), *S. granarius* (Wen & Brower, 1994; Steidle & Schöller, 2002), *R. dominica* (Menon et al., 2002), *L. serricorne* (Steidle et al., 2006), *S. paniceum*, *A. obtectus* and *Ptinus clavipes* (Reichmuth, 2000). *Habrobracon hebetor* (Say) (Hymenoptera: Braconidae) develops on larvae of many Lepidoptera, mainly members of the family Pyralidae (Schöller, 1998; Kovalenkov

& Meschcheryakova, 1983; Balevsky, 1984; Gerling, 1971; Amir-Maafi & Chi, 2006; Shoukat, 2012; Payne et al., 2011; Farag et al., 2012; Ba et al., 2013; Huang, 1986; Hopper, 2003; Grieshop et al., 2006). Generally, establishment of biological control agents in stored products is difficult and this is the reason for limited research in this area of biological control involving stored product pests (e.g. Bostrichidae, Bruchinae, Curculionidae, Dermestidae, Tenebrionidae, etc.). Hence, faunistic surveys are necessary to find new country records, new species, new host and natural enemies in the various stored products in Iran.




Figure 1. Number of reported species of stored product beetles by province.

Acknowledgments

The Iranian Research Institute of Plant Protection supported this research. We express our sincere appreciation.

Conflict of Interests

The author declares that there is no conflict of interest regarding the publication of this paper.

References

- Abai, M. (1969) List of Cerambycidae family in Iran. *Journal of Applied Entomology and Phytopathology*, 28, 47–54.
- Abai, M. (1984) *List of pests of forest trees and shrubs of Iran*. Plant Pest and Diseases Research institute, Tehran. 147 pp.
- Abd-Rabou, S., Ghahari, H., Huang, J. & Boucek, Z. (2005) New records of aphelinid and pteromalid wasps (Hymenoptera: Chalcidoidea: Aphelinidae, Pteromalidae) from Iran. *Egyptian Journal of Agricultural Research*, 83 (4), 1619–1623.
- Abivardi, C. (2001) Iranian Entomology, Volume 2: Applied Entomology. Schriftenreihe der Stiftung Franz Xaver Schnyder von Wartensee. *Zentralbibliothek Zurich*, 59, 445–1033.
- Adeli, E. (1972) Beitrag zur Kenntnis der Forstschädlichen Insekten des Iran. I. Coleoptera. *Zeitschrift für Angewandte Entomologie*, 70, 8–14. <https://doi.org/10.1111/j.1439-0418.1972.tb02144.x>
- Afshar, D.J. (1944) Les noms scientifique de quelques coleopteres de l' Iran et leurs importance en agriculture. Ministry of Agriculture, Tehran. 22 pp.
- Akbari Asl, M.H., Talebi, A. A. & Kamali, H. (2009) Stored product pests and their parasitoid wasps in Mashhad, Iran. *Advances in Environmental Biology*, 3 (3), 239–243.
- Alavi, A. (1974) *The Opium poppy*. Plant Pest and Diseases Research Institute, Tehran, 306 pp.
- Allahvaysi, S. (2013) Controlling *Lasioderma serricorne* F. (Col.: Anobiidae) by Fumigation and Packaging. *World Applied Sciences Journal*, 28 (12), 1983–1988.
- Alipour, M., Talebi, A., Fathipour, Y. & Mehrabadi, M. (2020) Life table parameters of *Cephalonomia tarsalis* (Hym.: Bethyridae), parasitoid of *Oryzaephilus surinamensis* (Col.: Silvanidae). *Plant Protection (Scientific Journal of Agriculture)*, 42(4), 39–48.
- Amini, S., Hosseini, R. & Sohani, M.M. (2013) A faunal study of bark beetles (Coleoptera: Curculionidae: Scolytinae) in Guilan province in North of Iran. *Entomofauna*, 34, 169–176.
- Amir-Maafi, M. & Chi, H. (2006) Demography of *Habrobracon hebetor* (Hymenoptera: Braconidae) on two pyralid hosts (Lepidoptera: Pyralidae). *Annals of the Entomological Society of America*, 99(1), 84–90. [https://doi.org/10.1603/0013-8746\(2006\)099\[0084:DOHHHB\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2006)099[0084:DOHHHB]2.0.CO;2)
- Anlas, S. & Newton, A.F. (2010) Distributional checklist of the Staphylinidae (Coleoptera) of Iran, with new and additional records. *Linzer biologische Beiträge*, 335–388.
- Anton, K.W. (1998) Results of the Czechoslovak-Iranian entomological expeditions to Iran, 1970, 1973 and 1977. Coleoptera Bruchidae. *Casopis Narodniho Muzea Rada Prirodovedna*, 167 (1-4), 73–90.
- Anton, K.W. (2010) Bruchinae. In: Löbl, I. & Smetana, A. (eds.), *Catalogue of Palaearctic Coleoptera*. Vol. 6. Chrysomeloidea. Apollo Books, Stenstrup, pp. 347–353.
- Arbogast, R.T. & Mullen, M.A. (1990) Interaction of maize weevil (Coleoptera: Curculionidae) and parasitoid *Anisopteromalus calandrae* (Hymenoptera: Pteromalidae) in a small bulk of stored corn. *Journal of Economic Entomology*, 83, 2462–2468. <https://doi.org/10.1093/jee/83.6.2462>
- Aref, S.P. & Valizadegan, O. (2015) *Eucalyptus krueseana* Muel essential oil: chemical composition and insecticidal effects against the lesser grain borer, *Rhyzopertha dominica* F. (Coleoptera: Bostrichidae). *Biharean Biologist*, 9, 93–97.
- Ashouri, S. & Shayesteh, N. (2009) Insecticidal activities of Black Pepper and Red Pepper in powder form on adults of *Rhyzopertha dominica* (F.) and *Sitophilus granarius* (L.). *Pakistan Entomologist*, 31(2), 122–127.
- Azimi, A.R., Rabee, A., Siahpoosh, A. & Mozaffari, M. (1993) Report of some Nitidulid species in corn fields of Khuzestan province. *Proceedings of 11th Iranian Plant Protection Congress, 28 August-2 September, Rasht, Iran*. University of Guilan, pp. 85.
- Ba, M., Baoua, I., N'diaye, M., Dabire-Binso, C., Sanon, A. & Tamo, M. (2013) Biological control of the millet head miner *Heliochelioides albipunctella* in the Sahelian region augmentative releases of the

- parasitoid wasp *Habrobracon hebetor*: effectiveness and farmers' perception. *Phytoparasitica*, 41 (5), 569-576. <https://doi.org/10.1007/s12600-013-0317-x>
- Bagheri-Zenouz, E. (1986) *Les animaux nuisibles aux produits entreposes. Vol. I, les Coleopteres de predateurs de produits alimentaires et industriels*. Sepehr Publications., Tehran. 309 pp.
- Bagheri-Zenouz, E. (2010) *Pests of Stored Products and Management to Maintain. Bioecology of Insects, Acari and Microorganism*. 2nd edition. University of Tehran Publication, Iran, 449 pp. [in Persian].
- Bagheri-Zenouz, E. (2013) *Pests of stored products and management to maintain*. 2nd Edition. University of Tehran Press, Tehran. 449 pp.
- Bagheri, M.R. & Nasr Isfahani, M. (2011) The fauna of harmful and beneficial arthropods of medicinal and range plants in Isfahan. *Journal of Entomological Research*, 3(2), 119-132. [in Persian, English summary].
- Balachowsky, A. (1949) Coloptères Scolytides. *Faune de France*, 50, 1-320.
- Balachowsky, A.S. (1969) Entomologie appliquee a l'agriculture. Tome I, Vol. II, Coleopteres. *Journal of Applied Entomology and Phytopathology*, 564 pp.
- Balevsky, N. (1984) Use of the parasite *Habrobracon hebetor* Say for biological control. *Rastitelna Zashchita*, 32, 19-28.
- Baroughi, H. (1977) Etude de la fauna entomologique (Tabriz et environs). Partie. II, order des coleopteres. *Journal of Applied Entomology and Phytopathology*, 8, 1-47.
- Behdad, E. (1982) *Pest of field crops in Iran*. Plant Pest and Diseases Research Institute, Esfahan, 589 pp. (In Persian).
- Behdad, E. (1984) *Pest of fruit crop in Iran*. Neshat publication, Esfahan, Iran, 822 pp. (In Persian).
- Behdad, E. (1988) *Pest and diseases of forest trees and shrubs of Iran*. Sepehr publication, 808 pp. (In Persian).
- Belles, X. (1985) *Sistemática, filogenia y biogeografía de la subfamilia Gibbiinae (Coleoptera, Ptinidae)*. Treballs del Museu de Zoologia. Vol. 3, Museu de Zoologia. Ajuntament de Barcelona. Barcelona, 94 pp.
- Boháč, J. (1981) Results of the Czechoslovak-Iranian entomological expeditions to Iran. Coleoptera, Staphylinidae. Three new species of Staphylinids from Iran. *Acta Entomologica Musei Nationalis Pragae*, 40, 355-358.
- Bonadona, P. (1970) Contribution a la fauna de l'Iran. 18. Coleopteres Anthicidae. *Annales de la Société entomologique de France, New Series*, 6 (2), 379-383.
- Borowiec, L. & Anton, K.W. (1993) Materials to the knowledge of seed beetles of the Mediterranean Subregion (Coleoptera, Bruchidae). *Annals of the Upper Silesian Museum in Bytom Entomology*, 4, 99-152.
- Borowiec, L. (1983) Survey of seed-beetles of Bulgaria (Coleoptera, Bruchidae). *Polskie Pismo Entomologiczne*, 53, 107-127.
- Borowiec, L. (1987) The seed-beetles (Coleoptera, Bruchidae) from the Middle East. *Polskie Pismo Entomologiczne*, 57, 601-616.
- Borowski, J. (2007) Gibbiinae and Ptininae. In: Lobl, I. & Smetana, A. (eds.), *Catalogue of Palaearctic Coleoptera. Vol. 4, Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea, Cucujoidea*. Apollo Books, Stenstrup, Denmark, pp. 328-339.
- Borror, D.J., Triplehorn, C.A. & Johnson, N.F. (1989) *An Introduction to the study of insects*, 6th ed. New York, Saunders College Publishing. 824 pp.
- Bouchard, P., Bousquet, Y., Davies, A.E., Alonso-Zarazaga, M.A., Lawrence, J.F., Lyal, C.H.C., Newton, A.F., Reid, C.A.M., Schmitt, M., Slipinski, S.A. & Smith, A.B.T. (2011) Family group names in Coleoptera (Insecta). *ZooKeys*, 88, 1-972. <https://doi.org/10.3897/zookeys.88.807>
- Bousquet, Y., Bouchard, P., Davies, A.E. & Sikes, D.S. (2013) Checklist of beetles (Coleoptera) of Canada and Alaska. Second edition. *Zookeys*, 360, 1-44. <https://doi.org/10.3897/zookeys.360.4742>

- Brodský, O. & Winkler, J.R. (1988) Results of the Czechoslovak – Iranian Entomological Expeditions to Iran 1970, 1973 and 1977. Coleoptera, Cleridae. *Acta Entomologica Musei Nationalis Pragae*, 42, 293–305. <https://doi.org/10.3906/zoo-1212-8>
- Bulak, Y., Yildirim, E., Kadej, M. & Háva, J. (2013) Contribution to the knowledge of the Dermestidae (Coleoptera) fauna of Turkey. *Turkish Journal of Zoology*, 37, 621–626. <https://doi.org/10.3906/zoo-1212-8>
- Campbell, J.F., Arthur, F.H. & Mullen, M.A. (2004) Insect management in food processing facilities. *Advanced Food Nutrient Research*, 48, 239–295. [https://doi.org/10.1016/S1043-4526\(04\)48005-X](https://doi.org/10.1016/S1043-4526(04)48005-X)
- Chandler, D.S., Uhmman, G., Nardi, G. & Telnov, D. (2008) Family Anthicidae Latreille, 1819. In: Löbl, I. & Smetana, A. (eds.), *Catalogue of Palaearctic Coleoptera*. Vol. 5. Apollo Books, Stenstrup, pp. 421–455.
- Chokouhian, A. (1973) Biology et nombre de generation de *Callosobruchus maculatus* F. *Journal of Applied Entomology and Phytopathology*, 34, 1–40.
- Dashan, M., Sadeghi, S., Bakhshi, Y. & Malek-Hosseini, M.J. (2014) First record and redescription of *Niptus hololeucus* (Faldermann, 1835) from Kangohar Cave (Coleoptera: Ptinidae). *Iranian Journal of Animal Biosystematics*, 10 (2), 81–85.
- Davatchi, A. (1946) Note sur les bruchides nuisibles les legumineuses en Iran. *Journal of Applied Entomology and Phytopathology*, 2, 1–8.
- Davatchi, A. (1947) Quelques insectes nuisibles aux cereals a la farine (2). *Journal of Applied Entomology and Phytopathology*, 4, 1–16.
- Davatchi, A. (1950) Les coleopteres trouvee dans le d'origine americaine. *Journal of Applied Entomology and Phytopathology*, 10, 1–2.
- Decelle, J. & Lodos, N. (1989) Contribution to the knowledge of the legume weevils of Turkey (Coleoptera: Bruchidae). *Bulletin et Annales de la Société royale belge d'Entomologie*, 125, 163–212.
- Esmaili, M. (1983) *The important pests of fruit trees*. Sepehr publications, Tehran. 578 pp.
- Eyidozohi, K., Khormali, S., Ravan S. & Barahoei, H. (2013) Introduction of seven wasps` parasitoid species associated with stored food product pests in Golestan Province. *Plant Pests Research*, 3, 69–72.
- Farag, M.M.A., Sayeda, S.A. & El-Husseini, M.M. (2012) Life history of *Habrobracon hebetor* (Say) (Hymenoptera: Braconidae) parasitizing *Cadra (Ephestia) cautella* (Walker) (Lepidoptera: Pyraliadae) on dried date fruits. *Egyptian Journal of Biological Pest Control*, 22 (1), 73–77.
- Farahbakhsh, G.H. (1961) Checklist of important insect and other enemies of plants and agricultural products in Iran. *Ministry Agriculture Department Plant protection, Tehran*, 1, 1–153.
- Fauvel A. (1874) Faune Gallo-Rhénane ou descriptions des insectes qui habitent la France, la Belgique, la Hollande, le Luxembourg, les provinces Rhénanes et la Valais avec tableaux synoptiques et planches gravées. *Bulletin de la Société Linnéenne de Normandie*, 8(2), 167–340. [Note: This publication includes the 2e supplément (HERMAN 2001)].
- Fauvel, A. (1875) *Faune Gallo-Rhénane Catalogue systématique des staphylinides de la Faune Gallo-Rhénane avec l'addition synonymique des espèces européennes, siberiennes, caucasiennes et Méditerranéennes et descriptions nouvelles*. Tome 2. Livraison 6. Caen: Le Blanc-Hardel, pp. 545–738.
- Forghani, S.H.R. & Marouf, A. (2015) An introductory study of storage insect pests in Iran. *Biharean Biologist*, 9 (1), 59–62.
- Gerling, D. (1971) Occurrence, abundance, and efficiency of some local parasitoid attacking *Spodoptera littoralis* (Lepidoptera: Noctuidae) in selected cotton fields in Israel. *Annals of the Entomological Society America*, 64, 492–499. <https://doi.org/10.1093/aesa/64.2.492>
- Ghahari, H. & Borowiec, L. (2017) A checklist of seed-beetles (Coleoptera: Chrysomelidae: Bruchinae) from Iran. *Zootaxa*, 4268 (2), 215–237. <https://doi.org/10.11646/zootaxa.4268.2.3>

- Ghahari, H. & Colonelli, E. (2012) Curculionoidea from Golestan province, Northern Iran. *Fragmenta Entomologica*, 44, 101–161. <https://doi.org/10.4081/fe.2012.37>
- Ghahari, H., Huang, J., Ostovan, H. & Rastegar, J. (2010) Notes on the Iranian fauna of Pteromalidae (Hymenoptera). *Efflatounia*, 10, 21–25.
- Ghahari, H. & Lim, J. (2012) A checklist on Iranian Bethyloidea (Hymenoptera: Chrysidoidea). *Linzer biologische Beiträge*, 44 (1), 1071–1077.
- Grieshop, M.J., Flinn, P.W. & Nechols, J.R. (2006) Biological control of indianmeal moth (Lepidoptera: Pyralidae) on finished stored products using egg and larval parasitoids. *Journal of Economic Entomology*, 99 (4), 1080–1084. <https://doi.org/10.1093/jee/99.4.1080>
- Gussakovskii, V.V. (1949) *Verdnie zhivotnye srednei azii: Coleoptera*. Izdatelstvo Akademii Nauk, Moskva, Leningrad. 288 pp.
- Háva, J. & Kalík, V. (1999) The dermestid beetles (Coleoptera: Dermestidae) of Iran. Part 1: genus *Dermestes*. *Acta Societatis Zoologicae Bohemicae*, 63, 443–450.
- Háva, J. (2003a) Faunistic records of Dermestidae from Libya (Coleoptera: Bostrichoidea). *Genus*, 14(1), 43–47.
- Háva, J. (2003b) Notes on Dermestidae (Coleoptera) with description of eight new species. *Annales historico-naturales Musei Nationalis Hungarici*, 95, 19–35.
- Háva, J. (2004) World keys to the genera and subgenera of Dermestidae (Coleoptera), with descriptions, nomenclature and distributional records. *Acta Musei Nationalis Pragae, Series B, Natural History*, 60 (3–4), 149–164.
- Háva, J. (2006) Notes on the genus *Phradonoma* from the Middle East with description of nine new species (Coleoptera: Dermestidae: Megatominae). *Genus*, 17 (1), 79–94.
- Háva, J. (2007) Dermestidae. In: Lobl, I. & Smetana, A. (eds.) *Catalogue of Palaearctic Coleoptera*. Vol. 4, *Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea, Cucujoidea*. Apollo Books, Stenstrup, Denmark, pp. 299–320.
- Háva, J. & Herrmann, A. 2011. New faunistic records of Dermestidae (Coleoptera). Part 6. *Boletín de la Sociedad Entomológica Aragonesa*, 49, 227–228.
- Háva, J. (2015) *World Catalogue of Insects*. Volume 13. Dermestidae (Coleoptera). Brill, Leiden, The Netherlands.
- Habibian, A. (1987) *Sitophilus oryzae* (L.) in Gilan (Caspian coastal). *Journal of Applied Entomology and Phytopathology*, 54 (1, 2), 45–54.
- Hagstrum, D.W. & Subramanyam, B. (2009) *Stored-product insect resource*. AACC International Press, USA, 518 pp.
- Hagstrum, D.W., Klejdysz, T., Subramanyam, B. & Nawrot, J. (2013) *Atlas of stored product insect and mites*. AACC International Press, USA. 588 pp.
<https://doi.org/10.1016/B978-1-891127-75-5.50009-5>
- Haines, C.P. (1991) *Insects and arachnids of tropical stored products: their biology and identification*. 2nd ed. Natural Resources Institute, Overseas Development administration, Kent, United Kingdom. 246 pp.
- Hedjazi, R. & Soleimani, P. (1976) Studied on the occurrence of natural resistance to the *Lyctus brunneus* Steph. In some species of wood plants. *University of Tehran Faculty of Forest*, 8, 1–40
- Herman, L.H. (2001) Catalog of the Staphylinidae (Insecta: Coleoptera). 1758 to the end of the second millennium. Volumes I–VII. *Bulletin of the American Museum of Natural History*, 265, 1–4218.
<https://doi.org/10.1206/0003-0090.265.1.1>
- Herrmann, A. & Háva, J. (2008) Two new *Phradonoma* species from Iran. *Journal of Entomological Society of Iran*, 28(1), 87–91.
- Herrmann, A., Háva, J. & Fallahzadeh, M. (2010) Contribution to the genus *Anthrenus* from Iran (Coleoptera: Dermestidae). *Journal of the Entomological Research Society*, 12, 113–115.

- Hoffmann, A. (1945a) *Fauna de France. 44, Coleopters bruchides et anthribidies*. Paul Lechevalier, Paris. 184 pp.
- Hoffmann, A. (1945b) *Fauna de France. 59, Coleopters curculionides*. *Journal of Applied Entomology and Phytopathology*, 2, 487-1208.
- Hopper, K.R. (2003) United States Department of Agriculture-agricultural research service research on biological control of arthropods. *Pest Management Science*, 59, 643-653. <https://doi.org/10.1002/ps.707>
- Hosseinzadeh, A., Shayesteh, N., Zolfaghari, H.R., Babaei, M., Zareshahi, H., Mostafavi, H.A. & Fatollahi, H. (2010) Gamma radiation sensitivity of different stages of saw-toothed grain beetle *Oryzaephilus surinamensis* L. (Coleoptera: Silvanidae). *Journal of Plant Protection Research*, 50(3), 250-255. <https://doi.org/10.2478/v10045-010-0044-y>
- Huang, X.F. (1986) Use of *Habrobracon hebetor* Say in granary pest control. *Chinese Journal Biological Control*, 2, 78-80.
- Jarrige, J. (1971) Contribution a la faune de l' Iran 21. Coléoptères brachelytra. *Annales de la Société Entomologique de France*, (N.S.) 7 (2), 483-502.
- Johnson, C., Otero, J.C. & Leschen, R.A.B. (2007) Catalogue Cryptophagidae. In: Lobl, I. & Smetana, A. (eds.) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, Copenhagen, Denmark, pp. 65-70.
- Karami, Z., Bagheri Zenooz, E. & Zhantiev, R.D. (2004) Notes on the *Anthrenus* species (Col.: Dermestidae) in Fars province. *Proceedings of 16th Iranian Plant Protection Congress, 2004, 27 August - 1 September, Tabriz, Iran*. Ministry of Agriculture, Tehran, pp. 143.
- Katlav, A., Hajiqanbar, H. & Talebi, A.A. (2014) First record of the genus *Acanthomastix* Mahunka, 1972 (Acari: Dolichocybidae) from Asia, with the description of a new species. *International Journal of Acarology*, 40 (1), 7-14. <https://doi.org/10.1080/01647954.2014.883429>
- Kazemi, F., Talebi, A.A. & Fathipour, Y. (2008) A Study on the Biological and Behavioural Parameters of *Anisopteromalus calandrae* (Hym.: Pteromalidae), a Parasitoid of *Callosobruchus maculatus* (Col.: Bruchidae) on Chickpea. *Journal of Science and Technology of Agriculture and Natural Resources*, 12, 313-322.
- Kazemi, F., Talebi, A.A., Fathipour, Y. & Farahani, S. (2009) A Comparative Study on the Effect of Four Leguminous Species on Biological and Population Growth Parameters of *Callosobruchus maculatus* (F.) (Col.: Bruchidae). *Advances in Environmental Biology*, 3 (3), 226-232.
- Khaleghizaden, A. & Sehhatisabet, M.E. (2006) Contribution to the knowledge of the diet of Iranian birds. *Ekologia*, 15 (1-2), 145-150.
- Keshavarzi, D., Moemenbellah-Fard, M.D., Fereidooni, M. & Montazeri, M. (2015) First report of *Dermestes frischii* Kugelann (Coleoptera: Dermestidae) on a human corpse, south of Iran. *International Journal of Forensic Science and Pathology*, 3 (4), 113-115. <https://doi.org/10.19070/2332-287X-1500027>
- Khoobdel, M., Ma'rouf, A., Farajzadeh, D., Vatani, H., Riazipour, M. & Joneydi, N. (2011) Frequency and diversity of pest arthropods in stored cereals in a military unit. *Iranian Journal of Military Medicine*, 13 (2), 81-87.
- Khormali, S., Taheri, M.S. & Borumand, H. (2002) Faunal investigation on stored pests in Gonbad and Minoodasht. *Journal of Applied Entomology and Phytopathology*, 70 (1), 13-23 [in Persian, English abstract].
- Khorramshahi, A. (1982) *Collection and identification of storage pests in Tabriz and vicinity*. University of Tabriz, Faculty of Agricultural. 21 pp.
- Knížek, M. (2011) Platypodinae and Scolytinae. In: Löbl, I. & Smetana, A. (eds.) *Catalogue of Palaearctic Coleoptera*. Vol. 7. Curculionoidea I. Apollo Books, Stenstrup, pp. 201-251.
- Kolibáč, J. (2013) Trogossidae: A review of the beetle family, with a catalogue and keys. *Zookeys*, 366, 1-194. <https://doi.org/10.3897/zookeys.366.6172>

- Kovalenkov, V.G. & Meshcherykova, T.V. (1983) Reserve plantations for *Trichogramma* and *Habrobracon*. *Zashchita Rastenii*, 12, 16-17.
- Lattin, J.D. (2000) Minute pirate bugs (Anthocoridae). In: Schaefer C.W. & Panizzi A.R. (eds.) *Heteroptera of economic importance*. CRC Press, Washington, pp. 607-637. <https://doi.org/10.1201/9781420041859.ch26>
- Legalov, A.A., Ghahari, H. & Arzanov, Yu.G. (2010) Annotated catalogue of Curculionidae-beetles (Coleoptera: Anthribidae, Rhynchitidae, Attelabidae, Brentidae, Brachyceridae, Dryophthoridae and Curculionidae) of Iran. *Amurian Zoological Journal*, II (3), 191-244.
- Lucas, E. & Riudavets, J. (2002) Biological and mechanical control of *Sitophilus oryzae* (Coleoptera: Curculionidae) in rice. *Journal of Stored Product Research*, 38(3), 293-304. [https://doi.org/10.1016/S0022-474X\(01\)00033-9](https://doi.org/10.1016/S0022-474X(01)00033-9)
- Mahdneshin, Z., Safaralizadeh, M.H. & Hgosta, Y. (2009) Study on the efficacy of Iranian isolates of *Beauveria bassiana* (Balsamo) Vuillemin and *Metarhizium anisopliae* (Metsch.) Sorokin against *Rhyzopertha dominica* F. (Coleoptera: Bostrichidae). *Journal of Biological Science*, 9, 170-174. <https://doi.org/10.3923/jbs.2009.170.174>
- Majka, C.G. & Langor, D. (2010) Contributions towards an understanding of the Cryptophaginae (Coleoptera, Cryptophagidae) of Atlantic Canada. *ZooKeys*, 35, 13-35. <https://doi.org/10.3897/zookeys.35.314>
- Marouf, A., Shayesteh, N. & Rostam-kolaii, S.E. (2009) Assessment of the efficacy of current recommendations for fumigation with Phosphine to control *Lasioderma serricornis* F. (Col., Anobiidae) in tobacco storages. *Journal of Entomological Research Society*, 1 (3), 249-259.
- Mehrabadi, M., Bandani, A.R., Saadati, F. & Mahmudvand, M. (2011) A-amylase activity of stored products insects and its inhibition by medicinal plant extracts. *Journal of Agricultural Science and Technology*, 13, 1173-1182.
- Ménétrières, E. (1832) *Catalogue raisonné des objets de zoologie recueillis dans un voyage au Caucase et jusqu'aux frontières actuelles de la Perse entrepris par ordre de S.M. L'Empereur*. St. Petersburg, L'Académie Impériale des Sciences, 271 pp. <https://doi.org/10.5962/bhl.title.63878>
- Menon, A., Flinn, P.W., Barry, A. & Dover, B.A. (2002) Influence of temperature on the functional response of *Anisopteromalus calandrae* (Hymenoptera: Pteromalidae), a parasitoid of *Rhyzopertha dominica* (Coleoptera: Bostrichidae). *Journal of Stored Product Research*, 38, 463-469. [https://doi.org/10.1016/S0022-474X\(01\)00050-9](https://doi.org/10.1016/S0022-474X(01)00050-9)
- Mifsud, D. & Knížek, M. (2009) The bark beetles (Coleoptera: Scolytidae) of the Maltese Islands (Central Mediterranean). *Bulletin of the Entomological Society of Malta*, 2, 25-52.
- Mirzayans, H. (1950) Quelques Cerambycidae de l'Iran. *Journal of Applied Entomology and Phytopathology*, 10, 12-21.
- Modarres Awal, M. (1994) *List of agricultural pests and their natural enemies in Iran*. Ferdowsi University Press, Mashhad. 364 pp.
- Modarres Awal, M. (1997) *List of agricultural pests and their natural enemies in Iran* (revised edition). Ferdowsi University Press, Mashhad. 429 pp.
- Mound, L., (1989) *Common insect pests of stored food products. A guide to their identification*. 7th edition. London, British Museum (Natural History). 68 pp.
- Mroczkowski, M. (1968) Distribution of the Dermestidae (Coleoptera) of the world with a catalogue of all known species. *Annales Zoologici*, 26, 15-191.
- Naeem, A. & Akhyani, A. (1988) the fig souring disease in Kashan and its transmission factors. *Journal of Applied Entomology and Phytopathology*, 55(1,2), 31-42.
- Ngamo, T.S.L., Kouninki, H., Ladang, Y.D., Ngassoum, M.B., Mapongmestem, P.M. & Hance, T. (2007) Potential of *Anisopteromalus calandrae* (Hymenoptera: Pteromalidae) as biological control

- agent of *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae). *Africa Journal of Agricultural Research*, 2, 168–172.
- Nikbakhtzadeh, M. & Sadeghiani, C. (1999) Dermatite due à deux espèces de *Paederus* au sud de l'Iran. *Bulletin de la Société de Pathologie Exotique*, 92 (1), 56.
- Nikbakhtzadeh M. & Tirgari, S. (1999) Two medically important beetles of Fars province, Iran: *Paederus* spp. (Col: Staphylinidae) & its linear dermatitis. *Iranian Journal of Public Health*, 28 (1-4), 73–86. [in Persian].
- Nikbakhtzadeh, M. & Tirgari, S. (2008) Medically important beetles (Insecta: Coleoptera) of Iran. *Journal of Venomous Animals and Toxins including Tropical Diseases*, 14 (4), 597–618. <https://doi.org/10.1590/S1678-91992008000400004>
- Nikdel, M., Fathalizadeh, A. & Moniri, V. (2016) Identification of seed feeder Bruchidae of pasture legumes in East Azarbaijan province. *Journal of Applied Entomology and Phytopathology*, 83 (2), 269–173.
- Niloufari, P. (1984) *Wood destroying insect of Iran*. University of Tehran, Faculty of Natural Resource, 87 pp.
- Niloufari, P. (1985) Studies on the infestation behaviour of the powder-post beetle *Lyctus brunneus* (Steph.) and its physical control in the wood yards of the Caspian forests of Iran. *International Research Group on Wood Preservation*, 3 pp.
- Olsen, A.R. (1998) Regulatory action criteria for fi lth and other extraneous materials. II. Allergenic mites: an emerging food safety issue. *Regulatory Toxicology & Pharmacology*, 28, 190–198. <https://doi.org/10.1006/rtp.1998.1270>
- Otero, J.C. (2013) Cryptophaginae (Coleoptera) de la Región Paleártica occidental. *Coleoptera Monographs*, 4, 1–296.
- Payne, W., Tapsoba, H., Baoua, I., Malick, B., N'diaye, M. & Dabire-Binso, C. (2011) On-farm biological control of the pearl millet heard miner. Realization of 35 years of unsteady progress in Mali, Burkina Faso and Niger. *International Journal of Agricultural Sustainability*, 9(1), 186–193. <https://doi.org/10.3763/ijas.2010.0560>
- Picard, F. (1949) *Fauna de France*. 20. *Coleopteres Cerambycidae*. Librairie de la Facult des Sciences, Paris. 166 pp.
- Plavilstshikov, N.N. (1940) *Fauna de l'USSR. Coleoptera*. Academie de sciences de l'USSR, Moscow, Leningrad, 758 pp.
- Ratti, E., & Nardi, G. (2011) Silvanidae, Cucujidae e Laemophloeidae di Sardegna: catalogo provvisorio (Coleoptera: Cucujoidea). *Conservazione habitat invertebrate*, 5, 461–492.
- Rees, D. (1990) The effect of temperature and relative humidity on population growth of three *Liposcelis* species (Psocoptera: Liposcelidae) infesting stored products in tropical countries. *Bulletin of Entomological Research*, 80, 353–358. <https://doi.org/10.1017/S0007485300050562>
- Reichmuth, C. (2000) Biological control in stored product protection, *IOBC/WPRS Bulletin*, 23(10), 11–23.
- Reichmuth, C., Schöller, M. & Ulrichs, C. (2007) *Stored product pests in grain. Morphology –biology–damage– control*. AgroConcept Verlagsgesellschaft, Bonn. 170 pp.
- Sakenin, H., Samin, N., Hava, J., Bunalski, M., Naderian, H. & Kubisz, D. (2018) A contribution to the study of Iranian Coleoptera with some new records. *Wiadomości Entomologiczne*, 37 (2), 96–109.
- Salavatian, M. (1959) List of pests of cotton, kanaf, flax, suger-beet, suger-cane, tobacco, sesame and castor bean. *Journal of Applied Entomology and Phytopathology*, 18, 62–75.
- Salehi Ardakani, A. & Nasserzadeh, H. (2014) First record and biology of coffee bean weevil, *Araecerus fasciculatus* De Geer, on pesticide plant, *Melia azedarach* L. from Iran. *International Journal of Biosciences*, 5(12), 486–491. <https://doi.org/10.12692/ijb/5.12.486-491>

- Scheerpeltz, O. (1961) Die von den Herren W. Richter und Dr. F. Schäuuffele in den Jahren 1954 und 1956 im Sudiran aufgefundenen Staphyliniden (Col.). *Stuttgarter Beiträge zur Naturkunde*, 50, 1-31.
- Schöller, M. (1998) Biologische Bekämpfung vorratsschädlicher Arthropoden mit Räubern und Parasitoiden-Sammelbericht und Bibliographie. In: Reichmuth, Ch. (eds.) *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft*, Parey, Berlin, Heft, pp. 85-189.
- Sepasgosarian, H. (1968) Ein neuer lagerschadling im Iran, *Sitophagus hololeptoides* Canst. (Col.: Tenebrionidae). *Journal of Applied Entomology and Phytopathology*, 27, 147-149.
- Sepasgosarian, H. (1978) *Storage pests of Iran and their control*. University of Tehran, 1026, 287 pp.
- Shahhosseini, M.J. & Kamali, K. (1989) Checklist of insect, Mites and rodents affecting stored products in Iran. *Journal of Entomological Society of Iran*, Supplement, 5, 1-47.
- Shayesteh, N. (1957) Studied of phostoxin for flour beetle under laboratory and natural conditions. *Journal of Applied Entomology and Phytopathology*, 38, 41-46.
- Shires, S.W. (1982) Observations on *Liposcelis bostrychophilus* Badonnel (Psocoptera). *Tropical Stored Products Information*, 44, 11-14.
- Shoukat, G.A. (2012) Larval parasitoids of *Lobesia botrana* (Denis and Schiffermüller, 1775) (Lepidoptera: Tortricidae) in Orumieh vineyards. *Journal of Agricultural Science and Technology*, 14(2), 267-274.
- Sinha, R.N. & Watters, F.L. (1985) *Insect pests of flourmills, grain elevators, and feed mills and their control*. Research Station Winnipeg, Manitoba Publication. 177 pp.
- Smetana, A. (2004) Family Staphylinidae (except subfamilies Pselaphinae and Scaphidiinae). In: LÖBL I. & A. SMETANA (eds), *Catalogue of Palaearctic Coleoptera*. Volume 2. Hydrophiloidea, Histeroidea, Staphyliniidea. Apollo Books, Stenstrup, pp. 237-698.
- Somerfield, K.G. (1981) Recent aspects of stored product entomology in New Zealand. *New Zealand Journal of Agricultural Research*. 24, 403-408. <https://doi.org/10.1080/00288233.1981.10423407>
- Soleimani, P. & Knopf, H.E. (1976) *Stomatium fulvum* vill. (= *unicolor* ol.) (Col., Cerambycidae) in Iran. *Journal of Applied Entomology*, 82 (1-4), 372-376. <https://doi.org/10.1111/j.1439-0418.1976.tb03426.x>
- Steidle, J.L.M. & Schöller, M. (1997) Olfactory host location and learning in the granary weevil parasitoid *Lariophagus distinguendus* (Hymenoptera: Pteromalidae). *Entomologia Experimentalis Applicata*, 95, 185-192.
- Steidle, J.L.M. & Schöller, M. (2002) Fecundity and ability of the parasitoid *Lariophagus distinguendus* (Hymenoptera: Pteromalidae) to find larvae of the granary weevil *Sitophilus granarius* (Coleoptera: Curculionidae) in bulk grain. *Journal of Stored Products Research*, 38, 43-53. [https://doi.org/10.1016/S0022-474X\(00\)00044-8](https://doi.org/10.1016/S0022-474X(00)00044-8)
- Steidle, J.L.M., Gantert, C., Prozell, S. & Schöller, M. (2006) Potential der Lagererzwespe *Lariophagus distinguendus* zur Bekämpfung des Tabakkäfers *Lasioderma serricornis*. *Mitteilungen Deutsche Gesellschaft für Allgemeine und Angewandte Entomologie*, 15, 295-298.
- Stejskal, V. & Verner, P.H. (1996) Long-term changes of cockroach infestations in Czech and Slovak food-processing plants. *Medical Veterinary Entomology*, 10, 103-104. <https://doi.org/10.1111/j.1365-2915.1996.tb00090.x>
- Stejskal, V., Kučerová, Z. & Háva, J. (2005) *Trogoderma longisetosum* and *Trogoderma variabile* (Coleoptera, Dermestidae) as two new stored product pests for the Czech Republic. *Plant Protection Science*, 41(1), 42-45. <https://doi.org/10.17221/2738-PPS>
- Taheri, M.S. (1991) The effect of low temperature on development stage of *Callosobruchus maculatus* F. (Coleoptera: Bruchidae). *Journal of Applied Entomology and Phytopathology*, 58 (1-2), 29-36.
- Telnov, D. & Ghahari, H. (2018) An annotated checklist of the Anthicidae and pediline Pyrochroaie (Insecta: Coleoptera) of Iran, with thirteen new country records. *Zootaxa*, 4497 (4), 451-491. <https://doi.org/10.11646/zootaxa.4497.4.1>

- Tezcan, S., Karsavuran, Y., Pehlivan, E. & Háva, J. (2004) Contribution to the Dermestidae (Coleoptera) fauna of Turkey along with new records. *Türkiye Entomoloji Dergisi*, 28(1), 27–37.
- Thind, B.B. & Dunn, J.A. (2002) A laboratory evaluation of a regulated airflow through wheat at four combinations of temperature and humidity on the productivity of three species of stored product mites. *Experimental and Applied Acarology*, 27, 89–102.
<https://doi.org/10.1023/A:1021522321940>
- Tirgari, S. & Nikbkkhtzadeh, M.R. (2002) *Paederus* beetles (Coleoptera: Staphylinidae): an urban problem in Iran. *Proceedings of the 4th International Conference on Urban Pests, Charleston, South Carolina*, pp. 401–407.
- Valizadegan, O., Pourmirza, A.A. & Sataralizadeh, M.H. (2009) Combination of microwaves radiation and cold storage for control of *Oryzaephilus surinamensis* (L.) (Col. Silvanidae). *Journal of biological Sciences*, 9(3), 231–236. <https://doi.org/10.3923/jbs.2009.231.236>
- Wegrzynowicz, P. (2007) Cucujidae and Laemophloeidae. In: Löbl, I. & Smetana, A. (eds.) *Catalogue of Palaearctic Coleoptera. Vol. 4, Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea, Cucujoidea*. Apollo Books, Stenstrup, Denmark, pp. 502–507.
- Weidner, H. & Rack, G. (1984) *Tables de détermination des principaux ravageurs des denrées entreposées dans les pays chauds*. GTZ Schriftenreihe No. 129; Eschborn, Germany. 148 pp.
- Wen, B. & Brower, J.H. (1994) Suppression of maize weevil, *Sitophilus zeamais* (Coleoptera: Curculionidae), populations in drums of corn by single and multiple releases of the parasitoid *Anisopteromalus calandrae* (Hymenoptera: Pteromalidae). *Journal of Kansas Entomological Society*, 67, 331–339.
- Williams, R.N. & Floyd, E.H. (1971) Effect of two parasitoids *Anisopteromalus calandrae* and *Choetospila elegans*, upon population of maize weevil under laboratory and natural conditions. *Journal of Economic Entomology*, 64, 1407–1408. <https://doi.org/10.1093/jee/64.6.1407>
- Zahradnik, J. (1977) *A field guide in color to insects*, Translated by Olga Kuthanova. Octopus Books Limited, London. 319 pp.
- Zahrddnik, P. (2007) Ptinidae (except Gibbiinae and Ptininae). In: Löbl, I. & Smetana, A. (eds.) *Catalogue of Palaearctic Coleoptera. Vol. 4, Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea, Cucujoidea*. Apollo Books, Stenstrup, Denmark, pp. 339–362.
- Zargari, O., Kimyai-asadi A., Fathalikhani, F. & Panahi, M. (2003) *Paederus* dermatitis in northern Iran: a report of 156 cases. *International Journal of Dermatology*, 42 (8), 608–612.
<https://doi.org/10.1046/j.1365-4362.2003.01771.x>
- Zehzad, B., Kiabi, B.H. & Madjnoonian, H. (2002) The natural areas and landscape of Iran: an overview. *Zoology in the Middle East*, 26, 7–10. <https://doi.org/10.1080/09397140.2002.10637915>
- Zhantiev, R.D. (1976) Zhuki kozheedy fauny SSSR. [The skin eaterseater's family Dermestidae of fauna of the USSR.] *Izdatelstvo Moskovskogo Universiteta, Moscow*. [In Russian].
- Ziaee, M., Safaralizadeh, M.H. & Shayesteh, N. (2006) Effects of temperature and exposure interval on the toxicity of Silico Sec® against two stored products insects. *Pakistan Entomologist*, 28 (1), 45–50.

چک لیست سوسک‌های محصولات انباری ایران (Insecta: Coleoptera)

نجمه ابراهیمی*

بخش تحقیقات حشره‌شناسی، مؤسسه تحقیقات گیاهپزشکی کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران.
* پست الکترونیکی نویسنده مسئول مکاتبه: n.ebrahimi@iripp.ir

| تاریخ دریافت: ۰۷ بهمن ۱۳۹۸ | تاریخ پذیرش: ۲۷ خرداد ۱۳۹۹ | تاریخ انتشار: ۰۹ تیر ۱۳۹۹ |

چکیده: مطالعه گسترده‌ای برای تعیین لیست سوسک‌های محصولات انباری در ایران انجام شد. از مجموع ۱۲۲ گونه سوسک‌های محصولات انباری، ۵۹ جنس و ۴۵ قبیله در ایران به ثبت رسیده است. خانواده‌های دارای بیشترین گونه شامل Dermestidae (۲۴ گونه) و Tenebrionidae (۲۴ گونه) هستند و سپس Chrysomelidae (۱۵ گونه)، Cryptophagidae (۱۳ گونه)، Ptinidae (۱۰ گونه)، Curculionidae (۶ گونه)، Staphylinidae (۴ گونه)، Nitidulidae (۴ گونه)، Bostrichidae (۳ گونه)، Laemophloeidae (۳ گونه)، Silvanidae (۳ گونه)، Trogossitidae (۲ گونه)، Anthicidae (۲ گونه)، Cleridae (۲ گونه) و خانواده‌های Carabidae، Cerambycidae، Anthribidae، Erotylidae، Latridiidae، Melyridae و Mycetophagidae دارای یک گونه می‌باشند. مطالعه حاضر هم یک گزارش فونستیک بوده و هم در طراحی و توسعه مدیریت تلفیقی سوسک‌های محصولات انباری بسیار مفید است.

واژگان کلیدی: سخت‌بالپوشان، چک لیست، ایران، سوسک‌های محصولات انباری