

## Article

<http://zoobank.org/urn:lsid:zoobank.org:pub: 4D5B30A3-02B1-425D-8486-1B29758A79DB>

# A new species of the genus *Vittacus* Keifer (Acari: Eriophyidae) from Iran

Jing-Feng Guo<sup>1</sup>, Hussein Sadeghi<sup>2\*</sup> & Ali Gol<sup>2</sup>, Xiao-Feng Xue<sup>1</sup>

1. Department of Entomology, Nanjing Agricultural University, Nanjing, Jiangsu 210095, China; Emails: 2012102106@njau.edu.cn, xfxue@njau.edu.cn

2. Department of Plant Protection, College of Agriculture, Ferdowsi University of Mashhad, Iran; Emails: sadeghin@um.ac.ir, ali.gol@stu-mail.um.ac.ir

\* Corresponding author

## Abstract

A new species of the genus *Vittacus*, *V. veronicae* Guo, Sadeghi & Xue **sp. nov.** is described and illustrated based on females collected from *Veronica anagallis-aquatica* L. (Plantaginaceae) in Ali Abad-e Katool, Golestan Province of Iran. This is the first report of the genus *Vittacus* from Iran. A key to the species of *Vittacus* is provided.

**Key words:** Trombidiformes, Eriophyoidea, taxonomy, plant feeding mites, *Veronica anagallis-aquatica*.

## Introduction

Recently, the fauna of Eriophyoidea of Iran has been focused by several investigations. For example, Xue *et al.* (2009) reviewed the Iranian related literature and gave a preliminary list of 89 species. Since then more eriophyoids have been recorded and described from Iran by several authors (e.g. Kamali & Jalaeeian 2011, 2013; Xue *et al.* 2011, 2012, 2013a, 2013b; Lotfollahi *et al.* 2012, 2013, 2014a, 2014b) increasing the known Iranian eriophyoid fauna at more than 120 species. However, eriophyoid fauna of many plant species in many parts of this country has remained undiscovered. In a survey of the Eriophyidae associated with *Veronica anagallis-aquatica* L. (Plantaginaceae) in Ali Abad-e Katool, Golestan Province of Iran, a new species was collected and is described herein.

## Material & methods

During a survey of the eriophyoid fauna in Ali Abad-e katool region, Golestan Province of Iran, which was conducted during seasonal growth of 2013, mite specimens were collected directly from *Veronica anagallis-aquatica* L. (Plantaginaceae) under a dissecting microscope. Mite specimens were cleared in lactic acid for 7–10 days at room temperature and then mounted in Hoyer's medium. No fibers were used in the mounting medium between slide and coverslip. Specimens were examined with the aid of a Zeiss A2 (Germany) research microscope with phase contrast and schematic drawings were made. The morphological terminology used herein follows Lindquist (1996) and the generic classification is made according to Amrine *et al.* (2003). Specimens were

measured following de Lillo *et al.* (2010). The holotype female measurement precedes the corresponding range for paratypes (given in parentheses). All measurements are in micrometres ( $\mu\text{m}$ ) and are lengths when not otherwise specified. The holotype and 5 paratypes are deposited in the Arthropod/Mite Collection of the Department of Entomology, Nanjing Agricultural University, Jiangsu Province, China. Also, two paratypes are held at the Department of Plant Protection, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran.

## Taxonomy

Family Eriophyidae Nalepa

Subfamily Phyllocoptinae, tribe Anthocoptini

Genus *Vittacus* Keifer

### *Vittacus veronicae* Guo, Sadeghi & Xue sp. nov. (Fig. 1)

#### Description

*Female* (n = 8) - Body fusiform, 168 (168–175), 62 (61–64) wide, 56 (56–57) thick; light yellow.

*Gnathosoma* - 20 (19–20), projecting obliquely down, pedipalp coxal setae *ep* 2 (2–3), dorsal pedipalp genual setae *d* 5 (5–6).

*Prodorsal shield* - 50 (50–51) including frontal lobe, 60 (60–61) wide; semi-circular; smooth; frontal shield lobe 10 (10–11); scapular tubercles 4 (3–4) on rear margin, 47 (43–48) apart, scapular setae (*sc*) 5 (5–6), projecting posteriorly and diverging.

*Coxigenital region* - with 6 (6–7) microtuberculated semiannuli between coxae and genitalia. Coxal plates I with few lines, coxal plates II smooth, anterolateral setae on coxisternum I (*1b*) 10 (9–10), 12 (12–13) apart; proximal setae on coxisternum I (*1a*) 18 (17–19), 9 (9–10) apart; proximal setae on coxisternum II (*2a*) 43 (41–45), 23 (23–24) apart; prosternal apodeme 6 (5–6).


*Leg I* - 28 (28–30), femur 12 (12–13), basiventral femoral setae *bv* 10 (8–10); genu 5 (4–5), antaxial genual setae *l''* 20 (17–20); tibia 6 (6–7), paraxial tibial setae *l'* 3 (3–4), located at 1/4 from base; tarsus 5 (4–5), tarsal setae *ft'* 18 (16–19), tarsal setae *ft''* 22 (20–24), tarsal setae *u'* 4 (3–4); tarsal empodium (*em*) 5 (5–6), simple, 4-rayed,  $\omega$  7 (7–8), slight knobbed.

*Leg II* - 25 (25–28), femur 11 (11–12), basiventral femoral seta (*bv*) 10 (10–11); genu 5 (5–6), antaxial genual setae *l''* 5 (4–5); tibia 4 (4–5) long; tarsus 5 (5–7), paraxial, fastigial, tarsal setae *ft'* 5 (4–5), antaxial, fastigial, tarsal setae *ft''* 17 (15–18), tarsal setae *u'* 4 (3–4); tarsal empodium (*em*) 5 (5–6), simple, 4-rayed, tarsal solenidion ( $\omega$ ) 7 (7–8), slight knobbed.

*Opisthosoma* - dorsally with 8 (8–9) broad semiannuli, smooth, ventrally with 69 (67–70) semiannuli, with rounded microtubercles on rear annular margins. Setae *c2* 9 (9–10), 54 (53–55) apart, on ventral semiannulus 13 (12–14); setae *d* 46 (46–50), 36 (35–36) apart, on ventral semiannulus 27 (26–28); setae *e* 15 (14–15), 17 (17–18) apart, on ventral semiannulus 45 (44–46); setae *f* 20 (18–20), 16 (16–18) apart, on 5th ventral semiannulus from rear; setae *h1* 3 (2–3), setae *h2* 60 (60–62).

*Female genitalia* - 20 (20–22) wide, 13 (13–14) long, coverflap with 10 (10–11) longitudinal ridges, setae *3a* 14 (13–14), 17 (17–18) apart.

Male - unknown.



**Figure 1.** Schematic drawings of *Vittacus veronicae* Guo, Sadeghi & Xue sp. nov.  
- D: Dorsal view of female; AL: Lateral view of anterior body region; CG: Female coxigenital region; em: Empodium; IG: Internal female genitalia; LO: Lateral view of annuli; L1: Leg I; PM: Lateral view of posterior opisthosoma.

#### Type material

Holotype - Female (slide number Iran D6.1, marked holotype), from *Veronica*

*anagallis-aquatica* L. (Plantaginaceae), Ali Abad-e Katool (36° 54' N, 54° 53' E), Golestan Province, Iran, 29 August 2013, coll. Ali Gol, deposited as a slide mounted specimen in the Arthropod/Mite Collection of the Department of Entomology, NJAU, Jiangsu Province, China.

*Paratypes* - Five females (slide number Iran D6.2–D6.6) and 2 females (slide number Iran D6.7–D6.8) from *V. anagallis-aquatica* L., same data as holotype, deposited in the Arthropod/Mite Collection of the Department of Entomology, NJAU, Jiangsu Province, China and Department of Plant Protection, Faculty of Agriculture, Ferdowsi University of Mashhad, Iran, respectively.

#### *Relation to host*

Vagrant. No damage to the host was observed.

#### *Etymology*

The specific designation *veronicae* is from the generic name of host plant, *Veronica*.

#### *Differential diagnosis*

This species is similar to *Vittacus mandshurica* Xue, Guo & Hong, 2013, but can be differentiated from the latter by prodorsal shield smooth (prodorsal shield with admedian and submedian lines in *V. mandshurica*), dorsal annuli smooth (dorsal annuli with fissures in *V. mandshurica*), semiannuli between coxae and genitalia with microtubercles (semiannuli between coxae and genitalia smooth in *V. mandshurica*).

#### Key to the world species of *Vittacus* Keifer

- |  |   |
|--|---|
| 1. Coxal area smooth.....  | 2   |
| - Coxal area with granules or short lines.....   | 4   |
| 2. Dorsal annuli with faint lines, median line on prodorsal shield present.....                          | <i>Vittacus bougainvilleae</i> Abou-Awad & El-Banhawy, 1991 |
| - Dorsal annuli smooth, median line absent.....  | 3   |
| 3. Prodorsal shield with a hexagon-shape in the middle.....  |   |
| .....  | <i>Vittacus pluchae</i> Abou-Awad & Nasr, 1986              |
| - Prodorsal shield with a U-shape in the middle.....   | <i>Vittacus cornutus</i> Xue, Song & Hong, 2010             |
| 4. Prodorsal shield smooth.....  | 5   |
| - Prodorsal shield with median, admedian or submedian lines.....   | 6   |
| 5. Dorsal annuli with short lines, empodium 3-rayed.....   |   |
| .....  | <i>Vittacus humuli</i> Xue, Song & Hong, 2005               |
| - Dorsal annuli smooth, empodium 4-rayed.....  |   |
| .....  | <i>Vittacus veronicae</i> Guo, Sadeghi & Xue sp. nov.       |
| 6. Scapular tubercles placed ahead of rear margin of prodorsal shield, dorsal annuli smooth.....         | <i>Vittacus fissistigmae</i> Wei & Kuang, 1997              |
| - Scapular tubercles on rear margin of prodorsal shield, dorsal annuli with short lines or fissures..... | 7   |
| 7. Median line on prodorsal shield present.....  |   |
| .....  | <i>Vittacus cannabis</i> Xue, Guo & Hong, 2013              |
| - Median line absent.....  | 8   |
| 8. Empodium 6-rayed, prodorsal shield with granules anteriorly.....                                      |   |

- ..... *Vittacus shaoguanicus* Xue, Song & Hong, 2010  
 - Empodium 4-rayed, prodorsal shield without granules..... 9  
 9. Dorsal annuli with 13 wide annuli, seta *h1* present.....  
 ..... *Vittacus mandshurica* Xue, Guo & Hong, 2013  
 - Dorsal annuli with 22 annuli, seta *h1* absent..... *Vittacus mansoni* Keifer, 1969

### Acknowledgements

This work was partly supported by Ferdowsi University of Mashhad, Mashhad, Iran. We thank Xiao Han of Nanjing Agricultural University, Nanjing, China for comments on an earlier draft of this manuscript.

### References

- Abou-Awad, B.A. & Nasr, A.K. (1986) An eriophyid mite, *Vittacus pluchaeae* sp. nov. (Acari: Eriophyoidea: Eriophyidae), on *Pluchea dioscoridis* L. (Compositae) from Egypt. *Acarologia*, 27(2): 159–161.
- Abou-Awad, B.A. & El-Banhawy, E.M. (1991) New mites on the family Eriophyidae from Kenya (Acari: Eriophyoidea). *Acarologia*, 32(4): 329–333.
- Amrine, J.W. Jr., Stasny, T.A. & Flechtmann, C.H.W. (2003) *Revised keys to world genera of Eriophyoidea* (Acari: Prostigmata). Indira Publishing House, West Bloomfield, MI, U.S.A., 244 pp.
- de Lillo, E., Craemer, C., Amrine, J.W. Jr. & Nuzzaci, G. (2010) Recommended procedures and techniques for morphological studies of Eriophyoidea (Acari: Prostigmata). *Experimental and Applied Acarology*, 51: 283–307.
- Kamali, H. & Jalaeian, M. (2011) Infestation of white mulberry trees to bud mite, *Aceria mori* Keifer (Acari: Eriophyoidea: Eriophyidae) in Iran. In: Kazemi, Sh. & Saboori, A. (Eds.), *Abstract and proceeding book of the First Persian Congress of Acarology*. International Center for Science, High Technology & Environmental Sciences, Kerman, Iran, p. 73.
- Kamali, H. & Jalaeian, M. (2013) Present status of gall mites (Acari: Prostigmata: Eriophyoidea) on Almond trees in Khorasan Razavi Province (North East of Iran). In: Joharchi, O. & Saboori, A. (Eds.) *Abstract and proceeding book of the Second International Persian Congress of Acarology*, Karaj, Iran, p. 17.
- Keifer, H.H. (1969) *Eriophyid studies* C-2. U.S. Department of Agriculture, Agricultural Research Service, 24 pp.
- Lindquist, E.E. (1996) External anatomy and notation of structures In: Lindquist, E.E., Sabelis, M.W. & Bruun, J. (Eds.), *Eriophyoid mites: their biology, natural enemies and control*. Elsevier, World Crop Pests, Vol. 6, pp. 3–31.
- Lotfollahi, P., Haddad Irani-Nejad, K., Khanjani, M., Moghadam, M. & de Lillo, E. (2012) Two new eriophyoid mite species (Acari: Prostigmata: Eriophyidae) on *Euphorbia* spp. (Euphorbiaceae) from Iran. *Zootaxa*, 3556: 55–60.
- Lotfollahi, P., Haddad Irani-Nejad, K., Khanjani, M., Moghadam, M. & de Lillo, E. (2013) Eriophyoid mites (Acari: Prostigmata: Eriophyidae) associated with Compositae in Iran. *Zootaxa*, 3664 (3): 349–360.
- Lotfollahi, P., Haddad Irani-Nejad, K. & de Lillo, E. (2014) Three new species from the subfamily Phyllocoptinae (Acari: Trombidiformes: Eriophyidae) in Iran. *Zookeys*, 426: 17–27.

- Lotfollahi, P., Haddad Irani-Nejad, K., Rikan, M.H. & de Lillo, E. (2014) Two new eriophyid mites (Acaria: Trombidiformes: Eriophyoidea) on *Rubia tinctorum* L. in Iran. *International Journal of Acarology*, 40(3): 225–229.
- Nalepa, A. (1892) Neue Arten der Gattung *Phytoptus* Duj. und *Cecidophyes* Nal. *Denkschriften der Kaiserlichen Akademie der Wissenschaften. Mathematisch-naturwissenschaftlich Klasse*, Wien, 59: 525–544.
- Wei, S.-G. & Kuang, H.-Y. (1997) Three new species of Phyllocoptinae from China (Acaria: Eriophyidae). *Acta Zootaxonomica Sinica*, 22(1): 45–49.
- Xue, X.F., Guo, J.F. & Hong, X.Y. (2013) Eriophyoid mites from Northeast China (Acaria: Eriophyoidea). *Zootaxa*, 3689: 1–123.
- Xue, X.F. & Hong, X.Y. (2005) A new genus and eight new species of Phyllocoptini (Acaria: Eriophyidae: Phyllocoptinae) from China. *Zootaxa*, 1039: 1–17.
- Xue, X.F., Sadeghi, H. & Hong, X.Y. (2009) Eriophyoid mites (Acaria: Eriophyoidea) from Iran, with descriptions of three new species, one new record and a checklist. *International Journal of Acarology*, 35: 461–483.
- Xue, X.F., Sadeghi, H. & Hong, X.Y. (2012) Four new eriophyoid mite species (Acaria: Eriophyoidea: Eriophyidae) from Iran. *Zootaxa*, 3544: 28–40.
- Xue, X.F., Sadeghi, H., Hong, X.Y. & Sinaie, S. (2011) Nine eriophyoid mite species from Iran. *ZooKeys*, 143: 23–45.
- Xue, X.F., Sadeghi, H., Hong, X.Y. & Sinaie, S. (2013) New species and records of eriophyid mites from Iran (Acaria: Eriophyidae). *Systematic & Applied Acarology*, 18(1): 41–52.
- Xue, X.F., Song, Z.W. & Hong, X.Y. (2010) Five new species of Anthocoptini from China (Acaria: Eriophyidae). *Zootaxa*, 2666: 29–44.

Received: 10 August 2014

Accepted: 26 November 2014

Published: 15 January 2015

#### COPYRIGHT

 Guo et al. Persian Journal of Acarology is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

## گونه جدیدی از جنس *Vittacus* Keifer (Acari: Eriophyidae) از ایران

جينگ-فنگ گو<sup>۱</sup>، حسين صادقی<sup>۲</sup>، علی گل<sup>۳</sup> و ژیائو-فنگ ژو<sup>۱</sup>

۱. گروه حشره شناسی، دانشگاه کشاورزی نانجینگ، نانجینگ، جیانگسو ۹۵۰۱۰۲، چین؛ رایانامه‌ها:

*xfxue@njau.edu.cn*, *2012102106@njau.edu.cn*

۲. گروه گیاهپردازی دانشکده کشاورزی دانشگاه فردوسی مشهد، ایران؛ رایانامه‌ها:

*ali.gol@stu-mail.um.ac.ir*, *sadeghin@um.ac.ir*

چکیده

گونه‌ای از جنس *V. veronicae* Guo, Sadeghi & Xue sp. nov. به نام *Vittacus* Keifer اساس نمونه‌های ماده جمع‌آوری شده از گیاه *Veronica anagallis-aquatica* L. (Plantaginaceae) در علی‌آباد کتول، استان گلستان ایران توصیف و ترسیم می‌شود. این نخستین گزارش جنس *Vittacus* از ایران است. کلید شناسایی گونه‌های جنس *Vittacus* در دنیا ارایه شده است. واژگان کلیدی: *Eriophyoidea*, *Trombidiformes*, *Rodebndi*, کنه‌های گیاه‌خوار.

تاریخ دریافت: ۱۳۹۳/۵/۱۹

تاریخ پذیرش: ۱۳۹۳/۹/۵

تاریخ چاپ: ۱۳۹۳/۱۰/۲۵