

Article

Mites of the families Trachyuropodidae Berlese and Urodiaspididae Trägårdh (Acari: Mesostigmata: Uropodina) from Iran

Shahrooz Kazemi*, Mojtaba Mohammad-Dustar-sharaf and Sepideh Saberi

Department of Biodiversity, Institute of Science and High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman, Iran; E-mails: shahroozkazemi@yahoo.com, sh.kazemi@kgut.ac.ir, mojtaba.doostar@gmail.com, saberi20202s@gmail.com

* Corresponding author

Abstract

Herein, two genera, *Urodiaspis* Berlese and *Urojanetia* Berlese, and four species, *Trachyuropoda hirschmanni* Pecina, 1980, *Urodiaspis tecta* (Kramer, 1876), *Urodiaspis pannonica* Willmann, 1951 and *Urojanetia excavata* (Wasmann, 1899), are reported for the first time from Iran. Additionally, *Urodiaspis pannonicasimilis* Bal and Özkan, 2009 is considered as junior synonym of *U. pannonica*, and new morphological data for this species represented.

Key words: Fauna; Monogynaspida; new synonym; soil inhabiting mites; Uropodoidea.

Introduction

The cosmopolitan monogynaspid mites of the infraorder Uropodina include four superfamilies, 35 families, 300 genera and about 2300 described species (Beaulieu *et al.* 2011; Halliday 2015). The families Urodiaspididae Trägårdh, 1944 and Trachyuropodidae Berlese, 1917 currently are classified within the superfamily Uropodoidea (Lindquist *et al.* 2009; Beaulieu *et al.* 2011). The Urodiaspididae includes three genera and 26 species (Beaulieu *et al.* 2011). One of these genera is *Urodiaspis* which was erected by Berlese (1916) to accommodate *U. tecta* (Kramer, 1876) as its type species. The genus was originally described as a member of the family Uropodidae Kramer, 1881 and this classification followed by some subsequent authors (e.g. Hirschmann and Zirngiebl-Nicol 1964; Kadite and Petrova 1977; Krantz and Ainscough 1990). On the other hand, some authors placed *Urodiaspis* in the family Urodinychidae Berlese, 1917 (e.g. Karg 1989; Mašán 2001), or under the family Dinychidae Berlese, 1916 (e.g. Lindquist *et al.* 2009), or in a separate family as Urodiaspididae (e.g. Beaulieu *et al.* 2011; Kontschán 2013). The *Urodiaspis* mites usually occur in habitats with high level of organic materials, such as litter, moss, decaying wood, humus. Also, they have been occasionally collected from nest of vertebrates, ants and bumble-bees (Wiśniewski and Hirschmann 1993; Karg 1989; Mašán 2001). Błoszyk *et al.* (2004) considered *Urodiaspis tecta* and *Urodiaspis pannonica* Willmann, 1951 as reproducing only (or almost exclusively) in parthenogenetic way. The family Trachyuropodidae includes 16 genera (Kontschán 2015), and about 100 described species (Wiśniewski and Hirsch-

mann, 1993; Beaulieu *et al.* 2011), and most of the family members have been reported in association with ants, but several species inhabit in soil, leaf litter and moss (Mašán, 2001; Kontschán and Starý 2013).

Iranian mites of Uropodina are poorly known, and so far only 38 species of this infraorder have been reported from Iran (Kazemi and Kontschán 2007, 2014; Kazemi and Rajaei 2013; Kontschán and Hajizadeh 2013; Arjomandi *et al.* 2013; Arjomandi and Kazemi 2014, 2016; Kazemi and Abolghasemi 2016). Herein, the genera *Urodiaspis* and *Urojanetia* represented by three species and one species of the genus *Trachyuropoda* Berlese, 1888 are reported for the first time from Iran. Additionally, we suggest *Urodiaspis pannonicasimilis* Bal and Özkan, 2009 as junior synonym of *Urodiaspis pannonica* and present new morphological data for this species.

Material and methods

Mite specimens were extracted from forest soil and litter by means of Berlese-Tullgren funnels, cleared in Nesbitt's fluid and then mounted in Hoyer's medium.

Morphological observations, measurements and illustrations were made using a compound microscope equipped with differential interference contrast and phase contrast optical systems and a drawing tube (Olympus BX51). Measurements are given in micrometers (μm). Dorsal and female epigynal shields length and width were taken from the anterior to posterior margins along the mid-line and from the lateral margins at the broadest point, respectively.

Mite specimens are deposited in the Acarological Collection, Institute of Science and High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman, Iran (ACISTE).

Results

Family Trachyuropodidae Berlese, 1917

Genus *Trachyuropoda* Berlese, 1888

Type species: *Trachyuropoda festiva* Berlese, 1888.

Trachyuropoda hirschmanni Pecina, 1980 (Fig. 1)

Trachyuropoda hirschmanni Pecina, 1980: p. 373.

Trachyuropoda hirschmanni – Mašán (2001): p. 235; Kontschán (2002): p. 53; Kontschán (2007): p. 51.

Studied materials – Two female specimens collected in soil, litter and cow manure in Shorum Forest, Tonekabon (47° 22' 93" N; 40° 63' 24" E), altitude 1318 m a.s.l., 1 June 2015.

Measurements of the Iranian specimens: Length and width of female idiosoma 729–741, 470–482, respectively.

Distribution – Europe (Pecina 1980; Mašán 2001; Kontschán 2002, 2007), and Iran (this study).




Figure 1. *Trachyuropoda hirschmanni* – Female body (dorsal view).

Genus *Leonardiella* Berlese, 1903

Type species: *Uropoda canestriniana* Berlese, 1891.

***Leonardiella riccardiana* (Leonardi, 1895)**

Note. This species has been reported from Kerman, Iran (Arjomandi and Kazemi 2014) as *Trachyuropoda riccardiana* (Leonardi, 1895), but the species currently transferred to the genus *Leonardiella*, and this genus can be easily distinguished from other genera of Trachyuropodidae by the following morphological characters: idiosoma subtriangular or pentagonal, anterior area of marginal shield broad, and opisthogastric region of ventral idiosoma with a pair of deep furrows (Kontschán 2015).

Genus *Urojanetia* Berlese, 1903

Type species: *Uropoda coccinea* Michael, 1891.

***Urojanetia excavata* (Wasmann, 1899) (Fig. 2)**

Glyphopsis coccinea Wasman var. *excavata* Wasmann, 1899: p. 166.

Urojanetia excavata – Balogh (1938): p. 108; Kontschán (2007): p. 47; Kontschán (2013): p. 111.

Trachyuropoda excavata – Pecina (1980): p. 369; Mašán (2001): p. 235.

Studied material – One male specimen from soil and litter in the Botanical Garden of Nowshahr (36° 38' N; 51° 30' E), altitude -13 m a.s.l., 15 September 2015.

Measurements of the Iranian specimen – Length and width of male idiosoma, respectively, 823, 576.

Distribution – Europe, former USSR (Pecina 1980; Kadite and Petrova 1977; Mašán 2001; Kontschán 2005, 2013), and Iran (this study).

Family Urodiaspididae Trägårdh, 1944**Genus *Urodiaspis* Berlese, 1916**

Type species: *Notaspis tectus* Kramer, 1876.

***Urodiaspis tecta* (Kramer, 1876) (Fig. 3)**

Type species: *Notaspis tectus* Kramer, 1876: p. 79.

Urodiaspis tecta – Berlese (1916): p. 25; Pecina (1980): p. 426; Kadite and Petrova (1977): p. 657; Karg (1989): p. 125; Błoszyk (1999): p. 148; Mašán (2001): p. 187; Bal and Özkan (2007): p. 16; Kontschán (2013): p. 11.

Studied materials – Two female specimens from soil and litter in Arasbaran Forest (38° 56' 25" N; 46° 45' 31" E), altitude 1380 m a.s.l., 20 June 2015.

Measurements of Iranian specimens – Length and width of female idiosoma,

respectively, 808–828, 624–638; genital shield length 213–217 and width 132–142.




Figure 2. *Urojanetia excavata* – Male body (dorsal view).

Distribution – Europe, Turkey, former USSR (Pecina 1980; Kadite and Petrova 1977; Karg 1989; Mašán 2001; Fend'a and Košel 2004; Bal and Özkan 2007; Kamczyc and Gwiazdowicz 2009; Kontschán 2013), and Iran (this study).




Figure 3. *Urodiaspis tecta* – Female body (ventral view).

***Urodiaspis pannonica* Willmann, 1951 (Figs. 4–7)**

Urodiaspis pannonica Willmann, 1951: p. 126.

Urodiaspis pannonica – Berlese (1916): p. 25; Pecina (1980): p. 426; Kadite and Petrova (1977): p. 654; Karg (1989): p. 125; Błoszyk (1999): p. 152; Mašán (2001): p. 187; Kontschán (2013): p. 11.

Urodiaspis pannonicasimilis **syn. nov.** – Bal and Özkan (2009): p. 334.



Figures 4–5. *Urodiaspis pannonica* (female) – 4. Peritreme; 5. Genital shield region.

Studied materials – Two female specimens from soil and litter in Arasbaran Forest (38° 53' 65" N; 46° 47' 64" E), altitude 1700 m a.s.l., 10 July 2015; three females from rotten woods, in Arasbaran Forest (38° 54' 67" N; 46° 47' 91" E), altitude 1250 m a.s.l., 8 August 2015.

Measurements of Iranian specimens – Length and width of female idiosoma, 504–530, 317–350 respectively; genital shield length 132–146 and width 70–80.

Distribution – Europe, former USSR, Turkey (Pecina 1980; Kadite and Petrova 1977; Karg 1989; Mašán 2001; Bal and Özkan 2009; Kontschán 2013), and Iran (this study).

Note. Kadite and Petrova (1977) stated that sternal setae *st5* are inserted in soft cuticle, but our examination of dissected specimens indicates that these setae are inserted on minute and narrow platelets laterad to sternal shield (Figs. 5, 7) which has been confirmed by the examination of European specimens (personal communication of Peter Mašán with the senior author, and the senior author personal observation of *U. pannonica* collected in Slovakia).

Bal and Özkan (2009) described *U. pannonicasimilis* from Turkey and although mentioned that it was very similar to *U. pannonica* they diagnosed the new species based on low value morphological differences and zoogeographical distribution. In comparison between the original description of *U. pannonicasimilis* with female specimens of *U. pannonica* collected in northwestern Iran (near Turkey), and some females and one deutonymph of *U. pannonica* collected in Slovakia, and also the original description of *U. pannonica* and some other related papers, we could not find any diagnostic character to separate these species. Therefore, we suggest *U. pannonicasimilis* as junior synonym of *U. pannonica*.



Figure 6. *Urodiaspis pannonica* – Female body (ventral view).




Figure 7. *Urodiaspis pannonica* (female) – Details of posterior region of genital shield.

Acknowledgments

We wish to thank Peter Mašán for his kind help in providing some information about European specimens of *U. pannonica* and sending some Slovakian specimens of *U. pannonica*. This paper was founded by a grant (No. 95/583) from Institute of Science and High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman Iran.

References

- Arjomandi, E., Kazemi, S. & Afshari, A. (2013) Fauna and diversity of the manure-inhabiting Mesostigmata (Acari) in Kerman County, South Eastern Iran. *Persian Journal of Acarology*, 2(2): 253–263.
- Arjomandi, E. & Kazemi, S. (2014) Edaphic mesostigmatic mites (Acari: Mesostigmata) fauna in parks and landscapes of Kerman City. *The Third Insect Pest Management Conference, Kerman, Shahid Bahonar University of Kerman, Kerman, Iran*, pp. 243–251.
- Arjomandi, E. & Kazemi S. (2016) First report of a genus and species of the family Dinychidae (Acari: Mesostigmata: Uropodoidea) from Iran. *Journal of Entomological Society of Iran* (In press).
- Bal, D.A. & Özkan, M. (2007) Re-description of *Urodiaspis tecta* (Kramer, 1876) (Acarina: Mesostigmata: Uropodina). *Munis Entomology & Zoology*, 2(1): 15–24.
- Bal, D.A. & Özkan, M. (2009) A new uropodine mite of the genus *Urodiaspis* Berlese, 1916 (Acarina, Mesostigmata) from Turkey. *Munis Entomology & Zoology*, 4(2): 333–339.

- Balogh, J. (1938) Magyarország hangyabolyban élő atkáiról I. *Folia entomologica hungarica*, 3: 106–109.
- Beaulieu, F., Dowling, A.P.G., Klompen, H., de Moraes, & Walter, D.E. (2011) Superorder Parasitiformes Reuter, 1909. In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148: 123–128.
- Berlese, A. (1916) Centuria prima di Acari nuovi. *Redia*, 12: 19–67.
- Błoszyk, J. (1999) *Geographical and ecological variability of mites of the cohort Uropodina (Acari: Mesostigmata) in Poland. I. Uropodina mites of oak-hornbeam forests (Carpinion betuli)*. Wydawnictwo Kontekst, Poznan, Poland, 245 pp.
- Błoszyk, J., Adamski, Z., Napierała, A. & Dylewska, M. (2004) Parthenogenesis as a life strategy among mites of the suborder Uropodina (Acari: Mesostigmata). *Canadian Journal of Zoology*, 82: 1503–1511.
- Fend'a, P. & Košel, V. (2004) Mites (Acarina: Mesostigmata) inhabiting caves of the Belianske Tatry Mts (Northern Slovakia). *Biologia*, 59 (supplementary 15): 35–40.
- Halliday, R.B. (2015) Catalogue of genera and their type species in the mite Suborder Uropodina (Acari: Mesostigmata). *Zootaxa*, 3972 (2): 101–147.
- Hirschmann, W. & Zirngiebl-Nicol, I. (1965) Gangsystematik der Parasitiformes, Teil 9, Uropodiden Bestimmungstabellen von 300 Uropodiden-Arten (Larven, Protonymphen, Deutonymphen, Weibchen, Männchen). *Acarologie. Schriftenreihe für Vergleichende Milbenkunde*, 8: 1–33.
- Kadite, B.A. & Petrova, A.D. (1977) Cohort Uropodina Kramer, 1881, Family Uropodidae, Berlese, 1892. In: Ghilyarov, M.S. & Bregetova, N.G. (Eds.), *Key to the Soil Inhabiting Mites: Mesostigmata*. Nauka, Leningrad, pp. 632–691.
- Kamczyc, J. & Gwiazdowicz, D. (2009) Soil mites (Acari: Mesostigmata) from Szczeliniec Wielki in the Stolowe Mountains National Park (SW Poland). *Biological Letter*, 46 (1): 21–27.
- Karg, W. (1989) *Acari (Acarina), Milben Unterordnung Parasitiformes (Anactinochaeta) Uropodina Kramer, Scheldkrötenmilben*. Gustav Fischer Verlag, 203 pp.
- Kazemi, S. & Abolghasemi, S. (2016) New species and records of Uropodina mites from Iran (Acari, Mesostigmata). *Zookeys*, 600: 25–34.
- Kazemi, S. & Kontschán, J. (2007) A review of the mite Uropodina mite fauna (Acari: Mesostigmata) of Iran and description of two new species. *Zoology in the Middle East*, 42: 79–82.
- Kazemi, S. & Kontschán, J. (2014) Three new records of the superfamily Uropodoidea (Acari: Mesostigmata: Uropodina) from Iran, and a key to the known Iranian species of Uropodoidea. *Persian Journal of Acarology*, 3(1): 9–16.
- Kazemi, S. & Rajaei, A. (2013) An annotated checklist of Iranian Mesostigmata (Acari), excluding the family Phytoseiidae. *Persian Journal of Acarology*, 2(1): 63–158.
- Kontschán, J. (2002) The first record of five *Trachyuropoda* species (Acari: Uropodina) from Hungary. *Opuscula Zoologica Budapest*, 34: 51–53.
- Kontschán, J. (2007) Trachyuropodid mites of the Carpathian Basin (Acari Uropodina: Trachyuropodidae). *Opuscula Zoologica Budapest*, 36: 43–56.
- Kontschán, J. (2013) Uropodina mites of the Balkan Peninsula (Acari: Mesostigmata). *Opuscula Zoologica Budapest*, 44 (1): 97–131.

- Kontschán, J. (2015) *Trachyibana sarawakiensis* gen. nov., sp. nov., a remarkable new genus and species from Malaysia (Acari: Uropodina: Trachyuropodidae). *Zootaxa*, 3915 (2): 272–278.
- Kontschán, J. & Hajizadeh, J. (2013) First record of the family Macrodirinychidae Hirschmann, 1975 (Acari: Uropodina) from Iran. *Persian Journal of Acarology*, 2 (1): 189–193.
- Kontschán, J. & Starý, J. (2013) Three new *Trachyuropoda* (Acari: Uropodina: Trachyuropodidae) species from the Neotropical region. *Turkish Journal of Zoology*, 37: 7–14.
- Kramer, P. (1876) Zur Naturgeschichte einiger Gattungen aus der Familie der Gamasiden. *Archiv für Naturgeschichte*, 42: 46–105 + plates IV–V (Not seen).
- Krantz, G.W. & Ainscough, B.D. (1990) Acarina: Mesostigmata (Gamasida). In: Dindal, D.L. (Ed.), *Soil Biology Guide*. Wiley & Sons, New York, USA, pp. 583–631.
- Lindquist, E.E., Krantz, G.W. & Walter, D.E. (2009) Order Mesostigmata. In: Krantz, G.W. & Walter, D.E. (Eds.) *A Manual of Acarology*, 3rd Edition. Texas Tech University Press, Lubbock, USA. pp. 124–232.
- Mašán, P. (2001) Mites of the cohort Uropodina (Acarina, Mesostigmata) in Slovakia. *Annotationes Zoologicae et Botanicae*, 223: 1–321 (In Slovak).
- Pecina, P. (1970) Contribution to the knowledge of Uropodidae (Berlese, 1892) Hirschmann et Zirngiebl-Nicol, 1964, of the environs of Prague (Acari, Mesostigmata). *Acta Universitatis Carolinae-Biologica*, 1968: 417–434.
- Pecina, P. (1980) Additional data on several Czechoslovak members of subfamily Trachyuropodinae Berlese, 1918 (Uropodidae, Mesostigmata). *Acta Universitatis Carolinae Biologica*, 1978: 357–388.
- Wasmann, E. (1899) Weitere Nachträge zum Verzeichniss der Ameisengäste von holländisch Limburg. *Tijdschrift voor Entomologie*, 42: 158–171 (Not seen).
- Willmann, C. (1951) Untersuchungen über die terrestrische Milbenfauna im pannonischen Klimagebiet Österreichs. *Sitzungsberichten der Österreich Akademie der Wissenschaften, mathematisch-naturwissenschaftlichen Klasse, Abt. 1*, 160: 91–176.
- Wiśniewski, J. & Hirschmann, W. (1993) Gangsystematik der Parasitiformes, Teil 548. Katalog der Ganggattungen, Untergattungen, Gruppen und Artender Uropodiden der Erde. *Acarologie. Schriftenreihe für Vergleichende Milbenkunde*, 40: 1–220.

Received: 9 June 2016

Accepted: 25 June 2016

Published: 15 July 2016

COPYRIGHT



Kazemi *et al.* Persian Journal of Acarology is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

کنه‌های خانواده‌های *Urodiaspididae* و *Trachyuropodidae* Berlese از ایران (Acari: Mesostigmata: Uropodina) Trägårdh

شهرز کاظمی*، مجتبی محمد دوستار شرف و سپیده صابری

گروه تنوع زیستی، پژوهشگاه علوم و تکنولوژی پیشرفته و علوم محیطی، دانشگاه تحصیلات تکمیلی صنعتی و فناوری پیشرفته، کرمان، ایران؛ رایانامه: shahroozkazemi@yahoo.com، saberi20202s@gmail.com، mojtaba.doostar@gmail.com، sh.kazemi@kgut.ac.ir

* نویسنده مسئول

چکیده

در اینجا دو جنس *Urodiaspis* Berlese و *Urojanetia* Berlese و چهار گونه *Urodiaspis pannonica* Willmann, 1951، *Trachyuropoda hirschmanni* Pecina, 1980 و *Urodiaspis tecta* (Kramer, 1876) و *Urojanetia excavata* (Wasmann, 1899) برای نخستین بار از ایران گزارش می‌شوند. افزون بر این، گونه *Urodiaspis pannonicasimilis* Bal and Özkan, 2009 به عنوان مترادف گونه *U. pannonica* معرفی و صفات ریخت‌شناسی جدیدی برای آن ارائه می‌شود.

واژگان کلیدی: فون؛ Monogynaspida؛ مترادف جدید؛ کنه‌های خاکزی؛ Uropodoidea.

تاریخ دریافت: ۱۳۹۵/۳/۲۰

تاریخ پذیرش: ۱۳۹۵/۴/۵

تاریخ چاپ: ۱۳۹۵/۴/۲۵