



Age-related Changes in some Blood Parameters of Ostrich (Short Communication)

Khodaei Motlagh, M¹ & Masoumi R²

¹Department of Animal Science, Faculty of Agriculture and Natural Resource, Arak University, Arak 38156-8-8349, Iran

²Department of Animal Science, Faculty of Agriculture, University of Zanjan, Zanjan, Iran

Poultry Science Journal 2016, 4(2): 167-171

Keywords

Age
Ostrich
Blood parameter

Corresponding author

Mahdi Khodaei
mmotlagh2002@gmail.com

Article history

Received: December 16, 2015

Revised: May 30, 2016

Accepted: November 14, 2016

Abstract

The aim of this study was to investigate blood parameters of 4-month old blue-neck male ostriches (*Struthio camelus*) after feeding a diet containing 3% sunflower oil for two months. Blood samples were collected from the wing vein of the ostriches at the beginning and end of the study. The plasma was harvested and analyzed to measure cholesterol, triglyceride, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, very low-density lipoprotein cholesterol, total protein, albumin, total immunoglobulin, and the activities of aspartate amino transferase and alanine amino transferase enzymes. From days 0 to 60, concentrations of high-density lipoprotein cholesterol decreased ($P < 0.05$) while very low-density lipoprotein cholesterol and triglyceride increased significantly ($P < 0.05$). Other blood parameters did not change significantly. The results of this study showed that in ostrich, with increasing age from four to six months of age, very low-density lipoprotein cholesterol and triglyceride concentrations increased whereas other factors did not.

Introduction

The ostrich (*Struthio camelus*) is a member of the ratite family of flightless birds and is native to semi-arid and desert areas of Africa (Smit, 1963). Ostrich farming is an activity that has been expanding worldwide for nearly 130 years (Feitosa *et al.*, 2013). In Iran, this economic activity is still very recent, considering that only in 1999 the first ostriches arrived for commercial purposes (Hajibabaei *et al.*, 2005) and present ostrich farms are distributed in all regions and climates of the country, especially in the central regions (Ghasemi *et al.*, 2012). Analysis of blood biochemistry is a tool for the early diagnosis of nutritional and metabolic disorders before the emergence of more serious symptoms (Miranda

et al., 2008). Physiological parameters in blood are known to be influenced by different factors include age, disease, management, and nutrition.

There is little information about blood parameters of ostrich. Evaluation of the ratite hematology and clinical chemistry has become increasingly important in veterinary care. The aim of this study was to investigate blood parameters of 4-month old blue-neck male ostriches (*Struthio camelus*) after feeding a diet containing 3% sunflower oil for two months.

Material and Methods

Animal and diet

The study was carried out on six, 4-month old ostriches with an average weight of 43.3 ± 2.1 kg

in Arak University (Markazi province, Iran). The ostriches were fed *ad libitum* with a standard mash diet which has been formulated according to the recommendations reported by Cooper and Horbanczuk (2004) for 60 days (Table 1). The ostriches were kept in a commercial farm with 150 m² open space and 100 m² covered space and had free access to fresh water throughout the

study. The experiment was approved by the animal group committee of the agriculture faculty of Arak University (Approval No: 2013-07/02 -366). The ostriches were under veterinary care by a veterinarian. They were closely checked twice each day for any signs of illness or behavioral changes throughout the study and did not show any clinical signs of any disease.

Table 1. Ingredients and chemical composition of diet

Ingredients	(%)
Corn, Grain	37.40
Alfalfa Meal (20% CP)	28.05
Soybean Meal (44% CP)	27.58
Dicalcium phosphate	2.30
Sunflower oil	3.00
Limestone	0.69
Common salt	0.31
Vitamin premix ¹	0.23
Mineral premix ²	0.23
DL-Methionine	0.14
L-Lysine HCl	0.07
<i>Chemical composition</i>	
ME (Kcal/kg)	2600
CP (%)	16

¹Vitamin premix supplied per kg of diet: vitamin A: 2.6 mg; vitamin D₃: 0.06 mg; vitamin E: 16 mg; vitamin K₃: 2 mg; thiamine: 1.8 mg; riboflavin: 6.2 mg; pantothenic acid: 10 mg; pyridoxine: 3 mg; cyanocobalamin: 0.015 mg; niacin: 28 mg; biotin: 0.1 mg; folic acid: 1.5 mg; choline chloride: 240 mg; antioxidant: 100 mg.

²Mineral premix supplied per kg of diet: Fe: 40 mg; Mn: 100 mg; Zn: 100 mg; Cu: 10 mg; I: 1 mg; Se: 0.2 mg.

Blood was collected from the ostriches at the beginning and the end of study, each time after 12 hrs of fasting. Blood samples were collected from the wing vein in heparinized tubes and transferred to the central laboratory of Arak city to prepare plasma by centrifugation at 2500 × g for 15 min. Plasma was stored at -21°C until analysis. Twice each day, ostriches were closely observed for any signs of illness or behavioral changes.

Growth performance

All ostriches were weighed in the morning, before drinking and feeding. Feed intake was calculated by measuring the amount of feed offered and residue left after 24 hrs. Feed conversion ratio (FCR) was calculated by dividing feed intake (FI) with weight gain.

Plasma biochemical analyses

Biochemical parameters consisted of total protein, albumin protein, total immunoglobulin, aspartate aminotransferase (AST), alanine aminotransferase (ALT), triglyceride, total cholesterol, high-density lipoprotein cholesterol

(HDL-C), very-low-density lipoprotein cholesterol (VLDL-C) and low-density lipoprotein cholesterol (LDL-C). All analyses were made using commercial kits (Pars Azmoon, Iran) and a standard auto analyzer apparatus (Biotechnica, BT-3000, Rome, Italia). The concentration of VLDL-C was calculated by dividing plasma triglyceride by five. The LDL-C value was calculated using the formula: LDL-C = Total cholesterol - HDL-C - VLDL-C.

Statistical analysis

The data were checked for normal distribution using Kolmogorov-Smirnov method before analysis. Paired t-test was performed using SAS software (2001) for data analysis. A value was considered to be statistically significant if the *P*-value was less than 0.05.

Results and Discussion

Growth performance of ostriches is reported in Table 2. Average daily feed intake (ADFI), water consumption per day, feed conversion ratio (FCR), and weight gain were different from that estimated by Mushi *et al.* (1998). Waugh *et al.*

(2006) reported the average daily weight gain (ADG) as 320 g/day for young ostriches which is slightly more than finding of present experiment (271.3 g/day). However, Cooper (2005) reviewed growth in the ostriches. He reported based on the previous researches that ostrich chicks have a high feed conversion to

body mass ratio. This ratio is 2:1 at hatching to 4 months old and 3.8:1 at 4-6 months old. Also, Kreibich and Sommer (1995) explained that FCR ranged from 1.4:1 to 1.6:1 for younger birds (four to six months old) while older birds ranged from 4:1 to 6:1.

Table 2. Growth performance of ostriches (mean \pm SE)

Initial Weight (kg)	Final Weigh (kg)	Gain (kg)	ADG (g)	FCR	ADFI (g)
43.3 \pm 2.1	59.6 \pm 1.1	16.3 \pm 1.01	271.3 \pm 11	2.648 \pm 0.6	718 \pm 25

No significant differences were observed in blood parameters of ostriches between the beginning and end of the study, except for triglyceride, HDL-C and VLDL-C concentrations (Table 3). These findings showed that ostriches at the end of the study had significantly higher plasma concentrations of VLDL-C, triglyceride, and HDL-C.

Total protein and total immunoglobulin values tended to increase by the end of study but this trend was insignificant. The average value of total protein for the two ages (4 and 6 months) in this study was 4.89 g/dL, which was similar to values reported by Bell and Freeman (1971), Quintavalla *et al.* (2001), and Penninks *et al.* (2012). The values of total immunoglobulin in this investigation were observed to be 0.88 - 0.93 g/dL. However, higher values have previously been reported (Ahmed *et al.*, 2006); Cantu-Martinez *et al.*, 2010; Omid and Ansari nik, 2013), as well as lower (Selvan *et al.*, 2012). Similar to the results of Durgun *et al.* (2005), the

effect of age on total immunoglobulin in this study was not significant, but, Quintavalla *et al.* (2001) and Penninks *et al.* (2012) reported that total protein and total immunoglobulin increased with age in ostriches until 24-36 months. There was no significant difference in serum albumin concentration between the two age groups. The range of blood albumin concentration was 4.27 \pm 0.46 to 4.53 \pm 0.26 g/dL in the two age groups which were similar to findings from Miranda *et al.* (2008), Bonadiman *et al.* (2009), and Omid and Ansari nik (2013). In contrast, higher concentrations of blood albumin were observed by Miranda *et al.* (2008) and Zahid *et al.* (2003).

Age had no significant effect on LDL-C concentration. HDL-C concentration in the two age groups was significantly affected with 4 month old group having 50.95 \pm 1.20 mg/dL and the six month old group having 47.08 \pm 1.13 mg/dL. HDL-C concentration decreased with an increase in age group.

Table 3. Effect of ostrich age on plasma biochemical parameters at beginning and end of the study

Variable	Day 0	Day 60	P-value
Triglyceride (mg/dL)	99.33 \pm 3.70 ^a	114.95 \pm 3.50 ^b	0.033
Cholesterol (mg/dL)	101.78 \pm 7.74	134.75 \pm 13.44	0.125
HDL-C (mg/dL)	50.95 \pm 1.20 ^a	47.08 \pm 1.13 ^b	0.041
LDL-C (mg/dL)	31.66 \pm 8.39	63.28 \pm 14.64	0.150
VLDL-C (mg/dL)	19.86 \pm 0.74 ^a	22.98 \pm 0.70 ^b	0.033
Total Protein (g/dL)	4.67 \pm 0.16	5.11 \pm 0.21	0.252
Albumin (g/dL)	4.27 \pm 0.46	4.53 \pm 0.26	0.710
Total Immunoglobulin (g/dL)	0.88 \pm 0.22	0.93 \pm 0.08	0.853
ALT (IU/L)	32.95 \pm 1.84	33.45 \pm 1.72	0.468
AST (IU/L)	34.65 \pm 1.48	32.73 \pm 0.84	0.284

Triglyceride concentration was significantly affected by age which was dissimilar to the findings of Moniello *et al.* (2006). The concentrations of triglyceride in four and six

month old ostriches were 99.33 \pm 3.00 and 114.95 \pm 3.50 mg/dL, respectively. Lower values were observed by Durgun *et al.* (2005), Moniello *et al.* (2006), and Omid and Ansari nik (2013).

Age had no significant effect on blood cholesterol concentration which was similar to the findings of Palomeque *et al.* (1991) and Durgun *et al.* (2005). Cholesterol concentration increased with (101.78±7.74 mg/dL at four months *vs.* 134.75±13.44 mg/dL at six months). Lower values were recorded by Palomeque *et al.* (1991), Durgun *et al.* (2005), and Selvan *et al.* (2012), while a higher value was reported by Omid and Ansari nik (2013).

There were no significant differences in activities of AST and ALT. Their averages were 33.69 and 33.02 IU/L for 4 and 6 month olds, respectively. In other studies, the AST values were reported 131 to 372.2 IU/L (Levy *et al.*, 1989; Angel, 1996), while ALT values were reported from 2 to 26.62 IU/L (Levy *et al.* 1989; Palomeque *et al.* 1991). This result was in disagreement with

the finding reported by Cilliers *et al.* (1995). This variation can include sensitivity of laboratory instruments and age of ostriches.

Conclusion

In summary, as ostriches grew from four to six months of age, VLDL-C and triglyceride concentrations increase while other blood parameters including total protein, albumin, total immunoglobulin, of AST and ALT are not changed.

Acknowledgement

The author gratefully acknowledges the financial support of Arak University and the kind help of Animal Science Department, especially Ali Hassani and Sepehr Nematpour for completing this study.

Reference

- Ahmed NHS, Ibrahim MT, Mahmoud ZN & Ali OS. 2006. Normal concentrations of some blood constituents in ostrich (*Struthio camelus*). Journal of Science and Technology, 7: 1-6. [\[Link\]](#)
- Angel CR. 1996. Serum chemistries and vitamin D metabolites in ostriches, emus, rheas and cassowaries. In Proceedings of an International Conference: Improving Our Understanding of Ratites in a Farming Environment, Manchester, pp: 122-124.
- Bell DJ & Freeman BM. 1971. Physiology and Biochemistry of the Domestic Fowl. Vol. 2. Academic Press, London, New York.
- Bonadiman SF, Stratievsky GC, Machado JA, Albernaz AP, Rabelo GR & DaMatta RA. 2009. Leukocyte ultrastructure, hematological and serum biochemical profiles of ostriches (*Struthio camelus*). Poultry Science, 88: 2298-2306. [\[Link\]](#)
- Cantu-Martinez AM, Silva-Paez MDL, Riojas-valdes VM, Santoyode-Estefano FA & Garza-Zermeno MV. 2010. Blood biochemical profile in ostrich from northeast Mexico. Journal of Animal and Veterinary Advances, 9: 226-228. [\[Link\]](#)
- Cilliers SC, Du Preez JJ, Maritz JS & Hayes JP. 1995. Growth curves of ostriches (*Struthio camelus*) from Oudtshoorn in South Africa. Animal Science, 61: 161-164. [\[Link\]](#)
- Cooper RG, & Horbanczuk JO. 2004. Ostrich nutrition: a review from a Zimbabwean perspective. Revue scientifique et technique International Office of Epizootics, 23: 1033-1042. [\[Link\]](#)
- Cooper RG. 2005. Growth in the Ostriches (*Struthio camelus* var. *domesticus*). Animal Science Journal, 76: 1-4. [\[Link\]](#)
- Durgun Z, Keskin E, Col R & Atalay B. 2005. Selected haematological and biochemical values in ostrich chicks and growers. Archiv fur Geflugelkunde, 69: 62-66. [\[Link\]](#)
- Feitosa TF, Vincius Longo R, Rodrigues A, Braga FR, Dantas ES, Vieira VD & Bezerra LA. 2013. Anthelmintic efficacy of pumpkin seed (*Cucurbita pepo* Linnaeus, 1753) on ostrich gastrointestinal nematodes in a semiarid region of Paraíba State, Brazil. Tropical Animal Health Production, 45: 123-127. [\[Link\]](#)
- Ghasemi HA, Kazemi-Bonchenari M, Khaltabadi-Farahani AH & Khodaei Motlagh, M. 2012. The effect of feeding rations with different ratios of concentrate to alfalfa hay on blood hematological and biochemical parameters of farmed ostriches (*Struthio camelus*). Tropical Animal Health Production, 45: 1635-40. [\[Link\]](#)
- Hajibabaei A, Mosavi SM & Vahedi H. 2005. Ostrich breeding in Iran, Proceedings of the 3rd International Ratite Science Symposium of the World's Poultry Science Association (WPSA), Madrid, Spain, 14th-16th October, 359-361.
- Levy A, Perelman B, Waner T, Van Grevenbroek M, Van Creveld C & Yagil R. 1989. Reference

- blood chemical values in ostriches (*Struthio camelus*). American Journal of Veterinary Research, 50: 1548-1550. [\[Link\]](#)
- Kreibich A & Sommer M. 1995. Ostrich Farm Management. 1st Ed. Landwirtschaftsverlag GnbH. 222 Pages. [\[Link\]](#)
- Miranda RL, Mundim AV, Costa AS, Guimarães RV & Silva FOC. 2008. Serum biochemistry of 4-day-old ostriches (*Struthio camelus*). Pesquisa Veterinária Brasileira, 8: 423-426. [\[Link\]](#)
- Moniello G, Bovera F, Riu ND, Piccolo G, Pinna W & Meo CD. 2006. Gender effect on the metabolic profile of ostriches (*Struthio Camelus domesticus*). Italian Journal of Animal Science, 5: 229-235. [\[Link\]](#)
- Mushi EZ, Binta MG, Chabo RG, Lsa JFW & Modisa L. 1998. Serum biochemical values of farmed Ostrich (*Struthio camelus*) in Botswana. Onderstepoort Journal Veterinary Research, 65: 189-193. [\[Link\]](#)
- Omidi A & Ansari nik H. 2013. Selected biochemical values of yearling African blue neck ostriches (*Struthio camelus*) in Iran. Comparative Clinical Pathology, 22: 601-604. [\[Link\]](#)
- Palomeque J, Pintò D & Viscor G. 1991. Haematological and blood chemistry values of the Masai ostrich (*Struthio camelus*). Journal of Wild Life Disease, 27: 34-40. [\[Link\]](#)
- Penninks AH, van Mierlo GJD, Kuper F, Snel CJ, Ganderup NC & Wolterbeek APM. 2012. Juvenile immunodevelopment in mini pigs, Pediatric Nonclinical Drug Testing: Principles, Requirements, and Practices, 231-254. [\[Link\]](#)
- Quintavalla F, Bigliardi E & Bertoni P. 2001. Blood biochemical baseline values in the ostrich (*Struthio camelus*). Annali Facoltà Medicina Veterinaria, 21: 61-71. [\[Link\]](#)
- Selvan ST, Gopi H & Vairamuttu P. 2012. Haematology and blood chemistry values in Juvenile Ostriches (*Struthio camelus*) under sub-tropical conditions of India. Research Journal of Poultry Science, 5: 60-63. [\[Link\]](#)
- Smit DJ. 1963. Ostrich farming in Little Karroo. Bulletin No. 358, Department of Agricultural Technical Services, Pretoria. [\[Link\]](#)
- Waugh EE, Aganga AA, Seabo D, Omphile UJ, and Tsopito CM. 2006. Growth rate and feed conversion rate of ostriches fed ration with or without grit in Botswana. International Journal of Poultry Science, 5: 470-473. [\[Link\]](#)
- Zahid A, Yükses N, Altuğ N, Kaya A & Baydaş B. 2003. Studies on Some Haematological and biochemical parameters in clinically healthy ostriches. Turkish Journal of Veterinary and Animal Sciences, 27: 161-163. [\[Link\]](#)



تغییرات وابسته به سن در برخی از فراسنجه‌های خونی شترمرغ (مقاله کوتاه)

Khodaei Motlagh, M¹ & Masoumi R²

¹ گروه علوم دامی، دانشکده کشاورزی و منابع طبیعی، دانشگاه اراک، اراک، ایران

² گروه علوم دامی، دانشکده کشاورزی، دانشگاه زنجان، زنجان، ایران

Poultry Science Journal 2015, 4(2): 167-171

DOI: 10.22069/PSJ.2016.10081.1166

چکیده

هدف از این مطالعه بررسی فراسنجه‌های خون در شترمرغ‌های چهار ماهه پس از تغذیه ۳ درصد روغن آفتابگردان در مدت دو ماه بود. در آغاز و پایان آزمایش از زیر بال پرندگان خونگیری انجام شد. فراسنجه‌های خون شامل: کلسترول، تری گلیسرید، HDL-کلسترول، LDL-کلسترول، VLDL-کلسترول، پروتئین کل، آلبومین، ایموگلوبولین کل و فعالیت آنزیم‌های آسپارات آمینوترانسفراز و آلانین آمینوترانسفراز اندازه‌گیری شد. از روز صفر تا ۶۰ آزمایش غلظت HDL-کلسترول کاهش یافت ($P < 0.05$). در حالی که غلظت VLDL-کلسترول و تری-گلیسرید به طور معنی‌داری افزایش یافت ($P < 0.05$). سایر فراسنجه‌های خون به‌طور معنی‌داری تغییر نکردند. نتیجه این مطالعه نشان داد که با افزایش سن شترمرغ‌ها از چهار ماه به شش ماه، میزان VLDL-کلسترول و تری گلیسرید افزایش یافت در حالیکه سایر فراسنجه‌ها تغییری نکرد.

کلمات کلیدی

سن
شتر مرغ
فراسنجه‌های خون

نویسنده مسئول

Mahdi Khodaei
mmotlagh2002@gmail.com

تاریخچه مقاله

دریافت: ۱۶ دسامبر ۲۰۱۵

ویرایش: ۳۰ می ۲۰۱۶

پذیرش: ۱۴ نوامبر ۲۰۱۶

Please cite this article as: Khodaei Motlagh, M & Masoumi R. 2016. Age-related Changes in Some Blood Parameters of Ostrich (Short Communication). Poul. Sci. J. 4(2): 167-171.

© 2016 PSJ. All Rights Reserved