

# Journal of Drought and Climate change Research (JDCR)



Spring 2023, Vol.1, No.1, pp 1 - 18

## The impact of climate change on severity, duration, and magnitude of drought using

### SPI and RDI in the Semnan region

Mahdi Delghandi<sup>1\*</sup>, Sanaz Joorablou<sup>2</sup>, Zahra Ganji Nowroozi<sup>3</sup>

- 1. Assistant Professor, Department of Water and Soil, Faculty of Agricultural Engineering, Shahrood University of Technology, Shahrood, Iran.
- 2. Ph.D. Student, Faculty of Civil Engineering, Shahrood University of Thecnology, Shahrood, Iran.
- 3. Assistant Professor, Department of Water and Soil, Faculty of Agricultural Engineering, Shahrood University of Technology, Shahrood, Iran.

\*Corresponding Author: delghandi@shahroodut.ac.ir

Keywords:	Extended abstract
Climate Change, Emission	Introduction
Scenarios, Drought, RDI, SPI.	In recent years human activities have induced increases in atmospheric carbon dioxide
	(CO2). Increases in [CO2] caused global warming and climate change (IPCC, 2001;
	IPCC, 2007). Climate change is anticipated to cause negative consequences, some of
	which will have directly obvious impacts, such as intense precipitation, surface water
	resources and flooding while some will have long-term impacts such as sea level rise,
	diseases and droughts. Globally, droughts have a devastating impact on agriculture,
	the economy, and society. Drought monitoring is an essential component of drought
	risk management. Drought indices are indispensable tools to detect, monitor, and eval-
	uate drought events. These indices were classified as meteorological drought indices.
Received: Dec/18/2022	The Reconnaissance Drought Index (RDI) and standardized precipitation index (SPI)
	are the most popular drought assessment Indices.
	As global warming cascades into the next generation, a change in drought intensity
Revised: Feb/25/2022	and duration as considered by many studies is becoming a threat to humanity. There-
	fore, any measures for mitigation and adaptation require careful assessment of histor-
	ical events and accurate representation of future drought based on new methodolog-
	ical approaches. Since Semnan is located in the arid and semi-arid region, drought
Accepted:	assessment is very important. To be prepared for the negative impacts of drought, it
Mar/28/2023	is essential to predict their properties. Therefore, the aim of this paper is to examin of
	climate change impacts on drought frequency, severity and duration using two drought
	indices (SPI and RDI) for Semnan region located in Iran.

#### How to cite this article:

Delghandi, M., Joorablou, S., & Ganji Nowroozi, Z. (2023). The impact of climate change on severity, duration and magnitude of drought using SPI and RDI in Semnan region. *Journal of Drought and Climate change Research*, 1(1), 1 - 18. 10.22077/JDCR.2023.5909.1004.



Copyright: © 2022 by the authors. Licensee *Journal of Drought and Climate change Research* (JDCR). This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<u>https://creativecommons.org/licenses/by/4.0/)</u>.

#### Material and method

#### Study Area

The synoptic station of Semnan, with an elevation of 1127 meters has longitude of 53°23 ' East, latitude of 35°35 ' North. The annual precipitation at this station is 140 mm, the average Tmin and Tmax is, respectively, 12.5 and 23.8 °C, respectively.

#### Generating climate precipitation scenarios

The outputs from 13 AOGCMs following two emission scenarios RCP8.5 and RCP4.5 were utilized to produce monthly rainfall and temperature climate scenarios. The periods 1971-2000, 2010-2039, 2040-2069 and 2070-2099 were selected as baseline and future periods, respectively. The LARS-WG model was used for spatial and temporal downscaling of temperature and precipitation time series and daily precipitation and temperature scenarios were generated at probability levels of 0.50, 0.75 and 0.90 for the three future periods.

#### Drought indices

#### Standardized precipitation index (SPI) and Reconnaissance drought index (RDI)

For the calculation of SPI and RDI, the long-term data (precipitation for SPI, precipitation and evapotranspiration for RDI) are fitted to a gamma distribution. Drought occurs when SPI and RDI reach a value of -1.0 or less.

#### Evapotranspiration and precipitation deficit (PD)

There are several models for the estimation of reference evapotranspiration  $(ET_{o})$ . In this study, FAO Penman–Monteith (FAO PM) and Hargreves-Samani methods were used to estimate ETo.

#### Drought characteristics

Yevjevich (1967) proposed the Run theory to identify drought parameters (duration, magnitude and severity) which are described in Fig 1. The most key element for extracting these parameters is the threshold level, which may be a constant or a function of time.

#### Fig. 1. Drought characteristics



#### **Results and Discussion**

As a result, the FAO Penman-Monteith method (FAO PM) was found to be more precise than the Hargreves-Samani method for estimating EToTherefore, in this study FAO PM used for estimation of RDI in climate change scenarios.

Temporal changes of SPI and RDI at 1-year timescale, under the risk level of 0.5 are illustrated in Fig 2.

Fig 2. Temporal changes of SPI and RDI at 1-year timescale, under the risk level of 0.5 (a): RCP4.5 (b): RCP8.5



The results revealed that as compared to the RCP4.5 scenario, in the RCP8.5 scenario, there will be more dry and wet periods. Drought duration predicted by RDI index compared to SPI index (at 1-year and 3-month time scales), under both emission scenarios in the future period (2010-2099) has decreased compared to the past period (1965-2009), but the number of droughts has increased. Additionally, increases and decreases in drought magnitude and severity were found at time scales 3-month and annually, respectively. These results provide observational evidence for the increasing risk of droughts as anthropogenic global warming progresses.

Journal of Drought and Climate change Research (JDCR)