

CASE STUDY

Effect of compost and humic acid in mobility and concentration of cadmium and chromium in soil and plant

A. Chaab, A.A. Moezzi*, G.A. Sayyad, M. Chorom

Department of Soil Science, College of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Iran

Received 1 April 2016; revised 21 May 2016; accepted 28 May 2016; available online 1 September 2016

ABSTRACT: The effect of compost and humic acid in mobility and concentration of cadmium and chromium in contaminated soil were investigated. Experiment was carried out with three levels of soil cadmium and chromium and two organic matters (compost and humic acid). The study was performed in a randomized complete block design with 3 replicates. Results indicated that application of organic substances enhanced movement of cadmium and chromium in soil column. Humic acid is more effective than compost on the mobility of cadmium and chromium in soil. Mobility of cadmium and chromium in the lower depths of soil column were increased. Cadmium and chromium concentration in shoots and roots enhanced due to increasing those concentration in soil and application of organic substances. Increase in cadmium in shoots can be attributed to the high mobility of this element in maize plant. Maize root chromium concentration was greater than shoot chromium concentration. Humic acid was more effective than compost as cadmium and chromium concentration in root and shoot was concerned. Low mobility of chromium in plant and accumulation of chromium in roots can be reasons of decreasing of chromium concentration in shoot of plant and its bioaccumulation.

KEYWORDS: Cadmium; Chromium; Compost; Humic acid; Mobility

INTRODUCTION

Heavy metal pollution in soils is a common problem in several countries (John *et al.*, 2009). Today soil pollution with heavy metals is a major environmental problem in the world. This pollution has harmful effects on the soil fauna, which reduces the yield and quality of agricultural products (Raymond and Okieimen, 2011). Physiological and biochemical process in plants were affected by high concentration of heavy metals such as cadmium and chromium (Kim *et al.*, 2004; Strobel *et al.*, 2005). Cadmium is one of the most toxic metals that effects on all biological processes of humans, animals and plants; such as leaf chlorosis, photosynthesis, reduce root growth, decrease absorption and transport of nutrient elements (Gabrijel *et al.*, 2009; Orrono;

Lavado, 2009; Karbassi *et al.*, 2015). The contents of Cd in soils range from 0.06 to 4.3 mg/kg and in plants range from 0.005-0.252 mg/kg (Kabata and Mukherjee, 2007; Wojcik *et al.*, 2005). Chromium in high concentration also affected human health and plant growth. The world soil average content of Cr in soils was 60 mg/kg and it rang in plants from 0.01-0.08 mg/kg (Kalantari *et al.*, 2014; Sanita *et al.*, 2010). Besides soil pH, the content of organic matter affect availability of heavy metals. Organic matter (OM) can bind with heavy metals and controlling their behavior in soils (Park *et al.*, 2011; Salati *et al.*, 2010; Tingqiang *et al.*, 2013; Wyszowski *et al.*, 2013; Karbassi and Pazoki, 2015). Soil organic matter is one of the important soil components that affects soil reactions such as form of ion uptake by plants (Ouni *et al.*, 2014). OM also effect plant growth by increasing in improvement of photosynthesis system, micro and macro element

✉ *Corresponding Author Email: moezzi251@gmail.com
Tel.: +98 916 313 9813; Fax: +98 61 3333 5011

Note: Discussion period for this manuscript open until December 1, 2016 on GJESM website at the "Show Article".

uptake and changing in growth of root cells (Pizzeghello et al., 2013). Some researchers believe that application of OM, such as compost, humic acid and sewage sludge, is a recommend way for mobilization of heavy metals in soils. In many of that experiments the stimulating substances were undoubtedly largely chelated elements, chiefly iron (Khan et al., 2000). They find that application of compost and humic acid in contaminated soils enhanced bioavailability and mobility of metals (Pizzeghello et al., 2013; Topcuoglu, 2012) while others showed that organic substances have the ability to bind metal ions and lead to stabilization of them (Zhang et al., 2013). Thus, the influence of OM such as compost and humic acid on availability and movement of Cd in soil and plant require further investigation. Therefore the research was aimed to investigate the effect of compost and humic acid on the mobility and accumulation of chromium and cadmium in soil and plant. This study has been performed in greenhouse of soil department of Shahid Chamran University of Ahvaz, Iran during April to September 2015.

MATERIALS AND METHODS

Experimental design

The study was carried out in greenhouse of the Shahid Chamran University, using soil columns. Treatments were performed with the level of Chromium (0, 100, 200 mg Cr/kg soil) using $K_2Cr_2O_7$, Cadmium (0, 25, 50 mg cd/kg soil) and using $CdCl_2$ along with two levels of organic matter; compost (0, 40 mg/kg soil) and humic acid (0, 2.5 g). Investigation of these elements and their value were performed due to reports of their contamination in the soils in Ahvaz, Iran (Jafarngadi et al., 2010; Parsafar and Marofi, 2013). Treatments were performed: 1) control: no Cd and Cr (T_0) + no organic substance (M_0); 2) compost (M_1) + T_1 (25 mg cd/kg soil+ 100 mg Cr/kg soil); 3) humic acid (M_2) + T_1 (25 mg cd/kg soil+ 100 mg Cr/kg soil); 4) compost (M_1) + T_2 (50 mg cd/kg soil+ 200 mg Cr/kg soil); 5) humic acid (M_2) + T_2 (50 mg cd/kg soil+ 200 mg Cr/kg soil). The treatments were performed in factorial randomized complete design with 3 replicates. Factorial design of treatments shows in Table.1. The surface of soil was mixed with 40 g compost/kg soil. Humic acid was purchased from the Flvka Institute, as the commercial sample (product No. 35069288) and as described by Hoop (1990). Seeds of plant (single grass 704) were sowed in plastic columns containing 14 kg of

soil. Up to harvest received proper operations. Prior to column filling, soil simultaneously mixed with required amount of Cd and Cr solution then incubated for one month followed by application of NPK as per soil test. Based on molecular weight of used salt for each understanding element, required amount of each salt for every treatment were calculated. The required quantity of each salt for each treatment was dissolved in 500 ml distilled water and sprayed over the soil informally. The soil texture was sandy loam and the general properties was provided in Table.2. At the end of the study, plants were harvested then roots were isolated from soli and washed with deionized water. The maize plant components (shoot and roots) then dried in an oven at 65° C for 72 h and then in followed by powdering the samples.

Determination of heavy metals

Concentration of heavy metals in plant was analyzed by atomic absorption spectroscopy at the following wavelengths: cadmium at 228.8 nm and chromium at 357.9 nm (Mireles, 2004). Cd and Cr concentration in soil column was measured in three depth (0-15, 15-30 and 30-45 cm) as previously described by Chapman and Pratt (1961). The samples were measured against a calibration graph which was obtained with measurements on matrix-compensated standards (Boner and Pohl, 2010).

Transportation index

The transportation index (Ti) was determined by the ratio between shoot and root heavy metal concentration (Marchiol et al., 2004).

$Ti = \text{Cd or Cr concentration in shoot (mg/kg)} / \text{Cd or Cr concentration in roots (mg/kg)}$

Bio accumulation factor

Bio accumulation factor is the ratio of the metal concentration in the plant to its concentration in the soil which is calculated from the following equation: (Parsafar and Marofi, 2013).

$$BAF = M_{pc}/M_{sc}$$

M_{pc} and M_{sc} represents the concentration of element in plant and soil mg/kg respectably.

Statistical analysis

Statistical analysis employed SAS windows version 9.1. The significance of differences between variables

Table 1: Factorial design of treatments

R1	M0T2	M1T1	M2T1	M0T0	M1T2	M0T1	M1T0	M2T0	M2T1
R2	M1T0	M0T0	M2T1	M2T0	M1T1	M2T1	M0T1	M1T2	M0T2
R3	M2T1	M1T2	M2T0	M1T0	M0T2	M2T1	M1T1	M0T0	M0T1

Table 2: Characteristics of soil and organic matter

Characteristics	texture	TOC* (%)	pH	Total N (%)	Total P (mg/Kg)	Total K (mg/Kg)	Total Cd (mg/Kg)	EC (ds/m)
Soil	sandy loam	0/68	7.8	0.06	12	105	0.043	2.2
Compost	---	21.3	7.53	1.39	0.65	2.1	0.02	2.5

*TOC: Total organic carbon




Fig. 1: Effects of compost and humic acid on Cd concentration in soil column. Values are means (n = 3) Similar letters are not significantly different at $P < 0.05$, according to Duncan's test. M0: without organic substance, M1: compost, M2: Humic acid.

at $P < 0.05$ was checked with a multiple comparison on Duncan's test.

RESULTS AND DISCUSSION

Concentration of Cd in soil column

Application of organic substances compared to control led to increase available Cd concentration in all treatments (Fig. 1). Highest concentration of Cd

recorded in depth 30-45 cm for M_2T_2 treatment. It was clear that application of organic substances enhanced Cd concentration by depth. At T_1 treatment, application of organic substances with no significant difference between compost and humic acid decreased Cd concentration in 0-15cm but increased in lower depths. In M_1T_2 treatment and depth 30-45cm, Cd concentration increased about 74 and 20% compare to depth 0-15




Fig. 2: Effects of compost and humic acid on Cr concentration in the soil column. Values are means (n = 3). Similar letters are not significantly different at $P < 0.05$, according to Duncan's test. M0: without organic substance, M1: compost, M2: Humic acid.

and 15-30cm respectively. Cadmium movement in soil column was more pronounced by humic acid than compost. In general over the test period, the concentration of cadmium decreased in 0-15 cm and accumulation of Cd was happened in the depth 30-45 cm. Therefore application of organic substances enhanced mobility of Cd. Increased heavy metals amount such as Pb and Cd by compost have been reported (Ben Achiba *et al.*, 2009). Enhancement in Cd availability and mobility eventually was due to multiple functional groups of organic substances (Zhang *et al.*, 2013) that can interacted with metal ions (Cd). This result is alike with other researchers (Angelova *et al.*, 2013; Meers *et al.*, 2005) indicating that Cd concentration was effected by soil properties such as organic substances.

Concentration of Cr in soil column

Fig. 2 is a presentation of the concentrations of Cr in soil column treated with organic substances.

Application of humic acid and compost increased the concentration of chromium in lower depths. Highest concentration of Cr observed in 30-45 cm and M_2T_2 treatment. Cr concentration in 30-45 cm and M_1T_1 was increased about 16.6 % compared to M_0T_1 . There was no significant difference between types of organic substances for 0-15 cm. Cr concentration in 15-30 cm enhanced about 50% in M_2T_2 and 43% in M_1T_2 compare to M_0T_0 . Humic acid was effective than compost as Cr concentration in lower depths. Concentration of Cr in soil column followed increasing trend but it was less than cadmium. Organic substances can form complex compounds with Cr (Barancikova and Makovnikova, 2003) and also maintaining Cr in an exchangeable situation. Cr competition with other elements such as Ca in the creation of complexes with organic materials (Mohammad *et al.*, 2013; Hernandez *et al.*, 2012) and low mobility in soil (Janos *et al.*, 2010) can be the reasons for those results.

Cd concentration in shoot and root

Data of the Table 3 showed that the concentration of Cd in shoot and root of plant (received Cd treatments) was enhanced with application of organic substances. At T₀ treatment, the Cd concentration of maize plant (shoot and root) showed no significant difference. humic acid in combination with T₁ and T₂ which was more effective than compost as increasing Cd concentration in root and shoot. Increasing in cadmium concentration in maize plant can be attributed availability and mobility of this element in soil (Fig. 1). At M₁T₂ and M₂T₂ treatments, Cd concentration increased by 26% and 132% in shoot and 4.5% and 80% in root as compared to control (M₀T₂) respectively. These results suggest that soil organic substances can effectively increase Cd concentration in maize plant in polluted soils. Furthermore, root Cd concentration exceeded that of shoot. M₂T₁ and M₂T₂ treatments increased cd concentration in root by 100 and 80% compared to control (M₀T₂), respectively. In order to preventing toxic accumulation of heavy metals in shoot, plant roots can reserve heavy metal by retention of them in vacuoles (Orrono and Lavado, 2009; Fischerova et al., 2005). Cadmium accumulation in root of maize plant than shoot can be due to connecting of Cd in cation exchange sites of roots (Zeng et al., 2011) and creating complexes with ligands containing -S group (Sulfhydryl) (Topcuoglu, 2012) which led to sediment and accumulation of cd in apoplast of root. Maximum Ti of Cd with no significant different between

organic substances was observed in T₀ treatment (Table 2). Compared with the respective T₀ treatment increment of soil Cd concentration decreased Ti significantly. Such an increment in Cd concentration in soil (Fig. 1) or in root (Table 2) will activate mechanisms of retention of Cd in root and causes hindrance of its translocation. Minimum amount of Ti was observed in M₀T₂. Even though no significant different was observed in Ti recorded for application of organic substances but increasing level of Cd concentration from T₁ to T₂ in combination with M₁ and M₂ increased Ti. The decreasing trend in T₀ to T₁ and increasing trend from T₁ to T₂ in combination with M₁ and M₂ treatment indicated that eventually increasing Cd concentration in root reduces Cd retention capacity which in turn increase Cd up ward translocation. Such reduction in Cd retention capacity might be due to neutralization of negative charges in apoplast (Park et al., 2013; Strobel et al., 2005) or limited capacity of phytochelation production of plant cells (Parsafar and Marofi, 2013). As a consequence, the transmission of Cd in plant (root to shoot) specified the distribution of Cd concentration in plant (Haliru et al., 2009).

Cr concentration in shoots and roots

Enhancement of Cr application increased it concentration in plant (shoot and root) (Table 4).The influence of organic substances on concentration of Cr in shoot followed decreasing trend. In M₁T₂ and M₂T₂ treatments, Cr concentration in shoot was

Table 3: Effect of application of heavy metals and type of organic substances in concentration and transportation index of Cd in maize plant (shoot and root)

Treatment	shoot Cd (mg/kg dw)			root Cd (mg/kg dw)			Transportation index (Ti)		
	M ₀	M ₁	M ₂	M ₀	M ₁	M ₂	M ₀	M ₁	M ₂
T ₀	0.08 ^c	0.07 ^c	0.07 ^c	0.09 ^c	0.08 ^c	0.08 ^c	0.88 ^a	0.87 ^a	0.87 ^a
T ₁ (Cr ₁₀₀ Cd ₂₅)	8.5 ^d	10.4 ^d	16.2 ^b	14.3 ^d	15.4 ^d	28.4 ^b	0.6 ^b	0.67 ^b	0.57 ^{bc}
T ₂ (Cr ₂₀₀ Cd ₅₀)	13.3 ^c	16.7 ^b	30.9 ^a	25.1 ^c	24.2 ^c	45.1 ^a	0.53 ^c	0.69 ^b	0.68 ^b

Treatments mean (n = 3). The same letters in each column showing no significant effect (P<0.05), by Duncan's test. M₀: without organic substance, M₁: compost, M₂: Humic acid.

Table 4: Effect of application of heavy metals and type of organic substances in concentration and transportation index of Cr in maize plant (shoot and root)

Treatment	shoot Cr (mg/kg dw)			root Cr (mg/kg dw)			Transportation index (Ti)		
	M ₀	M ₁	M ₂	M ₀	M ₁	M ₂	M ₀	M ₁	M ₂
T ₀	0.04 ^f	0.03 ^f	0.03 ^f	0.09 ^e	0.07 ^e	0.08 ^e	0.44 ^d	0.42 ^d	0.42 ^d
T ₁ (Cr ₁₀₀ Cd ₂₅)	71.6 ^b	30.4 ^c	24.3 ^e	54.3 ^d	61.7 ^d	98.1 ^b	1.31 ^a	0.49 ^d	0.24 ^e
T ₂ (Cr ₂₀₀ Cd ₅₀)	98.8 ^a	59.8 ^c	38.9 ^d	91.9 ^c	92.6 ^c	144.7	1.07 ^b	0.64 ^c	0.26 ^e

Treatments mean (n = 3). The same letters in each column showing no significant effect (P<0.05), by Duncan's test. M₀: without organic substance, M₁: compost, M₂: Humic acid.

Table 5: Bio accumulation factor (BAF) of heavy metals

Treatment	Cd BAF			Cr BAF		
	M ₀	M ₁	M ₂	M ₀	M ₁	M ₂
T ₀	1/09 ^a	1/05 ^a	0/9 ^b	3/09 ^a	2/06 ^c	2/7 ^b
T ₁ (Cr ₁₀₀ Cd ₂₅)	1/1 ^a	0/5 ^c	0/5 ^c	1/4 ^d	0/57 ^f	0/53 ^f
T ₂ (Cr ₂₀₀ Cd ₅₀)	1/16 ^a	0/44 ^e	0/38 ^d	1/06 ^e	0/7 ^f	0/6 ^f

Treatments mean (n = 3). The same letters in each column showing no significant effect ($P < 0.05$), by Duncan's test. M₀: without organic substance, M₁: compost, M₂: Humic acid.

decreased by 39.4 and 60.6% compared to control (M₀T₂), respectively. Minimum amount of Cr concentration in shoot was observed in M₂T₁ treatment. In T₁ and T₂ treatments, application of organic substances increased concentration of Cr in root compared to T₀. M₀ and M₁ in T₀ and T₁ treatments showed no significant difference. In M₁T₁ and M₂T₁ treatments, Cr concentration in root was increased by 13.6 and 80.6% compared to control (M₀T₁), respectively. Highest accumulation of Cr concentration in root was observed in M₂T₂ treatment. Low mobility of Cr in plant (root to shoot) can be reasons by saturation and accumulation of Cr in apoplast of cells (Kim et al., 2004; Topcuoglu, 2012). Root Cr concentration was greater than shoot Cr concentration. In general chromium is mainly accumulated in roots due to immobility of Cr in vacuoles of root cells (Park et al., 2011). Enhance in Cr concentration due to application of organic substances attributed to the formation of complex organic matter with metal. Therefore the highest amount of Cr uptake remain in roots. Experiments indicated that Cr has low mobility then Cd in plants (Haliru et al., 2009; Kalantari et al., 2014). Researchers find that Cd and Cr can complex by different functional groups of organic substances, this compounds were regulated heavy metal mobility and concentration in soil (Angelova et al., 2013; Lesage et al., 2005). In general chromium by containing negative charges, was immobilized in roots of plant and its translocation was low. The transportation index (Ti) of Cr is shown in Table 3. The fewest amount of Ti was in M₂T₂. In T₁ and T₂ treatments, Transportation index of Cr followed decreasing trend with application of organic substances. The low Transportation index of Cr represents the accumulation of Cr in roots. Application of organic substances decrease Ti. The concentration of Cr between root and shoot indicated that there was a limitation root to shoot movement of Cd and Cr (Wojcik et al., 2005).

Bio accumulation factor (BAF)

The Lowest values of BAF after control was observed in M₁T₂ (Table 5). This factor is directly related to the concentration of elements in plant shoot. So that increasing concentrations of cadmium and chromium in plant shoot (Tables 3, 4) increased this factor. High mobility of cadmium in the effect of organic substances in soil led to decline BAF in M₁ and M₂ treatments. In T₁ and T₂, application of organic matter was decreased BAF due to the low concentration of chromium in plant against soil. According to study bio accumulation factor, can be realized the behavior of plant in uptake of heavy metals. BAF was used for qualification of plant species for phytostabilization of contaminated soils. The BAF values in the range of 0.1-0.50 (as potential phytostabilization) and values from 0.5-1.0 (as partial phyto-stabilization) should not be neglected (Balabanova, et al., 2015).

CONCLUSION

In conclusion, it seems that chromium is likely to increase in plant roots enhances root to shoot movement of cadmium. Cr was accumulated more than Cd by the roots. Root heavy metal concentration was greater than shoot. Mobility of cadmium and chromium in soil was effected by application of organic substances. Organic substances by multiple functional groups can creation metal-humic complexes and increase heavy metals mobility in soil.

ACKNOWLEDGEMENT

The authors thanks the members of the Department of Soil Science, Shahid Chamran University of Ahvaz for their helps for perform this study.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest regarding the publication of this manuscript.

REFERENCES

- Angelova, V.R.; Akova, V.I.; Artinova, N.S.; Ivanov, K.I., (2013). The effect of organic amendments on soil chemical characteristics. *Bulg. J. Agric. Sci.*, 19: 958-971. **(12 pages)**
- Balabanova, B.; Stafilov, T.; Baceva, K., (2015). Bioavailability and bioaccumulation characterization of essential and heavy metals contents in *R. acetosa*, *S. oleracea* and *U. dioica* from copper polluted and referent areas. *J. Environ. Health Sci. Eng.*, 13: 21-29. **(8 pages)**
- Barancikova, G.; Makovnikova, J., (2003). The influence of humic acid quality on the sorption and mobility of heavy metals. *Plant, Soil Environ.*, 49: 565–571 **(11 pages)**.
- Ben Achiba, W.; Gabteni, N.; Lakhdar, A.; Du Laing, G.; Verloo, M.; Jedidi, N.; Gallali, T., (2009). Effects of 5-year application of municipal solid waste compost on the distribution and mobility of heavy metals in a Tunisian calcareous soil *Agriculture. Ecosyst. Environ.* 130: 156–163 **(7 pages)**.
- Boner, K.; Pohl, B., (2010). The Determination of Cd, Cr, Cu, Ni and Pb in a Concentrated CaCl₂/NaCl Solution by AAS. *Agilent Technologies*.
- Chapman, H.D.; Pratt, P.F., (1961). *Methods of analysis for soils, plants and waters*. University of California, Berkeley.
- Fischerova, Z.; Szakova, J.; Pavlikova, D.; Tlustos, P., (2005). The application of diffusive gradient technique (DGT) for assessment of changes in Cd, Pb, and Zn mobility in rhizosphere. *Plant Soil Environ.* 51: 532–538 **(6 pages)**.
- Gabrijel, O.; Davor, R.; Zed, R.; Marija, R.; Monika, Z., (2009). Cadmium accumulation by muskmelon under stress in contaminated organic soil. *Sci. Total Environ.*, 407: 2175–2182 **(7 pages)**.
- Haliru, M.; Ajibola, V.O.; Agbaji, E.B., (2009). Evaluation of the uptake and accumulation of metals by some commonly irrigated vegetables in soils treated with different concentration of these metals. *J. App. Sci.*, 9:1573 – 1577 **(4 pages)**.
- Hernandez-Soriano, M.; Jose, C.; Jimenez-Lopez, C., (2012). Effects of soil water content and organic matter addition on the speciation and bioavailability of heavy metals. *Sci. Total Environ.*, 423: 55–61 **(6 pages)**.
- Hoop, M.; Leeuwen, H.; Cleven, R., (1990). *Chim Anal Acta.* 191-232 **(40 pages)**.
- Jafaraghadi, A.; Sayyad, G.H.; Homei, M. (2010). Investigation of heavy metal values in soil of Khuzestan (Ahvaz) field. *National Conference of Environmental Crises*.
- Janos, P.; Vavrova, J.; Lucie, H., (2010). Effects of inorganic and organic amendments on the mobility (leachability) of heavy metals in contaminated soil: A sequential extraction study. *Geoderma* 159: 335–341 **(6 pages)**.
- John, R.; Ahmad, P.; Gadgil, K.; Sharma, S., (2009). Heavy metal toxicity: Effect on plant growth, biochemical parameters and metal accumulation by *Brassica juncea* L. *Intl. J. Plant Prod.* 3:1735-1743 **(8 pages)**.
- Kabata, A.; Mukherjee, B.A., (2007). *Trace Elements from Soil to Human*. New York: Springer. **450p**.
- Kalantari, K.M.; Pirooz, P.S.; Nasibi, F., (2014). A physiological analysis of sunflower under chromium stress: Impact on plant growth, bioaccumulation and oxidative stress induction on sunflower (*Helianthus annuus*). *J. Plant Biol.*, 11: 73-86 **(13 pages)**.
- Karbassi, A.R.; Pazoki, M., (2015). Environmental qualitative assessment of rivers sediments. *Global J. Environ. Sci. Manage.*, 1(2): 109-116 **(8 pages)**.
- Karbassi, A.R.; Tajziehchi, S.; Afshar, S., (2015). An investigation on heavy metals in soils around oil field area. *Global J. Environ. Sci. Manage.*, 1(4): 275-282 **(8 pages)**.
- Khan, A.G.; Kuek, C.; Chandhry, T.M.; Khoo, C.S.; Hayes, W.J., (2000). Role of plants, mycorrhizae and phytochelators in heavy metal contaminated land remediation, *Chemosphere*, 41: 197–207 **(10 pages)**.
- Kim, Y.J.; Kwon, Y.; Jeong, S., (2004). Combined effects of copper, cadmium, and lead upon *Cucumis sativus* growth and bioaccumulation. *Sci. Total Environ.* 326: 85–93 **(8 pages)**.
- Lesage, E.; Meers, E.; Vervaecke, P.; Lamsal, S.; Hopgood, M.; Tack, F.M.G.; Verloo, M.G., (2005). Enhanced phytoextraction: II. Effect of EDTA and citric acid on heavy metal uptake by *Helianthus annuus* from a calcareous soil. *Int. J. Phytoremed.* 7: 143–152 **(9 pages)**.
- Marchiol, L.; Assolari, S.; Sacco, P.; Zerbi, G., (2004). Phytoextraction of heavy metals by canola (*Brassica napus*) and radish (*Raphanus sativus*) grown on multicontaminated soil. *Environ. Pollut.*, 132: 21–27 **(5 pages)**.
- Meers, E.; Unamuno, V.; Vandegehuchte, M.; Vanbroekhoven, K.; Gebelen, W.; Samson, R.; Vangronsveld, J.; Diels, L.; Ruttens, A.; Laing, G.D.; Tack, F., (2005). Soil-solution speciation of Cd, as affected by soil characteristics in unpolluted and polluted soils. *Environ. Toxicol. Chem.*, 24: 499-509 **(10 pages)**.
- Mireles, A., (2004). Heavy metal accumulation in plants and soil irrigated with waste water from Mexico City. *Nuc. Instr. Method Phys. Res., B.* 220: 187-190 **(3 pages)**.
- Mohammad, J.K.; Muhammad, T.; Khalid, K., (2013). Effect of organic and inorganic amendments on the heavy metal content of soil and wheat crop irrigated with wastewater. *Sarhad J. Agric.* 29: 142-152 **(9 pages)**.
- Orrono, D.; Lavado, R., (2009). Heavy metal accumulation in *Pelargonium hortorum*: Effects on growth and development. *¥OTON: Revista Int. De Botanica Expert Int.*, 78: 75-84 **(9 pages)**.
- Ouni, Y.; Ghnaya, T.; Montemurro, C.; Abdely, A.; Lakhdar, A., (2014). The role of humic substances in mitigating the harmful effects of soil salinity and improve plant productivity. *Int. J. Plant Product.*, 8: 353-374 **(21 pages)**.
- Park, J.; Dane, L.; Periyasamy, P., (2011). Role of organic amendments on enhanced bioremediation of heavy metal (loid) contaminated soils. *J. Hazard. Mater.*, 185: 549-574 **(25 pages)**.
- Park, S.; Kim, K.; Kang, D.; Yoon, H.; Sung, K., (2013). Effects of humic acid on heavy metal uptake by herbaceous plants in soils simultaneously contaminated by petroleum hydrocarbons. *Environ. Earth Sci.*, 68(8): 2375-2384 **(10 pages)**.
- Parsafar, N.; Marofi, S., (2014). Investigation of transfer coefficients of Cd, Zn, Cu and Pb from soil to potato under wastewater reuse. *J. Water Soil Sci.*, 17(66): 199-209 **(11 pages)**.
- Pizzeghello, D.; Francioso, O.; Ertani, A.; Muscolo, A.; Nardi, S., (2013). Isopentenyladenosine and cytokinin-like activity of different humic substances. *J. Geochem. Ex.* 129: 70-75 **(5 pages)**.

- Raymond, A.; Okieimen, E., (2011). Heavy metal in contaminated soils. International Scholarly Research Network ISRN Ecology.
- Salati, S. Quadri G. Tambone F and Adani F. 2010. Fresh organic matter of municipal solid waste enhances phytoextraction of heavy metals from contaminated soil. *Environ. Pollut.* 158: 1899–1906 (7 pages).
- Sanita, L.; Fossati, F.; Musetti, R.; Mikerezi, I.; Favali, M.A., (2010). Effects of hexavalent chromium on maize and cauliflower plants. *J. Plant Nutr.* 25:701-717 (16 pages).
- Strobel, B.W.; Borggaard, O.K.; Hansen, H.C.B.; Andersen, M.K.; Aulund-rasmussen, K., (2005). Dissolved organic carbon and decreasing pH mobilize cadmium and copper in soil. *Eur. J. Soil Sci.*, 56: 189–196 (7 pages).
- Tingqiang, L.; Tao, Q.; Chengfeng, L.; Shohag, M.J.; Xiaoe, Y.; Sparks, D., (2013). Complexation with dissolved organic matter and mobility control of heavy metals in the rhizosphere of hyperaccumulator *Sedum alfredii*. *Environ. Pollut.*, 182: 248-255 (7 pages).
- Topcuoglu, B., (2012). The influence of humic acids on the metal bioavailability and phytoextraction efficiency in long-term sludge applied soil. Conference on International Research on Food Security, Natural Resource Management and Rural Development. Tropentag, Gottingen, Germany.
- Wojcik, M.; Vangronsveld, J.; Tukiendorf, A., (2005). Cadmium tolerance in *Thlaspi caerulescens*. I. Growth parameters, metal accumulation and phytochelatin synthesis in response to cadmium. *Environ. Exp. Bot.* 53: 151–161 (10 pages).
- Wyszkowski, M.; Zio³kowska, A., (2013). Content of polycyclic aromatic hydrocarbons in soils polluted with petrol and diesel oil after remediation with plants and various substances. *Plant, Soil Environ.*, 59: 287–294 (7 pages).
- Zeng, F.; Shafaqat, A.; Zhang, H., (2011). The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants. *Environ. Pollut.*, 159: 84-91 (6 pages).
- Zhang, Y.; Yang, X.; Tian, S.; Guo, W.; Wang, J., (2013). The influence of humic acids on the accumulation of lead and cadmium in tobacco leaves grown in different soils. *J. Soil Sci. Plant Nutr.*, 13: 43-53 (10 pages).

AUTHOR(S) BIOSKETCHES

Chaab, A., Ph.D. Candidate, Department of Soil Science, College of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Iran.
Email: ali.chaab87@gmail.com

Moezzi, A.A., Ph.D., Associate Professor, Department of Soil Science, College of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Iran.
Email: moezzi251@gmail.com

Sayyad, G.A., Ph.D., Assistant Professor, Department of Soil Science, College of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Iran.
Email: gsayyad@gmail.com

Chorom, M., Ph.D., Associate Professor, Department of Soil Science, College of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Iran.
Email: m.chorom@scu.ac.ir

COPYRIGHTS

Copyright for this article is retained by the author(s), with publication rights granted to the journal.
This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

HOW TO CITE THIS ARTICLE

Chaab, A.; Moezzi, A.A.; Sayyad, G.A.; Chorom, M., (2016). Effect of compost and humic acid in mobility and concentration of cadmium and chromium in soil and plant. *Global J. Environ. Sci. Manage.*, 2(4): 389-396.

DOI: 10.22034/gjesm.2016.02.04.008

URL: http://gjesm.net/article_19697.html

