

Original Article

Morphometric analysis of olfactory organ and telencephalon in maturing and mature migrants of Caspian lamprey (*Caspiomyzon wagneri*, Kessler 1870)

Ashraf Namdariyan Rad¹, Bagher Mojazi Amiri*¹, Soheil Eagderi¹, Allison Kupsc²

¹Department of Fisheries, Faculty of Natural Resources, University of Tehran, 31585-4314 Karaj, Iran.

²Department of Environmental Sciences, University of California, Riverside, CA 92521 USA.

Abstract: This study was conducted to provide a detailed information about changes of the olfactory organ and telencephalon morphology in spring and fall spawning run maturing and mature Caspian lamprey, *Caspiomyzon wagneri*, in the Shirud River, Sothern Caspian Sea basin, Iran. A total of 71 maturing and mature fish were collected during their spawning migration. The results showed that the thickness of the olfactory epithelium and the density of ciliated olfactory receptor cells (ORC) were lower in mature migrants. In addition, the nasal cavity, relative weight of olfactory organ and relative telecephalon area in mature migrants were larger indicating its more sensitivity to external queues. Based on the results, the olfactory organ and telencephalon of maturing migrants of Caspian lamprey have not developed completely and needs a period of rest in the river to its full development for spawning.

Article history:

Received 4 September 2016

Accepted 11 January 2017

Available online 25 February 2017

Keywords:

Caspian lamprey

Mature

Maturing Morphometry

Olfactory organ

Introduction

The Caspian lamprey, *Caspiomyzon wagneri*, is an endemic species to the Caspian Sea basin and migrates into its northern, western and southern rivers (Coad, 2016). Its populations in the southern part of the Caspian Sea have dramatically decreased due to river pollution, damming, and loss of spawning grounds (Kiabi et al., 1999). Now with “near threatened” status (IUCN, 1996), there is need for special protection of this species.

Caspian lampreys inhabit freshwater rivers during larval development and migrate to the Caspian Sea for feeding after metamorphosis. When sexual maturation approaches, adults migrate to the rivers for reproduction (Holcik, 1986). Spawning migrations to the southern Caspian Sea rivers (e.g. Shirud River, Northern Iran) occur in the middle of March to late-April in spring, and mid-September to late October in fall (Ahmadi et al., 2011) and these migrants can be classified into two sexual types: mature and maturing.

The olfactory organ of lampreys is located in a shallow nasal cavity on the dorsal surface of the head and its olfactory receptors are long having thin dendrites with rounded apical nodes (Thornhill, 1967). During sexual maturation and particular season, the olfactory sensitivity increases (Moore and Scott, 1992), e.g. Sea lampreys (*Petromyzon marinus*) use a specific chemical or pheromone to migrate back to the river, also males secrete pheromones to attract females to spawning grounds (Teeter, 1980; Bjerselius et al., 2000). Pheromones stimulate the olfactory receptors to send the appropriate message to the brain (Weiming et al., 2007).

Lamprey has a small brain comprising the telencephalon, diencephalon, mesencephalon and metencephalon. Their telencephalon, in particular, is very small. The shape of the brain does not change much through the adult period, but increase in size (Scott, 1887). There is significant coupling between the olfactory system and brain controlling areas. The

* Corresponding author: Bagher Mojazi Amiri
E-mail address: bmamiri@ut.ac.ir

Table 1. Body weight (g) and total length (mm) of maturing and mature males or females of the Caspian lamprey in spring and fall. Each value represents the mean \pm SE.

season	sex		total length (mm)	weight (g)
Spring	Male	mature	3788 \pm 362	79 \pm 9
		maturing	3917 \pm 110	93 \pm 8
	Female	mature	3876 \pm 40	102 \pm 5
		maturing	4274 \pm 390	127 \pm 8
Fall	Male	mature	3675 \pm 234	83 \pm 6
		maturing	3745 \pm 388	90 \pm 7
	Female	mature	3723 \pm 168	93 \pm 6
		maturing	3827 \pm 277	104 \pm 9

telencephalon is the olfactory center and mediates a variety of behaviors such as classical conditioning (Overmier and Hollis, 1983), processing of sensory information (Davis and Kassel, 1983; Hofmann, 2001), mating and reproduction (Demski and Beaver, 2001), social behavior (Huber et al., 1997; Kotrschal et al., 1998; Hofmann, 2001; Pollen et al., 2007), aggression (Davis and Kassel, 1983), schooling behavior (Davis and Kassel, 1983; Shinozuka and Watanabe, 2004), and avoidance learning (Portavella et al., 2003, 2004; Vargas et al., 2009). In addition, the size of the telencephalon is associated with residence in structurally complex habitats (Bauchot et al., 1977; Huber et al., 1997), superior spatial learning (Vargas et al., 2000, 2009), greater parental care (Gonzalez-Voyer et al., 2009), monogamy (Pollen et al., 2007), and sociality (Pollen et al., 2007). Sexual maturation can trigger changes in brain morphology (Jacobs et al., 1990; Jacobs and Spencer, 1994; Clayton et al., 1997; Sherry, 1998).

Little studies are available about the reproductive biology and physiology of the Caspian lamprey (Kiabi et al., 1999; Nazari and Abdoli, 2010; Ahmadi et al., 2011). To the best of our knowledge, no information is available about the olfactory organ and its target regions in the brain i.e. telencephalon of the Caspian lamprey. Hence, this study aimed to analysis morphometric features of the olfactory organ and telencephalon in maturing and mature migrants of Caspian lamprey. The findings of the present study can increases our understanding about the changes of the olfactory organ and telencephalon during reproductive spawning migration in mature

and maturing Caspian lamprey and provide valuable information on the spawning migration physiology in this endangered species.

Materials and Methods

Study site: Specimens were collected by hand at night in spring (middle of April) and fall (middle of October) 2013. A total of 71 maturing and mature fish were collected under the Shirud river bridge (200 m upstream from the river mouth; 36°51'N, 50°47'E). The mean water temperature was 14 \pm 3°C and 11 \pm 2°C during sampling period in spring and fall, respectively.

Sampling and biometry: Fish were anaesthetized using a solution of clove powder (250 ppm). A total of 40 specimens, including mature and maturing males and females were selected from the collected specimens in each sampling season (5 per each group). Mature males were opted from maturing males by applying moderate pressure to their abdomen to stimulate sperm ejaculation. Ejaculation was observed in mature males but not in maturing males. Mature and maturing females were separated according to their ovary color i.e. mature female had ovaries in bluish-green color, whereas those of maturing were yellowish. In addition, mature migrants had empty intestines unlike those of maturing migrants which had green and full intestines (Ahmadi et al., 2011). The body weight (g) and total length (mm) of fish were measured using a digital balance and measuring board, respectively (Table 1). The nasal cavity opening was measure from 2D pictures using ImageJ Software. Then, the olfactory organ (without inlet tube and surrounding




Figure 1. Nasal cavity (external feature, up and internal feature, down) in the Caspian lamprey (A: maturing and B: mature migrants).

cartilages) and brain of fish were removed (Figs. 1, 2) and their weight were measured using a digital balance to the nearest 0.01 g. The relative weight of olfactory organ was calculated using the following equation: nose weight/ body weight $\times 100$. Finally, the nasal and brain were fixed in Bouin's fixative.

Histological study: The olfactory organ of specimens was embedded in paraffin and histological sections were prepared at 4-6 μm thickness and stained with Delafield's hematoxylin and eosin based on Eagderi et al. (2013). Numbers of olfactory receptor cells (ORC) per 100 μm of the olfactory epithelium, and olfactory epithelium width were measured using Olympus BH-2 optical microscope equipped to a Dino-Capture camera and ImageJ Software. ORCs were distinguished by their ciliated olfactory knobs with basal bodies at the outer margins and analyzed according to van Denbossche et al. (1995). The area of brain and telencephalon region (as μm^2 in dorsal view) were examined under stereomicroscopy equipped to a Dino-Capture camera and ImageJ Software. The relative size of telencephalon was calculated using the equation: telencephalon area/

brain area $\times 100$.

Statistical Analysis: Differences between sex ratios were determined using Student's t-test. Mean data were compared using analysis of variance (ANOVA). T-test was used to compare data between maturing and mature fish. Data are expressed as mean \pm SD. P -value < 0.05 was considered significant.

Results

Nasal cavity opening in mature Caspian lampreys was larger than that of maturing fish in both spring and fall migrants (Fig. 1). The relative weight of the olfactory organ was significantly greater in mature fish than those of maturing migrants (0.3223 ± 0.002 vs. 0.2847 ± 0.0011 and 0.3347 ± 0.0023 vs. 0.3013 ± 0.0022 in spring and fall migrant males, respectively; 0.3067 ± 0.0019 vs. 0.2793 ± 0.0017 and 0.324 ± 0.0023 vs. 0.295 ± 0.0015 in spring and fall migrant females, respectively) ($P < 0.05$) (Fig. 3).

The thickness of olfactory epithelium (μm) in mature fish were significantly lower than that of maturing migrants (393 ± 1.2 vs. 249.67 ± 3.65 and




Figure 2. Nasal cavity (external feature, up and internal feature, down) in the Caspian lamprey (A: maturing migrants B: mature migrants).

389±1 vs. 256.6±2.2 in spring and fall migrant males, respectively; 354.9±1.1 vs. 276.63±3.31 and 364.2±1.6 vs. 279.3±3.5 in spring and fall migrant females, respectively) ($P<0.05$) (Fig. 4). The numbers of ORC/100 μm of olfactory epithelium in maturing migrants were closely spaced and greater, compared to those of matures (8.67 ± 3.65 vs.

3.33 ± 1.2 and 8.67 ± 2.2 vs. 3.66 ± 1 in spring and fall migrant males, respectively; 6.67 ± 3.31 vs. 4.67 ± 1.1 and 7.33 ± 3.5 vs. 4.67 ± 1.6 in spring and fall migrant females, respectively) ($P<0.05$) (Fig. 4).

Telencephalon area/ brain area (μm^2) (relative telecephalon area) was significantly greater in mature migrants than that of maturing fish (37.3 ± 0.2 vs. 29.4 ± 0.23 and $33.4.33\pm0.17$ vs. 28.1 ± 0.19 in spring and fall migrant males, respectively; 35.8 ± 0.15 vs. 29.1 ± 0.23 and 30.5 ± 0.12 vs. 26.9 ± 0.28 in spring and fall migrant females, respectively) ($P<0.05$). There was a high positive correlation ($R^2=0.98$) between the weight of olfactory organ and telecephalon weight in maturing and mature migrants during spring or fall.

Discussion

This study provided a detailed information about change of the olfactory organ and telencephalon region of maturing and mature migrants of *C. wagneri* during spring and fall migrations. The results showed that the nasal cavity and relative weight of the olfactory organ in mature migrants were larger. In sea lamprey, progressive enlargement of the nasal sac during metamorphosis and maturation has been reported (Lowe et al., 1973; van Denbossche et al., 1997).

The thickness of olfactory epithelium and ORC




Figure 3. The relative weight of the olfactory organ in maturing and mature male and female migrants in spring or fall (a, b=significant, a, a= not significant, $P<0.05$).




Figure 4. The olfactory lamella and the arrangement of sensory cells (A: maturing migrant, B: mature migrant, R: ORC (Arrowheads), B: basal cell, S: supporting cell, and M: mucous cells).

density were greater in non-mature specimens than in mature fish. Thomhill (1967) reported that the olfactory epithelium in adult *Lampretra fluviatilis* had ciliated ORC and our results also extend this observation to the maturing and mature stages. The range of ORC numbers in the olfactory epithelium of upstream migrant *P. marinus* was 2-12 (Zielinski et al., 1994). The enlarged olfactory organ may enable fish to receive large quantities of water and becomes more efficient at chemoreception as more water enters in the cavity and more odors can be detected (Kleerekoper, 1972; VanDenbossche et al., 1995). Maturing migrant lamprey must find food with a smaller nasal cavity, which may lead to an increase in ORC. In contrast, mature migrants not eat and build the reproductive products, resulting in a lower body weight and larger nasal cavity, which may explain the deformed ORC in current study.

Telencephalon area/ brain area was significantly

greater in mature fish compared to maturing migrants. During the larval period, the telencephalon grows slowly, then doubles in size during metamorphosis, and increases linearly until adulthood (Hardisty and Potter, 1971). Furthermore, olfactory sensitivity increases and changes are occurred in olfactory receptors during sexual maturation due to presence of pheromones in the water (Moore and Scott, 1992) maybe causing enlargement of the telencephalon region as observed in this study. Nanomolar concentrations of a sexual pheromone in males can stimulate the olfactory epithelium of adult females (Li et al., 2002). Behavioral responses in vertebrates are closely related to the olfactory system and olfactory-brain regions (Healey, 1972). Olfactory input is relayed on the medial part of the olfactory bulb, the posterior tuberculum, the mesencephalic locomotor region, and finally reach reticulospinal cells in the hindbrain. Activation of this olfactory-motor pathway generated rhythmic ventral root discharges and swimming movements (Derjean et al., 2010).

Based on the findings of the present study, mature fish (males or females) show larger relative nasal sac and telencephalon in both migration seasons indicating more sensitivity to external queues compare to maturing fish. Also, it can be concluded that the olfactory organ and telencephalon of maturing migrants of Caspian lamprey have not developed completely and needs a period of rest in the river to its full development for spawning.

Acknowledgments

This study was supported by Fisheries Department, University of Tehran.

References

- Ahmadi M., Mojazi Amiri B., Abdoli A., Fakharzade S.M.E., Hoseinifar S.H. (2011). Sex steroids, gonadal histology and biological indices of fall and spring Caspian lamprey (*Caspiomyzon wagneri*) spawning migrants in the Shirud River. Southern Caspian Sea. Environmental Biology of Fishes, 92: 229-235.
- Bauchot R., Bauchot M.L., Platel R., Ridet J.M. (1977). The brains of Hawaiian tropical fishes. Brain size and

- evolution. *Copeia*, 1977: 42-46.
- Bjerselius R., Li W., Teeter J.H., Seelye J.G., Johnsen P.B., Maniak P.J., Grant G.C., Polkinghorne C.N., Sorenson P.W. (2000). Direct behavioural evidence that unique bile acids released by larval sea lamprey (*Petromyzon marinus*) function as a migratory pheromone. *Canadian Journal of Fisheries and Aquatic Sciences*, 57: 557-569.
- Clayton N.S., Reboreda J.C., Kacelnik A. (1997). Seasonal changes of hippocampus volume in parasitic cowbirds. *Behavioural Processes*, 41: 237-243.
- Coad B.W. (2016). Review of the Lampreys of Iran (Family Petromyzontidae). *International Journal of Aquatic Biology*, 4(4): 256-268
- Davis R.E., Kassel J. (1983). Behavioral functions of the teleostean telencephalon. In: R.E. David, G. Northcutt (Eds.). *Fish neurobiology: higher brain areas and functions*. Vol. 2. University of Michigan, Ann Arbor. pp: 237-263.
- Demski L.S., Beaver J.A. (2001). Brain and cognitive function in teleost fishes. In: G. Roth, M.F. Wulliman (Eds.). *Brain Evolution and Cognition*. John Wiley and Sons Inc, New York. pp: 297-332.
- Derjean D., Moussaddy A., Atallah E., St-Pierre M., Auclair F., Chang S., Ren X., Zielinski B., Dubuc R. (2010). A novel neural substrate for the transformation of olfactory inputs into motor output. *PLoS Biology*, 8:e1000567.
- Eagderi S., Mojazi Amiri B., Adriaens D. (2013). Description of the ovarian follicle maturation of the migratory adult female bulatmai barbel (*Luciobarbus capito*, Gldenstdt 1772) in captivity. *Iranian Journal of Fisheries Sciences*, 12(3): 550-560.
- GonzalezVoyer A., Winberg S., Kolm N. (2009). Social fishes and single mothers: Brain evolution in African cichlids. *Home | Proceedings of the Royal Society of London B*, 276: 161-167.
- Hardisty M.W., Potter I.C. (1971). The behaviour, ecology and growth of larval lampreys. In: M.W. Hardisty, I.C. Potter (Eds.). *The Biology of Lampreys*, Vol. 1, Academic Press, London. pp: 85-125.
- Healey E.G. (1972). The central nervous system. In: M.W. Hardisty, I.C. Potter (Eds.). *The Biology of Lampreys*, Vol.2, Academic Press, London. pp: 307-372.
- Hofmann M.H. (2001). The role of the fish telencephalon in sensory information processing. In: E.G. Kapor, T.J Karas (Eds.). *Sensory Biology of Jawed Fishes*, New Insights. Science Publishers Inc, Enfield. pp: 255-274.
- Holcik J. (1986). *Caspiomyzon wagneri* (Kessler, 1870). In: J. Holcik (Ed.). *The freshwater fish of Europe*. Aula-Verlag, Wiesbaden. pp: 119-142.
- Huber R., Van Staaden M.J., Kaufman L.S., Liem K.F. (1997). Microhabitat use, trophic patterns, and the evolutions of brain structure in African cichlids. *Brain Behav. Evol.* 50: 167-182.
- IUCN. (1996). IUCN red list of threatened animals. International Union for Conservation of Nature and Natural Resources, Glandv.
- Jacobs L.F., Gaulin S.J.C., Sherry D.F., Hoffman G.E. (1990). Evolution of spatial cognition: Sex-specific patterns of spatial behavior predict hippocampal size. *Proceedings of the National Academy of Sciences*, 87: 6349-6352.
- Jacobs L.F., Spencer W.D. (1994). Natural space-use patterns and hippocampal size in kangaroo rats. *Brain Behav. Evol.* 44: 125-132.
- Kiabi B.H., Abdoli A., Naderi M. (1999). Status of the fish fauna in the south Caspian basin of Iran. *Zoology in Middle East*, 18: 57-65.
- Kleerkoper H., van Eekel G.A. (1960). The olfactory apparatus of *Petromyzon marinus*. *Canadian Journal of Zoology*, 38: 209-223.
- Kotrschal K., Van Staaden M.J., Huber R. (1998). Fish brain: Evolution and environmental relationships. *Reviews in Fish Biology and Fisheries*, 8: 373-408.
- Li W., Scott A.P., Siefkes M.J., Yan H., Liu Q., Yun S.S., Gage D.A. (2002). Bile acid secreted by male sea lamprey that acts as a sex pheromone. *Science*, 296: 138-141.
- Lowe D.R., Beamish F.W.H., Potter I.C. (1973). Changes in the proximate body composition of the landlocked sea lamprey, *Petromyzon marinus* (L.) during larval life and metamorphosis. *Journal of Fish Biology*, 5: 673-682.
- Moore A., Scott A.P. (1992). 17, 20-dihydroxy-4-pregnen-3-one 20 sulphate is a potent odorant in precocious male Atlantic salmon (*Salmo salar*) parr which have been pre-exposed to the urine of ovulated females. *Proceedings of Biological Sciences*, 249: 205-209.
- Nazari H., Abdoli A. (2010). Some reproductive characteristics of endangered Caspian lamprey (*Caspiomyzon wagneri* Kessler, 1870) in the Shirud River southern Caspian Sea, Iran. *Environmental Biology of Fishes*, 88: 87-96.
- Overmier J.B., Hollis K.L. (1983). The teleostean

- telencephalon in learning. In Northcut RG, Davis RE. (Eds.), Ann Arbor: The University of Michigan Press. Fish Neurobiology, 265-284.
- Pollen A.A., Dobberfuhr A.P., Scace J., Igulu M.M., Renn S.C.P. (2007). Environmental complexity and social organization sculpt the brain in Lake Tanganyikan cichlid fish. *Brain, Behavior and Evolution*, 70: 21-39.
- Portavella M., Salas C., Vargas J.P., Papini M.R. (2003). Involvement of the telencephalon in spaced-trial avoidance learning in the goldfish *Carassius auratus*. *Physiology and Behavior*, 80: 49-56.
- Portavella M., Torres B., Salas C. (2004). Avoidance response in goldfish: Emotional and temporal involvement of medial and lateral telencephalic pallium. *Journal of Neuroscience*, 24: 2335-2342.
- Scott W.B. (1887). Notes on the development of *Petromyzon*. *Journal of Morphology*, 1: 253-310.
- Sherry D.F. (1998). The ecology and neurobiology of spatial memory. In: R. Dukas (Ed.). *Cognitive Ecology: The Evolutionary Ecology of Information Processing and Decision Making*. Chicago, IL: University of Chicago Press. pp: 261-296.
- Shinozuka K., Watanabe S. (2004). Effects of telencephalic ablation on shoaling behavior in goldfish. *Physiology and Behavior*, 81: 141-148.
- Teeter J. (1980). Pheromone communication in sea lampreys (*Petromyzon marinus*): Implications for population management. *Canadian Journal of Fisheries and Aquatic Sciences*, 37: 2123-2132.
- Thornhill R.A. (1967). The ultrastructure of the olfactory epithelium of the lamprey, *Lampetra fluviatilis*. *Journal of Cell Science*, 274: 123-132.
- Thornhill R.A. (1970). Cell division in the olfactory epithelium of the lamprey, *Lampetra fluviatilis*. *Zeitschrift Fur Zellforschung Und Mikroskopische Anatomie*, 109: 147-157.
- Van Denbossche J.M., Seelye J.G., Zielinski B.S. (1995). The morphology of the olfactory epithelium in the larval, juvenile and upstream migrant stages of the sea lamprey, *Petromyzon marinus*. *Brain, Behavior and Evolution*, 45: 19-24.
- Van Denbossche J.M., Youson J.H., Pohlman D., Wong E., Zielinski B.S. (1997). Metamorphosis of the olfactory organ of the sea lamprey (*Petromyzon marinus*): morphological changes and morphometric analysis. *Journal of Morphology*, 231: 41-52.
- Vargas J.P., Rodriguez F., Lopez J.C., Arias J.L., Salas C. (2000). Spatial learning-induced increase in the argyrophilic nucleolar organizer region of the dorsolateral telencephalic neurons in goldfish. *Brain Research*, 865: 77-84.
- Vargas J.P., Lopez J.C., Portavella M. (2009). What are the functions of fish brain pallium? *Brain Research Bulletin*, 79: 436-440.
- Weiming L., Michael T., Michael J. (2007). Research to guide use of pheromones to control Sea Lamprey. *Journal of Great Lakes Research*, 33(Special Issue 2): 70-86.
- Wcislo W.T. (1989). Behavioral environments and evolutionary change. *Annual Review of Ecology, Evolution, and Systematics*, 20: 137-169.
- Zielinski B.S., Fredricks K., McDonald R., Zaidi A.U. (1994). Morphological and electrophysiological examination of olfactory sensory neurons during the early developmental prolarval stage of the sea lamprey (*Petromyzon marinus*). *Journal of Neurocytology*, 34: 209-216.

چکیده فارسی

تحلیل ریخت‌سنجی اندام بویایی و تَلن‌سفالون مغز در مارماهی دهان‌گرد خزری
(*Caspiomyzon wagneri*, Kessler 1870) بالغ و در حال بلوغ

اشرف نامداریان‌راد^۱، باقرمجازی امیری^{۲*}، سهیل ایگدری^۱، آلیسون کویسکو^۲

^۱گروه شیلات، دانشکده منابع طبیعی، دانشگاه تهران، ۴۳۱۴-۳۱۵۸۵، کرج، ایران.
^۲گروه علوم محیط زیست، دانشگاه کالیفرنیا، ریورساید CA۹۲۵۲۱، ایالت متحده آمریکا.

چکیده:

این مطالعه با هدف فراهم نمودن اطلاعات در مورد تغییرات ریختی اندام بویایی و تَلن‌سفالون مغز مارماهی دهان‌گرد خزری بالغ و در حال بلوغ در زمان مهاجرت‌های تولیدمثلی بهار و پاییزه به رودخانه شیروود، حوضه آبریز جنوب دریای خزر، ایران به اجرا درآمد. برای این منظور تعداد ۷۱ ماهی بالغ و در حال بلوغ در زمان مهاجرت تولیدمثلی جمع‌آوری شدند. نتایج نشان داد که ضخامت اپیتلیوم بویایی و تراکم سلول‌های گیرنده مژک‌دار بویایی در ماهیان بالغ کمتر بودند. به‌علاوه حفره بویایی، وزن نسبی اندام بویایی و مساحت نسبی تَلن‌سفالون در ماهیان مهاجر بالغ بیشتر بودند که بیانگر حساسیت بیشتر آن به محرک‌های خارجی می‌باشد. براساس نتایج اندام بویایی و تَلن‌سفالون مغز مارماهی دهان‌گرد مهاجر در حال بلوغ هنوز به‌طور کامل توسعه نیافته و نیازمند یک دوره استراحت در رودخانه برای تکمیل توسعه به‌منظور تخم‌ریزی است.
کلمات کلیدی: مارماهی دهان‌گرد خزری، بالغ، در حال بلوغ، ریخت‌سنجی، اندام بویایی.