

Original Article

Cadmium and arsenic bioaccumulation and bio-concentration in the endemic toothed carp *Aphanius arakensis* in salt water

Masoumeh Ariyae¹, Amir Hossein Hamidian^{*1}, Soheil Eagderi², Sohrab Ashrafi¹, Hadi Poorbagher²

¹Department of Environment, Faculty of Natural Resources, University of Tehran, Karaj, Iran.

²Department of Fisheries, Faculty of Natural Resources, University of Tehran, Karaj, Iran.

Abstract: Heavy metals are released to aquatic ecosystems from natural and anthropogenic recourses and accumulate to the body of organisms. This study aimed to assess the accumulation of As and Cd in the gill, liver, and muscle of the toothed carp *Aphanius arakensis* in salt water exposed to three concentrations of Cd and As (5, 10 and 20 mg L⁻¹) for 18 days. The specimens were collected from the Shoor River with an average weight of 1.5±0.3 g (mean±SD) and length of 3.4±0.4 cm. The findings showed that the bio-concentration factor (BCF) of Cd and As were in the following order: liver > gill > muscle, however, for 5 ppm of As the order was gill > liver > muscle. BCF in As concentrations were more than Cd concentrations. Also, the highest BCF was found at 5 ppm. The present study showed that the liver is the organ that accumulates the highest concentrations of As and Cd.

Article history:

Received 28 March 2017

Accepted 7 December 2017

Available online 25 February 2018

Keywords:

Toxicity

Heavy metals

Organs

BCF

Introduction

Fishes are suitable indicators for monitoring the land-based pollutions, particularly heavy metals which accumulate in fish tissues, being absorbed through the skin or indirectly through food (Kamrine, 2000). Heavy metals are important threat to the aquatic environments because of chemical persistent, low degradation, bioaccumulation and bio-magnification (Anbiaee et al., 2009). Among heavy metals, Cadmium is very toxic, 2-20 times more than those of others (Perez-Lopez et al., 2008). Arsenic is another hazardous heavy metal originating from fertilizers and effluents that enter to the surface and groundwater (Philip, 1978).

The heavy metals accumulate mostly in the muscle, liver and gill tissues of fishes due to their high metabolism rates (Khodabande, 2000). The liver plays an important role in accumulation of heavy metals and detoxification of wastes and its histopathological changes may be used as indicators of heavy metals (Wong et al., 2001). The gill is in direct contact with water and thus accumulates heavy metals (Oliveira-Filho et al., 2010). Bioconcentration is a process that

chemicals affect the organisms and bioconcentration factor (BF) is determined by dividing concentration of heavy metals in the organism tissues to concentration of heavy metals in water (Otitoloju et al., 2009).

The genus *Aphanius* is the only representative of the Cyprinodontidae (Teleostei, Cyprinodontiformes) in Eurasia. Iran and central Anatolia show the highest diversity of *Aphanius*, and 14 extant and one fossil species are known to occur in Iran based on data derived from fish morphometry and meristics, otoliths, scales and mtDNA sequences (Coad, 2016; Teimori et al., 2014). Twelve out of 14 Iranian *Aphanius* species are endemic to this country. This genus occurs in coastal (brackish) and landlocked (freshwater to saline) water bodies in the Mediterranean and Persian Gulf basins from Iberian Peninsula as far eastwards as Iran and Pakistan (Teimori et al., 2014). *Aphanius arakensis* is a newly described species of this genus from the Namak Lake basin in Iran (Teimori et al., 2012, 2014). This species tolerates a wide range of physicochemical parameters (Coad, 2016), then it can be an appropriate model to study the effect of heavy metal pollution since it is

*Corresponding author: Amir Hossein Hamidian
E-mail address: a.hamidian@ut.ac.ir




Figure 1. The sampling area (○) in the Shoor River.

found in both salt and fresh water in the Namak lake basin. Hence, this study was conducted to examine the accumulation of Cd and As in the muscle, gill and liver of *A. arakensis* and their bioconcentration factor in different concentrations of As and Cd in the laboratory condition.

Materials and Methods

In this study, a total of 375 female specimens of *A. arakensis* with an average weight of 1.5 ± 0.3 g (mean \pm SD) and length of 3.4 ± 0.4 cm were collected from the Eshtehard Shoor River ($35^{\circ}36'31''$ N, $50^{\circ}48'23''$ E), (Alborz Province, Iran) in June 2011 (Fig. 1). Dissolved oxygen (DO), salinity, water temperature and pH were measured during the sampling, which were 11.68 mg L⁻¹, 11-12 g L⁻¹, $12.85 \pm 6.22^{\circ}\text{C}$ and 7-8.5, respectively.

The fish samples were transported to the fisheries

laboratory of University of Tehran in polythene bags. Prior to the experiment, the fish were acclimatized to the laboratory conditions for five days in pre-cleaned 30 L glass aquaria (30×30×40 cm) filled with de-chlorinated tap water. During the experiment, the fish were fed with a commercial fish food (Biomar) at a rate of 3-5% body weight twice a day. The physicochemical conditions of the aquaria were as follows: temperature, $27.5 \pm 1.2^{\circ}\text{C}$, pH, 7.7 ± 0.5 , CaCO₃ hardness, 295 ± 18 mg L⁻¹ and DO, 7.9 ± 0.1 mg L⁻¹.

Cd and As were used as arsenic oxide (As₂O₃) and cadmium chloride (CdCl₂) salts (Merck, Germany). Twenty-one aquaria were filled with salt water of the sampling area and aerated throughout the experiment. The animals were randomly allocated to 21 aquaria as seven treatments each with three replicates containing 10 fish. Three aquaria received no chemical and was

Table 1. Detection limits, blanks and recoveries of the measurements.

Element	Detection limit ($\mu\text{g/g}$)	Recovery Mean (%) \pm SD	Blank Mean ($\mu\text{g/g}$) \pm SD
Cd	0.29	96.10 \pm 3.77	0.054 \pm 0.006
As	0.19	97.50 \pm 1.46	0.0 \pm 0.0

the control. The other 18 aquaria received only a concentration of Cd (5, 10 and 20 mg L⁻¹) or As (5, 10 and 20 mg L⁻¹) for 18 days.

At the end of the experiment, the animals were anaesthetized and euthanized. A piece of their gill, liver and muscle were removed. Five specimens from each aquarium were pooled to make an appropriate weight of sample (the gill and muscle \approx 1 g, the liver \approx 0.5 g). The organ samples were digested in a mixture containing HNO₃ and HClO₄ (Mansouri et al., 2012c). The tissues were then weighed accurately and put into 150 mL Erlenmeyer flasks. Then, 10 mL nitric acid (65%) were added to each sample. The samples were left overnight to be digested slowly (Baramaki et al., 2012; Mansouri et al., 2012). Finally, 5 mL perchloric acid (70%) were added to each sample. Digestion was performed on a hot plate (sand bath) at 90°C. The digested samples were diluted with 25 ml deionized water. The concentration of Cd and As were measured using ICP-OES (GBC Integra XL, Australia) and presented as mg g⁻¹ wet weight (ww). All the measurements were repeated three times and the average of the values was reported along with their standard deviations.

The following equation was employed to determine the bioconcentration factor in different tissues of *A. arakensis* in different concentrations of metals.

$$\text{BCF} = \text{C tissues} / \text{C water}$$

Where C tissues are average concentration of metals in different tissues and C water: average concentration of metals in water. The analyses of data were performed using SPSS (Version 16).

Results

Only 5% of the specimens died during the experiment. The detection limits, blanks and recoveries of the measurements of metals in the samples are presented in Table 1.

Average accumulation of As and Cd in different




Figure 2. Mean (\pm SD) As and Cd average accumulation in different organs of female *Aphanis arakensis*.

tissues of *A. arakensis* in salt water are shown in Figures 2. The metals were accumulated in different tissues in the following order of magnitude: liver > gill > muscle. The Figures 2 and 3 show that the highest accumulated Cd and As was found in the liver. On the other hand, the lowest accumulated Cd and As found in the muscle. The average accumulation of As and Cd in different tissues of *A. arakensis* in salt water are shown in the Figures 3 and 4. The lowest accumulation of As and Cd was observed in the control group and in the muscle, while the highest accumulation of As and Cd were found in the liver, and 10 and 20 ppm, respectively.

Bio-concentration Factor (BCF) was calculated in different tissues of the specimens. The highest BCF of As was found in the gill and in the concentration of 5ppm. The highest BCF of Cd was found observed in the liver. The lowest BCF of Cd and As were detected in the muscle and in the concentration of 20 ppm (Fig.




Figure 3. Mean (\pm SD) As and Cd accumulation in different organs of female *Aphanius arakensis* in concentrations of 5, 10, 20 ppm and control group.

4). BCF of As was more than that of Cd. The maximum BCF of As and Cd were found at the concentration of 5 ppm (5 ppm > 10 ppm > 20 ppm).

Discussion

Accumulation of metals in the aquatic environment suggests that fishes can serve as a suitable bioindicator for contaminating metals in aquatic environment, because they respond to variations with great sensitivity in aquatic ecosystems (Mansouri et al., 2011). Our findings confirm differences in accumulation of heavy metals in the different tissues. Based on the results of this study, the liver identified as target organ in salt water aquaria (liver > gill > muscle) which is in agreement with Kojadinovic et al. (2007), Monsef Zadeh (2008), Naseri et al. (2005), Norouzi et al. (2012) and Mansouri et al. (2012). Similarly, many studies showed that the liver can accumulate high concentrations of heavy metals due to its key role in the metabolism of the body. Therefore, the liver is suitable indicator for the




Figure 4. BCF of As and Cd in different tissues (ng g⁻¹ wet weight).

assessment of impact of environmental pollutants (Yılmaz, 2009). After the liver, the gill showed a high accumulation of Cd and As concentrations in salt water tanks. The gills are places that accumulate water ions and one of the first tissues that have direct contact with environmental pollutants. The gill is also exposed to the pollutants to a greater extent compared with other organs, causing a higher rate adsorption and accumulation of the pollutants than those of other organs. As in the liver, the gill can be considered as an indicator of pollutants in both marine and freshwater habitats (Oliveira-Filho et al., 2010).

The present study showed that accumulation of Cd and As was lower in the muscle than those of the liver and gill. This finding is similar to those of Mansouri et al. (2012) and Majnoni et al. (2013), where the rate of bioaccumulation in the muscle was lower than those of the liver and gill. Generally, tissues which have low metabolic activities accumulate high concentrations of heavy metals (Amini Ranjabr and Sotoodena, 2005). However, the muscle is not an active tissue to accumulate heavy metals which has been confirmed in this study and by others (Tekin-Ozan, and Kir, 2007;

Karadede, 2000; Wong et al., 2001). Terra et al. (2008) indicated that low concentrations of heavy metals in the muscle may be a result of a low level of binding proteins.

According to the finding of this study and other studies (Subathra and Karuppasamy, 2008; Senthil Murugan et al., 2008), the order of the magnitude of BCF for Cd and As in tissues were liver > gill > muscle but this order in 5 ppm of As was gill > liver > muscle which is similar to findings of Jayakumar and Issac Paul (2006) and Asagba et al. (2008). The liver is a place for storage, detoxification and biological transmission of heavy metals in fishes. Therefore, the highest accumulation of heavy metals is normally found in the liver (Kalay and Canli, 2000). It is shown that a large amount of metallothionein induction happens in the liver of fishes. Transmission of a large volume of water through the gills to obtain oxygen in stressful condition may increase rapidly toxic heavy metals in the gills (Karuppasamy, 2004). Hamidian et al. (2013) ignored results of this study that BCF of As was more than that of Cd. However, metabolic processes such as biological transmission, active uptake and fat metabolism affect bioaccumulation plus other factors including contamination gradients of water, temperature, salinity and interacting agents (Van Geest, 2010).

Acknowledgments

We would like to thank M. Khazaei for his help. This study was financially supported by the University of Tehran.

References

Amini Ranjbar G., Sotudenia F. (2005). Accumulation of heavy metals in muscle of Caspian Sea Golden mullet (*Mugil auratus*) in relation with biometric characteristic (standard length, weight, age and genus). Iranian scientific Fisheries Journal, 13-18.

Anbiaee A., Keykhosro A., Vatandoost J. (2009). Study of toxic effect of different concentrations of Zn and Cu in liver and muscle of common carp (*Cyprinus Carpio*). The 2th National Congress on Environmental Health, Shahid Beheshti University of Medical Science.

Asagba S.O., Eriyamremu G.E., Igberaese M.E. (2008).

Bioaccumulation of cadmium and its biochemical effect on selected tissues of the catfish (*Clarias gariepinus*). Fish Physiology and Biochemistry, 34: 61-69.

Baramaki R., Ebrahimpour M., Mansouri B., Rezaei M.R., Babaei H. (2012). Contamination of metals in tissues of *Ctenopharyngodon idella* and *Perca fluviatilis*, from Anzali Wetland, Iran. Bulletin of Environmental Contamination and Toxicology, 89: 831-835.

Coad B.W. (2016). Freshwater Fishes of Iran. www.briancoad.com. (accessed 30 January 2016).

Hamidian H., Zareh M., Poorbagher H., Vaziri L., Ashrafi S. (2013). Heavy metal bioaccumulation in sediment, common reed, algae and blood worm from the Shoor River, Iran. Toxicology and Industrial Health, 32(3): 398-409.

Jayakumar P., Issac Paul V. (2006). Patterns of cadmium accumulation in selected tissues of the catfish *Clarias ba trachus* (Linn.) exposed to sublethal concentration of cadmium chloride. Veterinary Archives, 76: 167-177.

Kalay M., Canli M. (2000). Elimination of essential (Cu and Zn) and non-Essential (Cd and Pb) metals from tissues of a freshwater fish *Tilapia zilli*. Turkish Journal of Zoology, 24: 429-436.

Karadede H., Unlu E. (2000). Concentrations of some heavy metals in water, sediment and fish species from the Ataturk dam Lake Euphrates, Turkey. Chemosphere, 41: 1371-1376.

Karuppasamy R. (2004). Evaluation of Hg concentration in the tissue of fish *Channa punctatus* (Bloch.) in relation to short and long-term exposure to phenylmercuricacetate. Journal of Platinum Jubilee Australia, 40: 197-204.

Kamrine M.A. (2000). Pesticide profiles, toxicity, environmental impact, and fate. CRC publisher. 704 p.

Khodabandeh S. (2000). Heavy metals accumulation in the sediments and aquatics of Caspian Sea. Water and Wastewater, 42: 19-29.

Kojadinovic J., Potier M., Le Corre M., Cosson M., Bustamante M. (2007). Bioaccumulation of trace elements in pelagic fish from the Western Indian Ocean. Environmental Pollution, 146: 548-566.

Majnoni F., Mansouri B., Rezaei M.R., Hamidian A.H. (2013). Contaminations of metals in tissues of Common carp, *Cyprinus carpio* and Silver carp, *Hypophthalmichthys molitrix* from Zarivar wetland, western Iran. Archives of Polish Fisheries, 21: 11-18.

Mansouri B., Ebrahimpour M., Babaei H. (2012). Bioaccumulation and elimination of nickel in the organs

- of black fish (*Capoeta fusca*). Toxicology and Industrial Health, 28: 361-368.
- Mansouri B., Pourkhabbaz A., Babaei H., Farhangfar H. (2011). Experimental studies on concentration and depuration of cobalt in the selected organs of fresh water fish *Capoeta fusca*. World Journal of Fish and Marine Sciences, 3: 387-392
- Monesef Zadeh F., 2008. Determination of Pb, Cd, Hg, Zn and Cu in liver and muscle of kutum fish (*Rutilus frisii kutum*) in the coast of Guilan province. Master of Science thesis in Environmental Engineering, University of Guilan.
- Naseri M., Rezaee M., Abedi A., AfsharNaderi A. (2005). Investigation of some heavy metal concentrations (Fe, Cu, Zn, Mg, Mn, Hg, Pb and Cd) in the edible and nonedible of lizadussumieri tissues in the coast of Bushehr. Journal of Marine Science and Technology, 4: 59-67.
- Nowrouzi M., Mansouri B., Hamidian A.H., Zarei I., Mansouri A. (2012). Metal contents in tissues of two fish species from Qeshm Island, Iran. Bulletin of Environmental Contamination and Toxicology, 89: 1004-1008.
- Oliveira-Filho E.C., Muniz D.H.F., Ferreira M.F.N. (2010). Cesar Koppe Grisolia evaluation of acute toxicity, cytotoxicity and genotoxicity of a nickel mining waste to *Oreochromis niloticus*. Bulletin of Environmental Contamination and Toxicology, 85: 467-471.
- Otitoloju A., Elegba O., Osibona A. (2009). Biological responses in edible crab, *Callinectes amnicola* that could serve as markers of heavy metals pollution. Journal of Environmental Sciences, 29: 37-46.
- Perez-Lopez M., Hermoso de Mendoza M., Lopez Beceiro A., SolerRodriguez F. (2008). Heavy metal (Cd, Pb and Zn) and metalloid (As) content in raptor species from Galicia (NW Spain). Ecotoxicology and Environmental Safety, 70: 154-162.
- Philip C., Singer E.D. (1973). Trace Metals and Metal-organic Interaction in Natural Waters, Ann Arbor Science Pcience Publishes, Inc. 580 p.
- Senthil Murugan S., Karuppasamy R., Poongodi K., Puvaneswari S. (2008). Bioaccumulation pattern of zinc in freshwater fish *Channa punctatus* (Bloch.) after chronic exposure. Turkish Journal of Fish Aquatic Sciences, 8: 55-59.
- Subathra S., Karuppasamy R. (2008). Bioaccumulation and depuration pattern of copper in different tissues of *Mystus vittatus*, related to various size groups. Archive of Environmental Contamination and Toxicology, 54: 236-244.
- Teimori A., Esmaeili H.R., Gholami Z., Zarei N., Reichenbacher B. (2012). *Aphanius arakensis*, a new species of tooth-carp (Actinopterygii, Cyprinodontidae) from the endorheic Namak Lake basin in Iran. ZooKeys, 215: 55-76.
- Teimori A., Esmaeili H.R., Erpenbeck D., Reichenbacher B. (2014). A new and unique species of the genus *Aphanius* (Teleostei: Cyprinodontidae) from Southern Iran: a case of regressive evolution. Zoologischer Anzeiger - A Journal of Comparative Zoology, 253(4): 327-337.
- Tekin-Ozan S., Kir I. (2008). Seasonal variations of heavy metals in some organs of carp (*Cyprinus carpio* L., 1758) from Beyşehir Lake (Turkey). Environmental Monitoring and Assessment, 138: 201-206.
- Terra B.F., Araujo F.G., Calza C.F., Lopes R.T., Teixeira T.P. (2008). Heavy metal in tissues of three fish species from different trophic levels in a tropical Brazilian river. Water, Air and Soil Pollution, 187(1): 275-284.
- Van Geast J. (2010). Bioaccumulation of sediment-associated contaminants in freshwater organism: development and standardization of a laboratory method, PhD thesis, University of Guelph, Canada. 232 p.
- Wong C.K., Wong P.P.K., Chu L.M. (2001). Heavy metal concentrations in marine fishes collected from fish culture sites in Hong Kong. Environmental Contamination and Toxicology, 40: 60-69.
- Yilmaz F. (2009). The comparison of heavy metal concentrations (Cd, Cu, Mn, Pb, and Zn) in tissues of three economically important fish (*Anguilla anguilla*, *Mugilcephalus* and *Oreochromis niloticus*) inhabiting Köycegiz Lake-Mugla (Turkey). Turkish Journal of Science and Technology, 4: 7-15.

چکیده فارسی

تغلیظ و تجمع زیستی کادمیوم و آرسنیک در گورماهی بومی *Aphanius arakensis* در آب شور

معصومه آریایی^۱، امیر حسین حمیدیان^{۱*}، سهیل ایگدری^۲، سهراب اشرفی^۱، هادی پورباقر^۲

^۱گروه محیط زیست، دانشکده منابع طبیعی، دانشگاه تهران، کرج، ایران.

^۲گروه شیلات، دانشکده منابع طبیعی، دانشگاه تهران، کرج، ایران.

چکیده:

فلزات از منابع طبیعی و انسانی به اکوسیستم‌های آبی منتقل می‌شوند و در بدن موجودات زنده انباشته می‌شوند. هدف از این مطالعه ارزیابی انباشت As و Cd در آبشش، کبد و عضله *Aphanius arakensis* در آب شور و در معرض سه غلظت Cd و As (۵، ۱۰ و ۲۰ میلی‌گرم در لیتر) برای ۱۸ روز بود. نمونه‌ها از رودخانه شور با میانگین وزنی $1/5 \pm 0/3$ گرم (میانگین $\pm SD$) و طول $3/4 \pm 0/4$ سانتی‌متر جمع‌آوری شد. یافته‌ها نشان داد که ضریب تغلیظ زیستی Cd و As به این ترتیب است: کبد < آبشش < عضله، اگرچه در غلظت ۵ ppm از As به ترتیب آبشش < کبد < عضله است. BCF آرسنیک بیشتر از کادمیوم بود. همچنین بیشترین میزان BCF در غلظت ۵ ppm مشاهده شد. مطالعه حاضر نشان داد که کبد اندامی است که بیشترین غلظت As و Cd را تجمع می‌دهد. کلمات کلیدی: سمیت، فلزات سنگین، اندام‌ها، تغلیظ زیستی.

Archive of SID