

Original Article

Contrasting fish size distributions between Neotropical run-of-river and storage reservoirs

Louise Cristina Gomes^{*1}, Pitágoras Augusto Piana², Dienes Aparecida de Oliveira Sereia³

¹Pós Graduação em Biologia Comparada, Universidade Estadual de Maringá, Brazil.

²Centro de Engenharias e Ciências Exatas, Universidade Estadual do Oeste do Paraná, Toledo, Paraná, Brazil.

³Centro de Ciências Biológicas, Universidade Tecnológica Federal do Paraná, Dois Vizinhos, Paraná, Brazil.

Abstract: The spectrum size has been used to assess environmental disturbances and to understanding the energy flow in ecosystems. The objective of this work was to investigate if the operation regimes of reservoirs, including run-of-river and storage systems, interfere with the biomass spectra of fish fauna. We tested the hypothesis that the run of river reservoirs present higher proportions of large individuals than storage system. Samplings of fish fauna were carried out between January 2005 and December 2007 at six Neotropical reservoirs belonging to the sub-basin Iguazu River and Coastal basin in São Jorge River, Brazil. The spectrum calculation was performed using the Pareto type I continuous distribution model. Reservoirs operated under run-of-river regime had significantly higher values than those operated under storage regimes. This study has elucidated some impacts of reservoir operating regime on biomass spectra and indicated differences in size spectra of fish assemblages among the sampled reservoirs. Therefore, it is important to incorporate management plans that take dam operating mode into account so that conservation of aquatic fauna, especially fish, is more effective.

Article history:

Received 7 March 2018

Accepted 24 April 2018

Available online 25 April 2018

Keywords:

Aquatic environments

Dam

Fish assemblage

Size spectrum

Introduction

Body size is a fundamental trait of living organisms, being linked with many important physiological and ecological processes (Giacomini et al., 2016; Wheeland and Rose, 2016; Brown, et al., 2004; Elton, 1927). The relationship between body size and abundance denominated size spectrum, is a concept initially developed by Sheldon et al. (1972), in which the authors assessed the distribution from bacteria biomass to whales and hypothesized that aquatic biomass is uniformly distributed among logarithmic class sizes. This approach provide important information about trophic ecology (Petchey and Belgrano, 2010; Gaedke, 1993; Gaedke, 1992) and environmental disturbances (Gamble et al., 2006; Shin et al., 2005), besides that, it has contributed for a better understanding of aquatic communities function and the regulatory processes of energy flow in these environments (Zhou et al., 2010; Layman et al., 2005; Thiebaut and Dickie, 1992; Rodríguez and Mullin, 1986).

As approaches based on body size are independent of the taxon, individuals with the same body size are considered energetically equivalent, and used as a reference for simplification of food webs. Size-based indicators has been used for studies of fish assemblages to assess impacts caused by fishing activity on stock size, marine food web dynamics (Gamble et al., 2006; Shin et al., 2005; Rice and Gislason, 1996) and fish communities in reservoirs (Kantoussan et al., 2009). Thus, the use of spectrum size models are relevant to fisheries science in the context of the ecosystem approach to fisheries management. The theory behind of spectrum size includes several mathematical derivations and estimation methods that are described in a number of detailed publications and their appendices (Andersen et al., 2015; Sprules and Barth, 2015; Reuman et al., 2008, White et al., 2007; Andersen and Beyer, 2006; Brown and Gillooly, 2003).

Freshwater ecosystem are vulnerable to anthropogenic disturbance due to landscape

*Corresponding author: Louise Cristina Gomes
E-mail address: louise-cristina@hotmail.com




Figure 1. Conceptual model about differences in water level fluctuations.




Figure 2. Location of the six Neotropical reservoirs sampled from January 2005 to December 2007. SO=Salto Osório, SS=Salto Santiago, SE=Segredo, FA=Foz do Areia, VO=Vossoroica and SM=Salto do Meio.

provided in Figure 1.

In this context, the aim of this study was to assess the size spectrum distributions in fish in Neotropical reservoirs, specifically to answer the following question: is there difference in fish size spectrum in reservoirs with different operating regimes? For this, we tested the hypothesis that the run of river reservoirs present higher proportions of large individuals than storage system.

Materials and Methods

Study Area: This study was conducted in six reservoirs located in the state of Paraná, southern Brazil. Four reservoirs were analyzed in the Paraná river basin, belonging to the following sub-basin: Iguaçú River and two reservoirs belonging to the Coastal basin in São Jorge River (Fig. 2). The Iguaçú River present an area of approximately 72.000 km² and the high unevenness of the basin constituted a great attraction for the hydroelectric exploitation, resulting in a cascade of five reservoirs that alters their physical, chemical and biological attributes (Baumgartner et al., 2012). The fish fauna of Iguaçú River present a high degree of endemism due to geographic isolation caused by Iguazu Falls.

The Coastal basin is formed by rivers draining directly into the Atlantic Ocean and present a total area of 5.630,8 Km². They are rivers with great slope, rapids and greater speed of current, as a result of this, several reservoirs were designed for the production of

influences on habitat and fishes (Van der Lee and Koops, 2015). Reservoirs are historically recent environments and considered as complex ecosystems, with characteristics distinct from those of natural rivers and lakes (Fernando and Holcik, 1991). The main Neotropical water basins are altered by dams, which were built for various purposes such as power generation, irrigation and urban water supply (Stenberg, 2006; Agostinho et al., 1999). These systems are exposed to natural and artificial forces that determine dynamic characteristics; in addition to climatological and hydrological events, reservoir operational regime, manner of construction, and use also interfere with the system dynamics and spatial and temporal organization (Straskraba et al., 1993). Further, reservoir operation significantly impacts river hydrology, changing the magnitude, frequency, duration, and timing of the flow regime (Suen, 2011). In this context, the reservoir operation mode can be classified according to the control mechanism for everted and turbined water. Run-of-river systems whose characteristics are more similar to natural lakes and storage systems in which water stock is kept high mainly in high rainfall periods, enables the maintenance of activity in dry periods. Therefore, the storage system tend to cause more disturb than run-of-river system due to higher water level fluctuations (Poff and Hart, 2002). These characteristics promote fragmentation and transformation of the river, as well as changes in species diversity, trophic structure and community composition (Poff et al., 1997; Ward and Stanford, 1995). The conceptual model about differences in water level fluctuations in reservoir are

Table 1. Physical factors of the analyzed reservoirs. Locations were paired as follows: Pair 1: FA and SE; Pair 2: SS and SO; Pair 3: VO and SM (FA=Foz do Areia, SE=Segredo, SS=Salto Santiago, SO=Salto Osório, VO=Vossoroca, SM=Salto do Meio and *=Data not found).

Physical factors	FA	SE	SS	SO	VO	SM
Operation regime	Storage	Run-of-river	Storage	Run-of-river	Storage	Run-of-river
Sub-basin	Iguaçu	Iguaçu	Iguaçu	Iguaçu	Coastal	Coastal
Fill year	1980	1992	1980	1975	1931	1931
Area (km ²)	139	80.6	208	55	5.1	0.1
Sampling	Quarterly	Quarterly	Bimonthly	Bimonthly	Quarterly	Quarterly
Average depth (m)	40.0	36.6	35	25.5	4.0	*
Altitude	742	607	506	397	814	722
Residence time (days)	105	47	50.8	16	110	*

energy (Rodrigues et al., 2005). These reservoirs present wide heterogeneity of morphometric and limnological characteristics. For analysis criteria, the reservoirs were grouped in pairs with one run-of river and one storage system each, according to the sub-basin in which they belong, area and proximity to the operating system (Table 1).

Sampling: Samplings were carried out between January 2005 and December 2007 (Table 1). Data collection took place in non-adjacent bank regions at points located upstream the dam in six reservoirs using mesh gill nets between 2.4 cm and 16 cm. Nets were exposed for 24 hours and collections took place at 8, 16 and 22 hours. Fish were anesthetized with benzocaine hydrochloride (250 mg/L), as required by Resolution No. 714/CFMV as of July 20, 2002, which regulates procedures and methods of euthanasia in animals (CFMV, 2002). After anesthesia fish were fixed in 10% formalin and 70% alcohol (Shibatta and Cheida, 2003) and assigned ID based on location, shift, and sampling month. Specimens were subsequently taken to the laboratory for analysis.

Analysis were performed at the Laboratory of Ichthyology at the State University of West Paraná, Toledo campus. Specimens were classified according to Graça and Pavanelli (2007), Britski et al. (1999) and Eschmeyer (1990). Weights and length were obtained for all specimens. The list of species and their respective abundances are provided in Table 2.

Data Analysis: The richness of species was assessed using the number of individual captured (See supplementary material). To represent the size spectrum, we used the biomass of all individuals captured in each reservoir. The spectrum calculation

was performed using the Pareto type I continuous distribution model for each reservoir. According to this model, the probability of finding individuals that are older or of a specific size decays on a logarithmic scale as size increases. This has been used for the analysis of biomass frequency distributions (White et al., 2008; Vidondo et al., 1997). The derivation of the Pareto type I model is described by the following equation:

$$\log_2(\text{Prob}(s \geq S)) = c * (\log_2(K) - \log_2(S))$$

Where $\text{Prob}(s \geq S)$ is the fraction of individuals over a given size (s) taken at random (S). The constant c is the slope of the size spectrum, representing the frequency distribution of fish body size. Thus, when $c=-1$, the frequency distribution is uniform between individual fish of different sizes; if $c<-1$, the frequency of smaller individuals is higher than large individuals; if $c>-1$, the frequency of large individuals is greater than smaller individuals. The Pareto model was fitted by non-linear regression using the least squares method. The possible effects of reservoir operation regime on the biomass spectra of fish assemblages were assessed using paired t test, both using the software Statistica 7.1® (Stat Soft, 2005). Results were considered significant at $P<0.05$.

Results

The slopes of the biomass spectra of fishes observed in each reservoir indicated that reservoirs Foz do Areia (storage), Salto Santiago (storage), Salto Osório (run of river) and Vossoroca (storage) tended to have higher concentrations of small individuals ($c<-1$), while Segredo (run of river) show equal distribution between large and small ($c \sim -1$). Salto do Meio (run of

Table 2. Species collected from January 2005 to December 2007 at six neotropical reservoirs and their respective abundances (FA=Foz do Areia, SE=Segredo, SS=Salto Santiago, SO=Salto Osório, VO=Vossoroica and SM=Salto do Meio).

TAXON	Common Name	FA	SE	SS	SO	VO	SM
Atheriniformes							
<i>Odontesthes bonariensis</i> (Valenciennes, 1835)	“Peixe-rei”	-	174	282	30	-	-
Characiformes							
<i>Apareiodon vittatus</i> (Garavello, 1977)	“Canivete”	67	716	88	294	-	-
<i>Astyanax altiparanae</i> Garutti & Britski, 2000	“Tambiú”	33	28	331	38	61	61
<i>Astyanax bifasciatus</i> Garavello & Sampaio, 2010	“Lambari-do-rabo-vermelho”	13902	2100	5270	1812	-	-
<i>Astyanax dissimilis</i> Garavello & Sampaio, 2010	“Lambari”	647	24	12	7	-	-
<i>Astyanax gymnodontus</i> (Eigenmann, 1911)	“Lambarizão”	184	256	1047	265	-	-
<i>Astyanax janeiroensis</i> (Eigenmann, 1908)	“Lambari-do-rio”	-	-	-	-	-	4
<i>Astyanax longirhinus</i> Garavello & Sampaio, 2010	“Lambari”	4	2	-	-	-	-
<i>Astyanax minor</i> Garavello & Sampaio, 2010	“Lambari-do-rabo-amarelo”	1651	1363	4100	292	-	-
<i>Astyanax serratus</i> Garavello & Sampaio, 2010	“Lambari”	7	-	-	-	-	-
<i>Bryconamericus ikae</i> Casciotta, Almirón & Azpelicueta, 2004	“Lambarizinho”	206	529	18	10	-	-
<i>Bryconamericus pyahu</i> Azpelicueta, Casciotta & Almirón, 2003	“Lambarizinho”	2	-	-	-	-	-
<i>Characidium</i> sp.	“Charutinho”	4	-	-	-	-	-
<i>Cyphocarax santacatarinae</i> (Fernández-Yépez, 1948)	“Saguaru”	4	-	-	-	-	-
<i>Deuterodon iguape</i> Eigenmann, 1907	“Lambari”	-	-	-	-	2265	1475
<i>Deuterodon</i> sp. A	“Lambari”	-	-	-	-	1262	384
<i>Deuterodon</i> sp. D	“Lambari”	-	-	-	-	12413	460
<i>Hoplias malabaricus</i> (Bloch, 1794)	“Traíra”	180	24	-	-	213	165
<i>Hoplias aff. malabaricus</i> (Bloch, 1794)	“Traíra”	-	-	38	18	-	-
<i>Leporinus elongatus</i> (Valenciennes, 1850)	“Piapara”	3	-	-	-	-	-
<i>Leporinus friderici</i> (Bloch, 1794)	“Piau”	2	-	-	-	-	-
<i>Leporinus macrocephalus</i> (Garavello & Britski, 1988)	“Piauçú”	-	-	-	-	-	-
<i>Leporinus obtusidens</i> (Valenciennes, 1837)	“Piau”	4	-	-	-	-	-
<i>Leporinus octofasciatus</i> (Steindachner, 1915)	“Piau-listrado”	2	-	-	-	-	-
<i>Leporinus</i> sp.	“Piau”	-	-	-	1	-	-
<i>Oligosarcus longirostris</i> Menezes & Géry, 1983	“Saicanga”	1372	262	256	306	-	-
<i>Prochilodus lineatus</i> (Valenciennes, 1837)	“Corimba”	116	-	-	-	-	-
<i>Salminus brasiliensis</i> (Cuvier, 1816)	“Dourado”	1	-	-	-	-	-
<i>Serrapinnus</i> sp.	“Piabinha”	-	-	-	-	-	1
Cypriniformes							
<i>Cyprinus carpio</i> (Linnaeus, 1758)	“Carpa-comum”	5	2	2	-	1	-
<i>Hypophthalmichthys nobilis</i> (Richardson, 1845)	“Carpa-cabeça-grande”	1	-	-	-	-	-
Gymnotiformes							
<i>Gymnotus inaequilabiatus</i> (Valenciennes, 1839)	“Morenita”	6	2	1	-	-	-
<i>Gymnotus sylvius</i> (Albert & Fernandes-Matioli, 1999)	“Morenita”	2	-	-	-	-	-
Perciformes							
<i>Australoheros kaaygua</i> (Casciotta, Almirón & Gómez, 2006)	-	3	-	-	-	-	-
<i>Australoheros</i> sp.	“Acará”	-	-	-	-	-	2
<i>Crenicichla iguassuensis</i> (Haseman, 1911)	“Joaninha”	69	122	27	30	-	-
<i>Crenicichla</i> sp.	“Joaninha”	17	15	6	14	-	-
<i>Geophagus brasiliensis</i> (Quoy & Gaimard, 1824)	“Cará”	227	168	50	15	842	260
<i>Micropterus salmoides</i> (Lacepède, 1802)	“Boca-grande”	-	-	-	-	77	8
<i>Oreochromis niloticus</i> (Linnaeus, 1758)	“Tilápia-do-nilo”	1	3	1	-	-	-
<i>Tilapia rendalli</i> (Boulenger, 1897)	“Tilápia”	17	24	4	3	136	1
Siluriformes							
<i>Ancistrus</i> sp.	“Cascudo-roseta”	4	3	-	-	-	-
<i>Corydoras ehrhardti</i> (Steindachner, 1910)	“Limpa-fundo”	-	-	-	-	53	26
<i>Corydoras paleatus</i> (Jenyns, 1842)	“Limpa-fundo”	889	1072	540	133	1	-
<i>Glanidium ribeiroi</i> (Haseman, 1911)	“Bocado”	26	93	4	18	-	-
<i>Hypostomus ancistroides</i> (Ihering, 1911)	“Cascudo-pintado”	-	-	-	-	2	2
<i>Hypostomus aspilogaster</i> (Cope, 1894)	“Cascudo”	-	-	-	-	1	-
<i>Hypostomus boulengeri</i> (Eigenmann & Kennedy, 1903)	“Cascudo”	-	-	-	-	-	1
<i>Hypostomus commersoni</i> (Valenciennes, 1836)	“Cascudo”	80	223	13	2	88	30
<i>Hypostomus derbyi</i> (Haseman, 1911)	“Cascudo”	36	73	4	9	-	-
<i>Hypostomus myersi</i> (Gosline, 1947)	“Cascudo”	7	6	1	2	-	-
<i>Hypostomus</i> sp. 1	“Cascudo”	-	-	-	-	-	3
<i>Hypostomus</i> sp. 2	“Cascudo”	-	-	-	-	-	1
<i>Isbrueckerichthys</i> sp.	“Cascudinho”	-	-	-	-	-	1
<i>Pimelodus britskii</i> Garavello & Shibatta, 2007	“Mandi-pintado”	1024	1397	-	-	-	-
<i>Pimelodus ortmanni</i> Haseman, 1911	“Mandi-pintado”	44	181	459	685	-	-
<i>Pseudoplatystoma corruscans</i> (Spix & Agassiz, 1829)	“Pintado”	-	-	-	-	-	1
<i>Rhamdia branneri</i> Haseman, 1911	“Bagre, jundiá”	3	5	1	7	-	-
<i>Rhamdia quelen</i> (Quoy & Gaimard, 1824)	“Bagre-amarelo”	-	-	-	-	30	42

Table 2. Continued.

TAXON	Common Name	FA	SE	SS	SO	VO	SM
<i>Rhamdia voulezi</i> Haseman, 1911	“Bagre, jundiá”	34	18	4	2	-	-
<i>Rineloricaria</i> sp.	“Cascudo-espada”	-	-	-	-	237	259
<i>Steindachmeridion melanodermatum</i> (Garavello, 2005)	“Surubim”	-	1	-	-	-	-
<i>Tatia jaracatia</i> Pavanelli & Bifi, 2009	“Bagre-sapo”	-	-	-	1	-	-
Total Richness		40	29	26	25	16	21




Figure 3. Size spectra of fish assemblages in the studied reservoirs, adjusted using the Pareto type I model.

river), showed a higher proportion of large individuals ($c > -1$) (Fig. 3). The reservoirs operated under run-of-river regime had significantly higher values than did those operated under storage regimes (paired t test: $t=3.777$, $P=0.032$).

Discussion

The results indicated differences in size spectra of fish assemblages among the sampled reservoirs. Here,

abiotic events that could to determine some relationships between body size distributions and the environment were not evaluate. However, the physical characteristics differences were considered in each reservoir and how these aspects can change the communities and to provide some information about ecological processes which regulate these environments.

Fluctuations in reservoir water level due to

accumulation operating regime cause flooding of bank areas considered critical for fish feeding (Hahn et al., 1998) and may act as a disturbance factor, altering the structure and stability of fish assemblages (Tundisi et al., 2003). The increase in nutrients from the decomposition of flooded vegetation provides an increase in primary productivity, and a consequent increase in smaller sized individuals. According to Agostinho et al. (2001), flooding may favor juveniles and may thus influence the size distribution patterns. The success of higher proportions of small individuals in accumulation regime reservoirs possibly is likely due to fluctuations in water level that provide better conditions for some species. For example, the genus *Astyanax* have adhesive eggs (Sato et al., 2006) and fractional spawning (Agostinho, 1999), facilitating the use of margin areas as sites for spawning and juvenile shelter when the reservoir contains abundant water (i.e., high water level). When the opposite occurs, that is, lower water level, the edges and vegetation are exposed and there is a tendency toward increased turbidity, which reduces the visibility for piscivorous fish and consequently, decreases predation rate (Robertis et al., 2003).

Important ecological interactions and physiological processes in aquatic ecosystems are dependent on body size (e.g., mortality, transfer efficiencies, metabolism and growth) (Kerr and Dick, 2001). Consequently, researchers have been incorporating biomass spectrum theory into models designed to estimate size-dependent ecological processes. For example, Benejam et al. (2015) examined the effect of land use on the size structure of fish communities in subtropical streams. Van der Lee and Koops (2015) investigated correlations between body size of fish with habitat loss. Jennings et al. (2002) quantified trophic transfer efficiencies using a combination of biomass spectrum data and production-body size and trophic level-body size relationships. Furthermore, the current understanding of regulatory processes and trophic dynamics of aquatic ecosystems has been focused on ecological studies (Jansson et al., 2007; Shin and Cury, 2004) and approaches using body size have been used to primarily to provide information

concerning conservation of aquatic environments (Petchey and Belgrano, 2010), trophic structure, and biotic interactions in fish assemblages (Emmrich et al., 2011; Jennings et al., 2001). In this sense, body size is crucial for predicting responses to energy flow, species diversity and population densities (Layman et al., 2005).

From an ecological point of view, trophic dynamics are highly sensitive to environmental disturbances resulting from fluctuations in water level (Tundisi et al., 2003), and these changes are key components for the reorganization of fish assemblages for which feeding grounds and reproduction are affected (Agostinho et al., 2007). Although reservoirs are dominant components of hydrological landscapes, studies of the effects of operating regime on biomass spectra are scarce, making it difficult to make inferences regarding size distribution and energy flow in communities. Besides that, others factors can be associate to the differences in sizes distributions. The size of individual in ecological community is affected by many kinds of processes, for example, environmental changes, species invasion and management (Petchey and Belgrano, 2010). According to metabolic theory of ecology, a greater part of variation among organisms, including their ecological roles and life history, is limited by their body sizes, chemical composition and operating temperatures (Brown et al., 2004).

This study has elucidated some impacts of reservoir operating regime on the size distributions and this suggests that energy transfer occurs more efficiently through fish fauna in reservoirs with lower fluctuations in water level, such as run-of-river regimes. However, more researches is needed to unreavel relationships between body size distributions and the ecological processes regulating the energy flow in reservoirs. In addition, researches which consider abiotic factors and others relevant aspects are need considering the acceleration of transformations at the landscape caused by impoundments. Finally, it is important to incorporate management plans that take dam operating mode into account so that conservation of aquatic fauna, especially fish, is more

effective.

Acknowledgments

We thank the Research Group on Fisheries, Resources and Limnology at the State University of West Paraná for logistical support and providing data for analysis. We thank CAPES for financial support.

References

- Agostinho A.A., Miranda L.E., Bini L.M., Gomes L.C., Thomaz S.M., Suzuki H.I. (1999). Patterns of colonization in Neotropical reservoirs, and prognoses on aging. In: J.G. Tundisi, M. Straskraba (Eds.). *Theoretical reservoir ecology and its applications*, International Institute of Ecology, Brazilian Academy of Science and Backhuys Publishers. São Carlos, São Paulo. pp: 226-265.
- Agostinho A.A., Gomes L.C., Zalewski M. (2001). The importance of floodplains for the dynamics of fish communities of the upper river Paraná. *Ecohydrology and Hydrobiology*, 1: 209-217.
- Agostinho A.A., Gomes L.C., Pelicice F.M. (2007). *Ecologia e manejo de recursos pesqueiros em reservatórios do Brasil*. Maringá, Eduem. 501 p.
- Andersen K.H., Beyer J.E. (2006). Asymptotic size determines species abundance in the marine size spectrum. *American Naturalist*, 168: 54-61.
- Andersen K.H., Jacobsen N.S., Farnsworth K.D. (2015). The theoretical foundations for size spectrum models of fish communities. *Canadian Journal of Fisheries and Aquatic Sciences*, 73: 575-588.
- Baumgartner G., Pavanelli C.S., Baumgartner D., Bifi A.G., Debona T., Frana V.A. (2012). *Peixes do baixo rio Iguaçu*. Maringá, Eduem. 203 p.
- Benejam L., Mello F.T., Meerhoff M., Loureiro M., Jeppesen E., Brucet S. (2015). Assessing effects of change inland use on size-related variables of fish in subtropical streams. *Canadian Journal of Fisheries and Aquatic Sciences*, 73: 547-556.
- Britski H.A., Silimon K.Z.S., Lopes B.S. (1999). *Peixes do Pantanal: manual de identificação*. Corumbá, Embrapa. 227 p.
- Brown J.H., Gillooly J.F. (2003). Ecological food webs: high-quality data facilitate theoretical unification. *Proceedings of the National Academy of Science of the USA*, 100: 1467-1468.
- Brown J.H., Gillooly J.F., Allen A.P., Savage V.M., West G.B. (2004). Toward a metabolic theory of ecology. *Ecology*, 85: 1771-1789.
- Conselho Federal de Medicina Veterinária (CFMV). (2002). Resolução N.714 Dispõe sobre procedimentos e métodos de eutanásia em animais, e dá outras providências. Available in: <http://portal.cfmv.gov.br/portal/lei/index/id/327>. Access in: 01/11/2014.
- Elton C. (1927). *Animal ecology*. Macmillan, New York. 260 p.
- Emmrich M., Brucet S., Ritterbusch D., Mehner D. (2011). Size spectra of lake fish assemblage: responses along gradients of general environmental factors and intensity of lake-use. *Freshwater Biology*, 56: 2316-2333.
- Eschmeyer W.N. (1990). *Catalog of the genera of recent fishes*. San Francisco, Academic of Sciences. 697 p.
- Fernando C.H., Holcik J. (1991). Fish in reservoir. *Internationale Revue der gesamten Hydrobiologie und Hydrographie*, 76: 149-167.
- Gaedke U. (1992). The size distribution of plankton biomass in a large lake and its seasonal variability. *Limnology and Oceanography*, 37: 1202-1220.
- Gaedke U. (1993). Ecosystem analysis based on biomass size distributions: a case study of the plankton community in a large lake. *Limnology and Oceanography*, 38: 112-127.
- Gamble A.E., Lloyd R., Aiken J., Johannsson O.E., Mills E.L. (2006). Using zooplankton biomass size spectra to assess ecological change in a well-studied freshwater lake ecosystem. *Oneida Lake, New York. Canadian Journal of Fisheries and Aquatic Sciences*, 63: 2687-2699.
- Giacomini H.C., Shuter B.J., Baum J.K. (2016). Size-based approaches to aquatic ecosystems and fisheries science: a symposium in honour of Rob Peters. *Canadian Journal of Fisheries and Aquatic Sciences*, 73: 471-476.
- Graça W.J., Pavanelli C.S. (2007). *Peixes da planície de inundação do alto rio Paraná e áreas adjacentes*. Maringá, Eduem. 241 p.
- Hahn N.S., Agostinho A.A., Gomes L.C., Bini L.M. (1998). Estrutura trófica da ictiofauna do reservatório de Itaipu (Paraná-Brasil) nos primeiros anos de sua formação. *Interciencia*, 23: 299-305.
- Jennings S., Pinnegar J.K., Polunin N.V.C., Boon T.W. (2001). Weak cross-species relationships between body size and trophic level belie powerful size-based trophic structuring in fish communities. *Journal of Animal Ecology*, 70: 934-944.
- Kantoussan J., Ecoutin J.M., Fontenelle G., Thiaw O.T.,

- Morais L.T., Laë R. (2009). The relevance of size parameters as indicators of fishery exploitation in two West African reservoirs. *Aquatic Ecology*, 43: 1167-1178.
- Kerr S.R., Dickie L.M. (2001). *The biomass spectrum: a predator-prey theory of aquatic production*. Columbia University Press. 352 p.
- Layman C.A., Winemiller K.O., Arrington D.A., Jepsen D.B. (2005). Body size and trophic position in a diverse tropical food web. *Ecology*, 86: 2530-2535.
- Petchey O.L., Belgrano A. (2010). Body-size distributions and size spectra: universal indicators of ecological status? *Biology Letters*, 1-5.
- Poff N.L., Allan J.D., Bain M.B., Karr J.R., Prestegard K.L., Richter B.D., Sparks R.E., Stromberg J.C. (1997). The natural flow regime: a paradigm for river conservation and restoration. *BioScience*, 47: 769-784.
- Poff N.L., Hart D.D. (2002). How dams vary and why it matters for the emerging science of dam removal. *BioScience*, 52: 659-668.
- Reuman D.C., Mulder C., Raffaelli D., Cohen J. (2008). Three allometric relations of population density to body mass: theoretical integration and empirical tests in 149 food webs. *Ecology Letters*, 11: 1216-1228.
- Rice J., Gislason H. (1996). Patterns of change in the size spectra of numbers and diversity of the North Sea fish assemblage, as reflected in surveys and models. *ICES Journal of Marine Science*, 53: 1214-1225.
- Robertis A., Ryer C., Veloza A., Brodeur R. (2003). Differential effects of turbidity on prey consumption of piscivorous and planktivorous fish. *Canadian Journal of Fisheries and Aquatic Sciences*, 60: 1517-1526.
- Rodrigues L., Thomaz S. M., Agostinho A. A., Gomes L.C. (2005). *Biocenoses em reservatórios: Padrões espaciais e temporais*. São Carlos, RiMa. 333 p.
- Rodríguez J., Mullin M. (1986). Relations between biomass and body-weight of plankton in a steady state oceanic system. *Limnology and Oceanography*, 21: 361-370.
- Sato Y., Sampaio E.V., Fenerich-Verani N., Verani J.R. (2006). *Biologia reprodutiva e reprodução induzida de duas espécies de Characidae (Osteichthyes, Characiformes) da bacia do São Francisco, Minas Gerais, Brasil*. *Revista Brasileira de Zoologia*, 23: 267-273.
- Sheldon R.W., Prakash A., Suctcliffe Jr W.F. (1972). The size distribution of particles in the ocean. *Limnology and Oceanography*, 17: 323-340.
- Shibatta O.A., Cheida C.C. (2003). *Composição em tamanho dos peixes (Actinopterygii, Teleostei) de ribeirões da bacia do rio Tibagi, Paraná, Brasil*. *Revista Brasileira de Zoologia*, 20: 469-473.
- Shin Y.J., Rochet M.J., Jennings S., Field J.G., Gislason H. (2005). Using size-indicators to evaluate the ecosystem effects of fishing. *ICES Journal of Marine Science*, 62: 384-396.
- Shin Y.J., Cury P. (2004). Using an individual-based model of fish assemblages to study the response of size spectra to changes in fishing. *Canadian Journal of Fisheries and Aquatic Sciences*, 61: 414-431.
- Sprules W.G., Barth L.E. (2015). Surfing the biomass size spectrum: some remarks on history theory, and application. *Canadian Journal of Fisheries and Aquatic Sciences*, 73: 477-495.
- Statsoft Inc. (2005) *STATISTICA (data analysis software system)*, version 7.1 www.statsoft.com
- Sternberg R. (2006). Damming the river: a changing perspective on altering nature. *Renewable and Sustainable Energy Reviews*, 10: 165-197.
- Straskraba M., Tundisi J.G., Duncan A. (1993). *Comparative reservoir limnology and water quality management*. Kluwer Academic Publisher, Netherlands. 294 p.
- Suen J.P. (2011). Determining the ecological flow regime for existing reservoir operation. *Water Resources Management*, 25: 817-835.
- Thiebaut M.L., Dickie L.M. (1992). Models of aquatic biomass size spectra and the common structure of their solutions. *Journal of Theoretical Biology*, 159: 147-161.
- Tundisi J.G., Matsumura-Tundisi T., Rodríguez L.S. (2003). *Gerenciamento e recuperação das bacias hidrográficas dos rios Itaqueri e do Lobo e da Represa Carlos Botelho (Lobo-Broa)*. São Carlos, IIE. 54 p.
- Van der Lee A.S., Koops M.A. (2015). Are small fishes more sensitive to habitat loss? A generic size-based model. *Canadian Journal of Fisheries and Aquatic Sciences*, 73: 716-726.
- Vidondo B., Prairie Y.T., Blanco J.M., Duarte C.M. (1997). Some aspects of the analysis of size spectra in aquatic ecology. *Limnology and Oceanography*, 42: 184-192.
- Ward J.V., Stanford J.A. (1995). The serial discontinuity concept: extending the model to floodplain rivers. *Researches and Management*, 10: 159-168.
- Wheeland L.J., Rose G.A. (2016). Acoustic measures of lake community size spectra. *Canadian Journal of Fisheries and Aquatic Sciences*, 73: 557-564.

- White E.P., Ernest S.K.M., Kerkhoff A.J., Enquist B.J. (2007). Relationships between body size and abundance in ecology. *Trends in Ecology and Evolution*, 22: 323-330.
- White E.P., Enquist B.J., Green J.L. (2008). On estimating the exponent of power-law frequency distributions. *Ecology*, 89: 905-912.
- Zhou M., Carlotti F., Zhu Y. (2010). A size-spectrum zooplankton closure model for ecosystem modelling. *Journal of Plankton Research*, 32: 1147-116.

Archive of SID